arXiv:1702.07062v1 [math.PR] 23 Feb 2017

Stochastic complex Ginzburg-Landau equation with space-time white noise

Masato Hoshino, Yuzuru Inahama and Nobuaki Naganuma

Abstract

We study the stochastic cubic complex Ginzburg-Landau equation with complex-valued space-time white noise on the three dimensional torus. This nonlinear equation is so singular that it can only be understood in a renormalized sense. In the first half of this paper we prove local well-posedness of this equation in the framework of regularity structure theory. In the latter half we prove local well-posedness in the framework of paracontrolled distribution theory.

Keywords: Stochastic partial differential equation, Complex Ginzburg-Landau equation, Regularity structure, Paracontrolled distribution, Renormalization.

MSC2010: 60H15, 82C28.

Contents

1 Introduction **2**

2 CGL by the theory of regularity structures **5**

2.1 Results on regularity structures 6

2.2 Regularity structures associated with CGL and admissible models 10

2.3 Abstract solution map 14

2.4 Renormalization 16

2.5 Convergence of Gaussian models 21

3 Proof of convergence of renormalized models **21**

3.1 Wiener chaos decomposition 22

3.2 Estimates of singularity of kernels 23

3.3 Proof of L^2-estimates (3.1) 24

3.3.1 $\Xi, \mathbf{I}, \mathbb{I}, \mathbf{V}, \mathbb{V}, \mathbb{V}$ 24

3.3.2 $X_i\mathbf{V}, X_i\mathbb{V}, \mathbf{V}, \mathbf{V}, \mathbf{V}, \mathbf{V}$ 26

3.3.3 . . . 27
3.4 Behaviors of renormalization constants . . . 30

4 CGL by the theory of paracontrolled distributions 32
4.1 Besov-Hölder spaces and paradifferential calculus . . . 34
4.1.1 Besov spaces . . . 34
4.1.2 Paraproducts and Commutator estimates . . . 35
4.1.3 Regularity of $\mathcal{C}^\alpha$-valued functions . . . 36
4.2 Definitions of driving vectors and solutions . . . 37
4.3 Properties of the integration map . . . 42
4.3.1 Properties of $\mathcal{M}^1$ . . . 42
4.3.2 Properties of $\mathcal{M}^2$ . . . 45
4.4 Local existence and uniqueness . . . 52
4.5 Renormalized equation . . . 56

5 Proof of convergence of driving vectors 60
5.1 Convergence criteria . . . 61
5.1.1 $\mathcal{C}^\alpha$-valued random variables . . . 62
5.1.2 Good kernels . . . 62
5.2 Definitions of driving vectors . . . 65
5.2.1 Ornstein-Uhlenbeck like process and its approximations . . . 65
5.2.2 Definition of driving vectors . . . 65
5.2.3 Itô-Wiener integral expressions of driving vectors . . . 66
5.3 Proof of convergence of driving vectors . . . 71
5.3.1 Useful estimates . . . 71
5.3.2 Lower order terms . . . 72
5.3.3 Higher order terms . . . 73
5.4 Properties of renormalization constants . . . 81
5.4.1 Expression of renormalization constants . . . 81
5.4.2 Divergence rate of renormalization constants . . . 82

A Complex multiple Itô-Wiener integral 84

1 Introduction

The cubic complex Ginzburg-Landau (CGL) equation is one of the most important nonlinear partial differential equations (PDEs) in applied mathematics and physics. It describes various physical phenomena such as nonlinear waves,

second-order phase transition, superconductivity, superfluidity among others. See [AK02] for example.

There are also many papers on its stochastic version, the CGL with a noise term ([BS04a, BS04b, KS04, Oda06, PG11, Yan04] to name but a few). In these preceding works, however, the noise is either non-white or multiplicative. Except when the space dimension $d = 1$ in [Hai02], the stochastic cubic CGL with *additive* space-time white noise has not been solved.

The difficulty in the case $d \geq 2$ is as follows. Since space-time white noise is so rough, a solution $u_t(x) = u(t, x)$ would be a Schwartz distribution in x, not a function, even if it existed. Consequently, the cubic nonlinear term $|u_t|^2 u_t$ does not make sense in the usual way. For this reason, well-definedness of the equation itself was unclear and the cubic CGL with space-time white noise was considered too singular when $d \geq 2$.

However, two new theories emerged recently, which can deal with quite singular stochastic PDEs of this kind. One is regularity structure theory [Hai14] and the other is paracontrolled distribution theory [GIP15]. They are both descendants of rough path theory and their deterministic part looks somewhat similar to the counterpart in rough path theory at least in spirit. However, their probabilistic part is more complicated than the counterpart in rough path theory since non-trivial renormalization of the noise has to be done. (There is another theory based on the theory of renormalization groups [Kup16] which will not be discussed in this paper, however.)

Although they are clearly different theories, examples of stochastic PDEs they can deal with are very similar. A partial list of singular stochastic PDEs which have been solved (locally in time) by these theories is as follows: Parabolic Anderson Model ($d = 2, 3$) [GIP15, Hai14, BBF15], KPZ equation and its variants ($d = 1$) [FH14, GP17, Hos16a, FH16], the dynamic Φ^4_3-model ($d = 3$) [Hai14, CC13], Navier-Stokes equation with space-time white noise ($d = 3$) [ZZ15], FitzHugh-Nagumo equation with space-time white noise ($d = 3$) [BK16].

The main objective of this paper is to prove local well-posedness of the stochastic cubic complex Ginzburg-Landau equation on the three-dimensional torus $\mathbf{T}^3 = (\mathbf{R}/\mathbf{Z})^3$ of the following form by using these two theories:

$$\partial_t u = (\mathsf{i} + \mu)\triangle u + \nu(1 - |u|^2)u + \xi, \qquad t > 0, \quad x \in \mathbf{T}^3. \tag{1.1}$$

Here, $\mathsf{i} = \sqrt{-1}$, $\mu > 0$, $\nu \in \mathbf{C}$ are constants and ξ is *complex-valued* space-time white noise, that is, a centered complex Gaussian random field with covariance

$$\boldsymbol{E}[\xi(t, x)\xi(s, y)] = 0, \qquad \boldsymbol{E}[\xi(t, x)\overline{\xi(s, y)}] = \delta(t - s)\delta(x - y),$$

where δ denotes the Dirac delta function.

We replace ξ by a smeared noise ξ^ϵ with a parameter $0 < \epsilon < 1$ so that $\xi^\epsilon \to \xi$ as $\epsilon \downarrow 0$ in an appropriate topology and consider a renormalized equation

$$\partial_t u^\epsilon = (\mathrm{i} + \mu)\triangle u^\epsilon + \nu(1 - |u^\epsilon|^2)u^\epsilon + \nu C^\epsilon u^\epsilon + \xi^\epsilon, \qquad t > 0, \quad x \in \mathbf{T}^3, \tag{1.2}$$

where C^ϵ is a suitably chosen complex constant (specified later) which diverges as $\epsilon \downarrow 0$. We show that the solution to (1.2) converges to some process in an appropriate topology. To this end, we use the theory of regularity structure by Hairer [Hai14] and the theory of paracontrolled distributions by Gubinelli-Imkeller-Perkowski [GIP15]. The smeared noises ξ^ϵ and the constants C^ϵ in the results by the regularity structure and the paracontrolled distributions are different. Consequently, we do not know at the moment whether the solutions obtained by these two theories coincide, although it is likely.

We now make a comment on the space dimension. When $d \geq 4$, CGL (1.1) is not subcritical in the sense of [Hai14] and therefore the equation cannot be solved (or does not even make sense) by any existing method. Though we do not give a proof in this paper, we believe that the case $d = 2$ is actually much easier than our case $d = 3$.

This paper is organized as follows. In Sections 2 and 3, following [Hai14], we apply the theory of regularity structures to the stochastic CGL (1.1). At the beginning of Section 2 we first present our main result (Theorem 2.1) in a precise form. Then we construct a regularity structure for (1.1) and prove local-wellposedness of (1.1) in a deterministic way. Section 3 is devoted to the probabilistic step, in particular, the renormalization procedure.

In Sections 4 and 5, we apply the paracontrolled calculus to (1.1). In Section 4, we precisely present our main result (Theorem 4.1) and deterministically solve (1.1) locally in time in a similar way to [MW16]. We prove the probabilistic part in Section 5 using a new method developped in [GP17].

Section A is an appendix, in which we recall the definition of complex multiple Itô-Wiener integrals. The product formula for them is frequently used in Sections 3 and 5.

Notations: We use the following notations: For two functions f and g, we write $f \lesssim g$ if there exists a positive constant C such that $f(x) \leq Cg(x)$ for any x. We write $f(x) \approx g(x)$ if both $f(x) \lesssim g(x)$ and $g(x) \lesssim f(x)$ hold. To indicate the argument x of a function f, we use both symbols $f(x)$ and f_x.

2 CGL by the theory of regularity structures

In this and the next sections, we study CGL equation by the theory of regularity structures. We begin by presenting the main result in Theorem 2.1 below.

We replace ξ by space-time smeared noise $\xi^\epsilon = \xi * \rho^\epsilon$ for $\epsilon > 0$, where ρ is non-negative, smooth and compactly supported function on $\mathbf{R}^4$ with $\int \rho = 1$, and $\rho^\epsilon(t,x) = \epsilon^{-5}\rho(\epsilon^{-2}t, \epsilon^{-1}x)$. We consider the classical solution u^ϵ of the equation

$$\partial_t u^\epsilon = (\mathrm{i}+\mu)\triangle u^\epsilon + \nu(1-|u^\epsilon|^2 + C^\epsilon)u^\epsilon + \xi^\epsilon, \quad (t,x) \in \mathbf{R}_+ \times \mathbf{R}^3,$$

with initial condition u_0, where $C^\epsilon = 2C_1^\epsilon - 2\overline{\nu}C_{2,1}^\epsilon - 4\nu C_{2,2}^\epsilon$ is a sum of diverging constants as $\epsilon \downarrow 0$ and precise behaviors of them are stated in Proposition 3.4. We write $\mathbf{R}_+ = (0,\infty)$. We define

$$\mathcal{C}^\eta = \mathcal{C}^\eta(\mathbf{T}^3, \mathbf{C}) = \{u \in \mathcal{B}^\eta_{\infty,\infty}(\mathbf{R}^3, \mathbf{C})\,;\, u(\cdot + k) = u(\cdot) \text{ for any } k \in \mathbf{Z}^3\}.$$

We denote by $C([0,T], \mathcal{C}^\eta)$ the set of all $\mathcal{C}^\eta$-valued continuous functions on $[0,T]$ endowed with the supremum norm $\|\cdot\|_{C([0,T],\mathcal{C}^\eta)}$.

Theorem 2.1. *Let $\eta \in (-\frac{2}{3}, -\frac{1}{2})$. Then for every $u_0 \in \mathcal{C}^\eta$, the sequence $\{u^\epsilon\}$ converges to a limit u in probability as $\epsilon \downarrow 0$. Precisely speaking, this means that there exists an a.s. strictly positive random time T depending on u_0 such that u and u^ϵ for every $\epsilon > 0$ belong to the space $C([0,T], \mathcal{C}^\eta)$ and we have*

$$\|u^\epsilon - u\|_{C([0,T],\mathcal{C}^\eta)} \to 0$$

in probability. Furthermore, u is independent of the choice of ρ.

We use the following notations in Sections 2 and 3:

- For $z = (t, x_1, x_2, x_3) \in \mathbf{R}^4$, we define $\|z\|_s = |t|^{\frac{1}{2}} + |x_1| + |x_2| + |x_3|$.
- For $k = (k_i)_{i=0}^3 \in \mathbf{Z}_+^4$, we define $|k|_s = 2k_0 + k_1 + k_2 + k_3$ and $\partial^k = \partial_t^{k_0}\partial_{x_1}^{k_1}\partial_{x_2}^{k_2}\partial_{x_3}^{k_3}$. Here $\mathbf{Z}_+ = \{0, 1, 2, \dots\}$.
- For $\varphi \in C(\mathbf{R}^4, \mathbf{C})$ and $\delta > 0$, we define the space-time scaling around $z = (t, x_1, x_2, x_3) \in \mathbf{R}^4$ by

  $$\varphi_z^\delta(t', x_1', x_2', x_3') = \delta^{-5}\varphi(\delta^{-2}(t'-t), \delta^{-1}(x_1'-x_1), \delta^{-1}(x_2'-x_2), \delta^{-1}(x_3'-x_3)).$$

- For $\alpha > 0$, we denote by $\mathcal{C}_s^\alpha$ the space of complex-valued functions φ on $\mathbf{R}^4$ such that

  $$\Big|\partial^k\varphi(z') - \sum_{|k+l|_s < \alpha} \frac{(z'-z)^l}{l!}\partial^{k+l}\varphi(z)\Big| \lesssim \|z'-z\|_s^{\alpha-|k|_s} \tag{2.1}$$

  holds locally in $z, z' \in \mathbf{R}^4$ and for every k with $|k|_s < \alpha$.

- Denote by $\mathcal{C}_s^0 = L^\infty_{\mathrm{loc}}(\mathbf{R}^4, \mathbf{C})$ the space of locally bounded functions.

- For $r > 0$, let $\mathcal{B}_r$ be the set of complex-valued smooth functions φ on $\mathbf{R}^4$ supported in the ball $B_s(0,1) = \{z; \|z\|_s \le 1\}$ and such that their derivatives of order up to r are bounded by 1. Let $\alpha < 0$ and $r = \lceil -\alpha \rceil$. Denote by $\mathcal{C}_s^\alpha$ be the space of Schwartz distributions $\xi \in \mathcal{S}' = \mathcal{S}'(\mathbf{R}^4, \mathbf{C})$ such that
$$\|\xi\|_{\alpha;K} := \sup_{z \in K, \varphi \in \mathcal{B}_r, \delta \in (0,1]} \delta^{-\alpha} |\langle \xi, \varphi_z^\delta \rangle| < \infty$$
for every compact set $K \subset \mathbf{R}^4$.

2.1 Results on regularity structures

First we recall basic concepts from the theory of regularity structures [Hai14].

Definition 2.2. We say that a triplet $\mathcal{T} = (A, T, G)$ is a *regularity structure* with *index set* A, *model space* T and *structure group* G, if

- A is a locally finite set of real numbers bounded from below and $0 \in A$.

- $T = \bigoplus_{\alpha \in A} T_\alpha$ with complex Banach spaces $(T_\alpha, \|\cdot\|_\alpha)$. Furthermore, $T_0 \simeq \mathbf{C}$ and its unit vector is denoted by $\mathbf{1}$.

- G is a subgroup of $\mathcal{L}(T)$, the set of continuous linear operators on T, such that, for every $\Gamma \in G$, $\alpha \in A$, and $\tau \in T_\alpha$,
$$\Gamma\tau - \tau \in T_\alpha^- := \bigoplus_{\beta < \alpha} T_\beta.$$
Furthermore, $\Gamma\mathbf{1} = \mathbf{1}$ for every $\Gamma \in G$.

Definition 2.3. Let $\mathcal{T}$ be a regularity structure. We say that a subspace $V = \bigoplus_{\beta \in A} V_\beta$ with $V_\beta \subset T_\beta$ is a *sector* of regularity $\alpha \le 0$ if V is invariant under G (i.e. $\Gamma V \subset V$ for every $\Gamma \in G$) and α is the minimal index such that $V_\alpha \neq \{0\}$. A sector with regularity 0 is called *function-like*.

For $\tau \in T$, we write $\|\tau\|_\alpha = \|\tau_\alpha\|_\alpha$, where τ_α is the component of τ in T_α.

Definition 2.4. Let $\mathcal{T}$ be a regularity structure and let $r = \lceil -\inf A \rceil$. A *model* $Z = (\Pi, \Gamma)$ is a pair of maps $\Gamma : \mathbf{R}^4 \times \mathbf{R}^4 \ni (z, z') \mapsto \Gamma_{zz'} \in G$ and $\Pi : \mathbf{R}^4 \ni z \mapsto \Pi_z \in \mathcal{L}(T, \mathcal{S}')$, the set of continuous linear operators from T to $\mathcal{S}'$, which satisfy
$$\Gamma_{zz'}\Gamma_{z'z''} = \Gamma_{zz''}, \quad \Pi_z \Gamma_{zz'} = \Pi_{z'}$$

for every $z, z', z'' \in \mathbf{R}^4$, and

$$\|\Gamma\|_{\gamma;K} := \sup_{\substack{\beta<\alpha<\gamma,\, \tau\in T_\alpha,\\ (z,z')\in K^2}} \frac{\|\Gamma_{zz'}\tau\|_\beta}{\|\tau\|_\alpha \|z-z'\|_s^{\alpha-\beta}} < \infty,$$

$$\|\Pi\|_{\gamma;K} := \sup_{\substack{\alpha<\gamma,\, \tau\in T_\alpha,\\ z\in K,\, \varphi\in\mathcal{B}_r,\, \delta\in(0,1]}} \frac{|\langle \Pi_z\tau, \varphi_z^\delta\rangle|}{\|\tau\|_\alpha \delta^\alpha} < \infty$$

for every $\gamma > 0$ and compact set $K \subset \mathbf{R}^4$. For models $Z = (\Pi, \Gamma)$ and $Z' = (\Pi', \Gamma')$ on $\mathcal{T}$, we write

$$|\!|\!| Z |\!|\!|_{\gamma;K} = \|\Gamma\|_{\gamma;K} + \|\Pi\|_{\gamma;K}, \quad |\!|\!| Z - Z' |\!|\!|_{\gamma;K} = \|\Gamma - \Gamma'\|_{\gamma;K} + \|\Pi - \Pi'\|_{\gamma;K}.$$

Following [Hai14, Section 6], we define the space of modelled distributions with singularity at $P = \{z = (t,x) \in \mathbf{R}^4 \,;\, t = 0\}$. For a subset $K \subset \mathbf{R}^4$, we denote by

$$K_P := \{(z,z') \in (K \setminus P)^2 \,;\, z \neq z', \|z - z'\|_s \le 1 \wedge |t|^{\frac{1}{2}} \wedge |t'|^{\frac{1}{2}}\}.$$

Definition 2.5. Let $Z = (\Pi, \Gamma)$ be a model on $\mathcal{T}$, $\gamma > 0$ and $\eta \in \mathbf{R}$. For a function $f : \mathbf{R}^4 \to T_\gamma^-$ and a subset $K \subset \mathbf{R}^4$, we define

$$\|f\|_{\gamma,\eta;K} := \sup_{\substack{\beta<\gamma\\ z=(t,x)\in K\setminus P}} (1 \wedge |t|^{\frac{\beta-\eta}{2}\vee 0}) \|f(z)\|_\beta,$$

$$|\!|\!| f |\!|\!|_{\gamma,\eta;K} := \|f\|_{\gamma,\eta;K} + \sup_{\substack{\beta<\gamma\\ (z,z')\in K_P}} (1 \wedge |t| \wedge |t'|)^{\frac{\gamma-\eta}{2}} \frac{\|f(z) - \Gamma_{zz'} f(z')\|_\beta}{\|z-z'\|_s^{\gamma-\beta}}.$$

We write $f \in \mathcal{D}_P^{\gamma,\eta} = \mathcal{D}_P^{\gamma,\eta}(Z)$ if $|\!|\!| f |\!|\!|_{\gamma,\eta;K} < \infty$ for every compact subset $K \subset \mathbf{R}^4$.

If f takes value in a sector V, we write $f \in \mathcal{D}_P^{\gamma,\eta}(V; Z)$. For models Z, Z' and $f \in \mathcal{D}_P^{\gamma,\eta}(Z)$, $f' \in \mathcal{D}_P^{\gamma,\eta}(Z')$, we define

$$|\!|\!| f; f' |\!|\!|_{\gamma,\eta;K} = \|f - f'\|_{\gamma,\eta;K} + \sup_{\substack{\alpha<\gamma,\\ (z,z')\in K_P}} (1 \wedge |t| \wedge |t'|)^{\frac{\gamma-\eta}{2}} \frac{\|f(z) - f'(z) - \Gamma_{zz'} f(z') + \Gamma'_{zz'} f'(z')\|_\alpha}{\|z-z'\|_s^{\gamma-\alpha}}.$$

We denote by $\mathcal{M} \ltimes \mathcal{D}_P^{\gamma,\eta}$ the set of all pairs (Z, f) of a model Z and $f \in \mathcal{D}_P^{\gamma,\eta}(Z)$. The topology on $\mathcal{M} \ltimes \mathcal{D}_P^{\gamma,\eta}$ is defined by the family of pseudo-metrics $\{|\!|\!| \cdot ; \cdot |\!|\!|_{\gamma,\eta;K}\}$.

Theorem 2.6 ([Hai14, Theorem 3.10 and Proposition 6.9])**.** *Let $Z = (\Pi, \Gamma)$ be a model on $\mathcal{T}$. Let V be a sector with regularity $\alpha \le 0$ and let $r = \lceil -\alpha \rceil$. If*

$\gamma > 0$, $\eta \le \gamma$, and $\alpha \wedge \eta > -2$, then there exists a unique continuous linear map $\mathcal{R} : \mathcal{D}_P^{\gamma,\eta}(V) \to \mathcal{C}_s^{\alpha\wedge\eta}$ such that, if K and K' are compact subsets of $\mathbf{R}^4$ such that K is included in the interior of K', then we have

$$|\langle \mathcal{R}f - \Pi_z f(z), \varphi_z^\delta \rangle| \lesssim \delta^\gamma \|\Pi\|_{\gamma;K'} |\!|\!| f |\!|\!|_{\theta,\eta;K'}, \tag{2.2}$$

uniformly over $f \in \mathcal{D}_P^{\gamma,\eta}(V)$, $\delta \in (0,1]$, $z \in K$ and $\varphi \in \mathcal{B}_r$ with φ_z^δ supported in K' and uniformly away from P. Furthermore, the map $\mathcal{M} \ltimes \mathcal{D}_P^{\gamma,\eta}(V) \ni (Z, f) \to \mathcal{R}f \in \mathcal{C}_s^{\alpha\wedge\eta}$ is locally uniformly continuous.

Remark 2.7 ([Hai14, Lemma 6.7])**.** The reconstruction operator $\mathcal{R}$ is local in the sense that, the behavior of $\mathcal{R}f$ on the compact set $K \subset \mathbf{R}^4$ is uniquely determined by the values of f and Π in an arbitrary neighborhood of K.

Next we introduce specific symbols and operators to describe (1.1) by regularity structure: the polynomial structure, product, integration against Green's function, and the complex conjugate.

We have the regularity structure $\mathcal{T}^{\mathrm{poly}}$ given by all polynomials in the symbols X_0, X_1, X_2, X_3, which denote the time and space directions, respectively. Denote $X^k = \prod_{i=0}^3 X_i^{k_i}$ for a multi-index $k \in \mathbf{Z}_+^4$, and $\mathbf{1} = X^{(0,0,0,0)}$. We endow these with the parabolic degrees $|X^k|_s = |k|_s$. Now we define the model space $T^{\mathrm{poly}} = \bigoplus_{n\in\mathbf{Z}_+} T_n^{\mathrm{poly}}$, where

$$T_n^{\mathrm{poly}} = \langle X^k \,;\, |k|_s = n \rangle.$$

The group $G = \mathbf{R}^4$ acts on T^{poly} by defining $\Gamma_h X^k = (X - h\mathbf{1})^k$ for every $h \in \mathbf{R}^4$. Now we have the regularity structure $\mathcal{T}^{\mathrm{poly}} = (\mathbf{Z}_+, T^{\mathrm{poly}}, \mathbf{R}^4)$. Furthermore, we have the canonial model (Π, Γ) on $\mathcal{T}^{\mathrm{poly}}$ given by

$$(\Pi_z X^k)(z') = (z' - z)^k, \quad \Gamma_{zz'} = \Gamma_{z'-z}, \tag{2.3}$$

for every $z, z' \in \mathbf{R}^4$.

Throughout this section, the regularity structure $\mathcal{T} = (A, T, G)$ contains $\mathcal{T}^{\mathrm{poly}}$, i.e. T^{poly} is contained as a sector and the restriction of G on T^{poly} coincides with $\{\Gamma_h \,;\, h \in \mathbf{R}^4\}$. The model (Π, Γ) acts on T^{poly} by (2.3). Furthermore, we assume that for every $n \in \mathbf{Z}_+$, $T_n = T_n^{\mathrm{poly}}$.

Proposition 2.8 ([Hai14, Proposition 3.28])**.** *Let V be a function-like sector which contains T^{poly} and such that $V \subset T^{\mathrm{poly}} + T_\alpha^+$ for some $\alpha > 0$, where $T_\alpha^+ := \bigoplus_{\alpha\le\beta} T_\beta$. Let $\gamma > \alpha$, $\eta \in \mathbf{R}$. Then for every $f \in \mathcal{D}_P^{\gamma,\eta}(V)$, $\mathcal{R}f$ coincides with the component of f in $V_0 = \langle \mathbf{1} \rangle$ and belongs to $\mathcal{C}_s^\alpha((0,\infty) \times \mathbf{R}^3)$, the space of functions φ such that the estimate* (2.1) *holds uniformly over $z, z' \in K$ for every compact set $K \subset (0,\infty) \times \mathbf{R}^3$.*

For a pair of sectors (V, W), a *product* $* : V \times W \to T$ is a continuous bilinear map such that

- $V_\alpha * W_\beta \subset T_{\alpha+\beta}$ for every $\alpha, \beta \in A$,
- $\mathbf{1} * w = w$ for every $w \in W$ and $v * \mathbf{1} = v$ for every $v \in V$,
- $\Gamma(v * w) = (\Gamma v) * (\Gamma w)$ for every $(v, w) \in V \times W$ and $\Gamma \in G$.

The canonical product on T^{poly} is given by $X^k * X^l = X^{k+l}$.

Proposition 2.9 ([Hai14, Proposition 6.12])**.** *Let (V, W) be a pair of sectors with regularities α_1, α_2, respectively, and product $* : V \times W \to T$. For every $f_1 \in \mathcal{D}_P^{\gamma_1, \eta_1}(V)$ and $f_2 \in \mathcal{D}_P^{\gamma_2, \eta_2}(W)$, the function $f = f_1 * f_2$ (projected onto T_γ^-) belongs to $\mathcal{D}_P^{\gamma, \eta}$ with $\gamma = (\gamma_1 + \alpha_2) \wedge (\gamma_2 + \alpha_1)$ and $\eta = (\eta_1 + \alpha_2) \wedge (\eta_2 + \alpha_1) \wedge (\eta_1 + \eta_2)$. Furthermore, this bilinear map is locally uniformly continuous with respect to the topology of $\mathcal{M} \ltimes \mathcal{D}_P^{\gamma, \eta}$.*

We say that a function $K : \mathbf{R}^4 \setminus \{0\} \to \mathbf{C}$ is a *regularizing kernel* (of order 2) if it can be written by $K = \sum_{n \geq 0} K_n$, where $\{K_n\}$ satisfies the following assumptions.

Assumption 2.10.
- *$K_n : \mathbf{R}^4 \to \mathbf{C}$ is smooth and supported in a ball $B_s(0, 2^{-n})$.*
- *There exists a constant $C > 0$ such that $\sup_z |\partial^k K_n(z)| \leq C 2^{(3+|k|_s)n}$ for every $n \geq 0$ and $k \in \mathbf{Z}_+^4$.*
- *There exists $r > 0$ such that $\int_{\mathbf{R}^4} K_n(z) z^k dz = 0$ for every $n \geq 0$ and k with $|k|_s \leq r$.*

For a sector V, an *abstract integration map* $\mathcal{I} : V \to T$ is a continuous linear map such that

- $\mathcal{I} V_\alpha \subset T_{\alpha+2}$ for every $\alpha \in A$ such that $\alpha + 2 \in A$,
- $\mathcal{I}\tau = 0$ for every $\tau \in V \cap T^{\mathrm{poly}}$,
- $(\mathcal{I}\Gamma - \Gamma\mathcal{I})V \subset T^{\mathrm{poly}}$ for every $\Gamma \in G$.

Given a sector V and an abstract integration map $\mathcal{I}$, we say that a model (Π, Γ) *realizes* a regularizing kernel K for $\mathcal{I}$, if for every $\alpha \in A$, $\tau \in V_\alpha$ and $z \in \mathbf{R}^4$ we have

$$\Pi_z \mathcal{I}\tau = K * (\Pi_z \tau) - \Pi_z \mathcal{J}(z)\tau,$$

where $\mathcal{J}(z)\tau = \sum_{|k|_s<\alpha+2} \frac{1}{k!} X^k (\partial^k K * \Pi_z \tau)(z)$. It is a consequence of Assumption 2.10 that $(\partial^k K * \Pi_z \tau)(z)$ is defined for all k with $|k|_s < \alpha + 2$.

For $f \in \mathcal{D}_P^{\gamma,\eta}(V)$, we define the modelled distribution $\mathcal{K}_\gamma f$ by

$$(\mathcal{K}_\gamma f)(z) = \mathcal{I} f(z) + \mathcal{J}(z) f(z) + (\mathcal{N}_\gamma f)(z),$$

where $(\mathcal{N}_\gamma f)(z) = \sum_{|k|_s<\gamma+2} \frac{1}{k!} X^k \partial^k K * (\mathcal{R} f - \Pi_z f(z))(z)$.

Proposition 2.11 ([Hai14, Proposition 6.16])**.** *Let V be a sector of regularity α and with an abstract integration map $\mathcal{I}$. Let $\gamma > 0$ and $\eta < \gamma$. Assume that $\eta \wedge \alpha > -2$ and $\gamma + 2, \eta + 2 \notin \mathbf{Z}_+$. Then, for every model Z realizing K for $\mathcal{I}$, the operator $\mathcal{K}_\gamma$ maps $\mathcal{D}_P^{\gamma,\eta}(V)$ into $\mathcal{D}_P^{\gamma',\eta'}$ with $\gamma' = \gamma + 2$, $\eta' = \eta \wedge \alpha + 2$, and for every $f \in \mathcal{D}_P^{\gamma,\eta}(V)$, we have*

$$\mathcal{R}\mathcal{K}_\gamma f = K * \mathcal{R} f.$$

Furthermore, the map $\mathcal{K}_\gamma : \mathcal{M} \ltimes \mathcal{D}_P^{\gamma,\eta}(V) \to \mathcal{D}_P^{\gamma',\eta'}$ is locally uniformly continuous.

Remark 2.12. Even in the case that $\eta \wedge \alpha \le -2$, if there exists a distribution $\mathcal{R} f \in \mathcal{C}_s^{\eta\wedge\alpha}$ which satisfies (2.2), then Proposition 2.11 still holds.

Given a sector V, we say that a continuous antilinear map $V \ni \tau \mapsto \overline{\tau} \in T$ (in the sense that $\overline{\lambda_1\tau_1 + \lambda_2\tau_2} = \overline{\lambda_1}\,\overline{\tau_1} + \overline{\lambda_2}\,\overline{\tau_2}$ for $\lambda_1, \lambda_2 \in \mathbf{C}$ and $\tau_1, \tau_2 \in V$) is *complex conjugate*, if $\overline{V_\alpha} \subset T_\alpha$ for every $\alpha \in A$, and

$$\overline{X^k} = X^k, \quad \Gamma\overline{\tau} = \overline{\Gamma\tau}$$

for every $\Gamma \in G$ and $\tau \in V$. Then $\overline{V} = \{\overline{\tau}\,; \tau \in V\}$ is also a sector. We assume that a model (Π, Γ) is compatible with the complex conjugate, i.e.

$$\Pi_z \overline{\tau} = \overline{\Pi_z \tau}$$

for every $z \in \mathbf{R}^4$ and $\tau \in T$. From these assumptions, we see that the map $\mathcal{D}_P^{\gamma,\eta}(V) \ni f \mapsto \overline{f} \in \mathcal{D}_P^{\gamma,\eta}$ is continuous antilinear, and $\mathcal{R}\overline{f} = \overline{\mathcal{R} f}$ holds.

2.2 Regularity structures associated with CGL and admissible models

For smooth ξ, the CGL equation (1.1) is equivalent to the mild form

$$u = G * \{\mathbf{1}_{t>0}(\xi + \nu(1 - u\overline{u})u)\} + Gu_0, \tag{2.4}$$

where $*$ denotes the space-time convolution, G is the fundamental solution of

$$\partial_t G = (\mathrm{i} + \mu)\triangle G, \quad (t,x) \in \mathbf{R}_+ \times \mathbf{R}^3, \tag{2.5}$$

with initial condition $G(0,\cdot) = \delta_0$, extended into the function $G : \mathbf{R}^4 \setminus \{0\} \to \mathbf{C}$ by $G(t,\cdot) \equiv 0$ if $t \le 0$, and Gu_0 denotes the solution of (2.5) with initial condition u_0.

We construct a regularity structure associated with (2.4) by following [Hai14, Section 8.1]. We assumed that polynomials $\{X^k\}$ are contained in our regularity structure. Additionally we have symbols Ξ (noise), an abstract integration $\mathcal{I}$ (space-time convolution with G), and the complex conjugate. Inspired from (2.4), we can define $\tilde{\mathcal{F}}$ as the smallest set of symbols such that $\{\Xi, X^k\} \subset \tilde{\mathcal{F}}$ and closed for the operations:

- If $\tau, \tau' \in \tilde{\mathcal{F}}$, then $\tau\tau' = \tau'\tau \in \tilde{\mathcal{F}}$.
- If $\tau \in \tilde{\mathcal{F}}$, then $\overline{\tau} \in \tilde{\mathcal{F}}$, where we set $\overline{X^k} = X^k$.
- If $\tau \in \tilde{\mathcal{F}} \setminus \{X^k\}$, then $\mathcal{I}\tau \in \tilde{\mathcal{F}}$.

For a fixed number $\alpha < -\frac{5}{2}$, we define the homogeneity of each variable by

$$|\Xi|_s = \alpha, \quad |X^k|_s = |k|_s, \quad |\tau\tau'|_s = |\tau|_s + |\tau'|_s, \quad |\overline{\tau}|_s = |\tau|_s, \quad |\mathcal{I}\tau|_s = |\tau|_s + 2.$$

However, $\tilde{\mathcal{F}}$ is too big. Precisely we consider the subsets $\mathcal{U}$ and $\mathcal{W}$, which are defined by the smallest sets such that $\{\Xi, X^k\} \subset \mathcal{W}$, $\{X^k\} \subset \mathcal{U}$ and

$$\tau \in \mathcal{W} \Rightarrow \mathcal{I}\tau \in \mathcal{U}, \quad \tau_1, \tau_2, \tau_3 \in \mathcal{U} \Rightarrow \tau_1\tau_2\overline{\tau_3} \in \mathcal{W}.$$

We set $\mathcal{F} = \mathcal{U} \cup \mathcal{W}$, and define

$$T_\beta = \langle \tau \in \mathcal{F} \,;\, |\tau|_s = \beta \rangle, \quad T = \langle \mathcal{F} \rangle, \quad U = \langle \mathcal{U} \rangle, \quad W = \langle \mathcal{W} \rangle.$$

We can see that T contains all polynomials T^{poly}, and furthermore, the abstract integration $\mathcal{I}$, the complex conjugate, and the product $U \times U \times \overline{U} \to W$ are well defined. Here $\overline{U} = \langle \overline{\tau} \,;\, \tau \in \mathcal{U} \rangle$.

Remark 2.13. We do not assume identifications of symbols

$$\overline{\overline{\tau}} = \tau, \quad \overline{\tau_1\tau_2} = \overline{\tau_1}\,\overline{\tau_2}$$

since $\overline{\overline{\tau}}$ and $\overline{\tau_1}\,\overline{\tau_2}$ are not involved in the definition of $\mathcal{F}$.

In order to define T as a model space of a regularity structure, the set $\{|\tau|_s\,;\tau\in\mathcal{F}\}$ must be bounded from below. A nonlinear SPDE is called *sub-critical*, if the nonlinear terms formally disappear in some scaling which keeps the linear part and the noise term invariant. This is equivalent to the property that all symbols except Ξ defined as above have homogeneities strictly greater than $|\Xi|_s$ ([Hai14, Assumption 8.3]). In the present case, this is equivalent to $|(\mathcal{I}\Xi)^2\overline{\mathcal{I}\Xi}|_s = 3(2+\alpha) > \alpha$, or $\alpha > -3$.

We need to define the structure group acting on T. Let T^+ be the complex free commutative algebra generated by abstract symbols

$$\mathcal{F}^+ = \{X^k\} \cup \{\mathcal{J}_k\tau, \overline{\mathcal{J}_k\tau}\,;\tau\in\mathcal{F}\setminus\{X^k\}, |k|_s < |\tau|_s + 2\}.$$

We define the homogeneity of each variable by

$$|X^k|_s = |k|_s, \quad |\tau\tau'|_s = |\tau|_s + |\tau'|_s, \quad |\mathcal{J}_k\tau|_s = |\overline{\mathcal{J}_k\tau}|_s = |\tau|_s + 2 - |k|_s.$$

In the following, we will view $\mathcal{J}_k$ as a map from T to T^+, by defining $\mathcal{J}_k\tau = 0$ if $\tau = X^k$ or $|\tau|_s + 2 - |k|_s \le 0$, and linearly extending it for all $\tau\in T$.

We construct two linear maps Δ and Δ^+ recursively as follows. The linear map $\Delta : T \to T\otimes T^+$ is defined by

$$\begin{aligned}
&\Delta\mathbf{1} = \mathbf{1}\otimes\mathbf{1}, \quad \Delta X_i = X_i\otimes\mathbf{1} + \mathbf{1}\otimes X_i \ \ (i=0,1,2,3), \quad \Delta\Xi = \Xi\otimes\mathbf{1},\\
&\Delta(\tau\tau') = (\Delta\tau)(\Delta\tau'), \quad \Delta\overline{\tau} = \overline{\Delta\tau},\\
&\Delta\mathcal{I}\tau = (\mathcal{I}\otimes\mathrm{Id}_{T^+})\Delta\tau + \sum_{l,m}\frac{X^l}{l!}\otimes\frac{X^m}{m!}\mathcal{J}_{l+m}\tau.
\end{aligned}$$

The linear map $\Delta^+ : T^+ \to T^+\otimes T^+$ is defined by

$$\begin{aligned}
&\Delta^+\mathbf{1} = \mathbf{1}\otimes\mathbf{1}, \quad \Delta^+ X_i = X_i\otimes\mathbf{1} + \mathbf{1}\otimes X_i \ \ (i=0,1,2,3),\\
&\Delta^+(\tau\tau') = (\Delta^+\tau)(\Delta^+\tau'), \quad \Delta^+\overline{\tau} = \overline{\Delta^+\tau},\\
&\Delta^+\mathcal{J}_k\tau = \sum_l\left(\mathcal{J}_{k+l}\otimes\frac{(-X)^l}{l!}\right)\Delta\tau + \mathbf{1}\otimes\mathcal{J}_k\tau.
\end{aligned}$$

Then by [Hai14, Theorem 8.16], the pair (T^+,Δ^+) is a Hopf algebra, i.e. Δ^+ satisfies the identity

$$(\mathrm{Id}_{T^+}\otimes\Delta^+)\Delta^+ = (\Delta^+\otimes\mathrm{Id}_{T^+})\Delta^+$$

and the algebra homomorphism $\mathbf{1}^* : T^+ \to T^+$ defined by $\mathbf{1}^*(\mathbf{1}) = \mathbf{1}$ and $\mathbf{1}^*(\tau) = 0$ for $\tau\in\mathcal{F}^+\setminus\{\mathbf{1}\}$ is a counit in the sense that

$$(\mathbf{1}^*\otimes\Delta^+)\Delta^+ = (\Delta^+\otimes\mathbf{1}^*)\Delta^+,$$

and furthermore, the algebra homomorphism $\mathcal{A} : T^+ \to T^+$ recursively defined by

$$\mathcal{A}X^k = (-X)^k, \quad \mathcal{A}\overline{\tau} = \overline{\mathcal{A}\tau}, \quad \mathcal{A}\mathcal{J}_k\tau = -\sum_l \mathcal{M}\left(\mathcal{J}_{k+l} \otimes \frac{X^l}{l!}\mathcal{A}\right)\Delta\tau,$$

where $\mathcal{M} : T^+ \otimes T^+ \ni \tau \otimes \tau' \mapsto \tau\tau' \in T^+$, is an antipode of T^+ in the sense that

$$\mathcal{M}(\mathrm{Id}_{T^+} \otimes \mathcal{A})\Delta^+ = \mathbf{1}^* = \mathcal{M}(\mathcal{A} \otimes \mathrm{Id}_{T^+})\Delta^+.$$

The pair (T, Δ) is a comodule over T^+, i.e. Δ satisfies the identity

$$(\mathrm{Id}_{T^+} \otimes \Delta^+)\Delta^+ = (\Delta^+ \otimes \mathrm{Id}_{T^+})\Delta^+.$$

We denote by G the set of algebra homomorphisms $g : T^+ \to \mathbf{C}$ such that $g(\overline{\tau}) = \overline{g(\tau)}$ for every $\tau \in T^+$. Then G is a group with the product $\circ$ defined by

$$g \circ g' = (g \otimes g')\Delta^+, \quad g, g' \in G.$$

The inverse of $g \in G$ is given by $g^{-1} = g\mathcal{A}$. Each $g \in G$ acts on T as the operator $\Gamma_g \in \mathcal{L}(T)$ defined by

$$\Gamma_g\tau = (\mathrm{Id}_T \otimes g)\Delta\tau, \quad \tau \in T.$$

The following theorem is a modification of [Hai14, Theorem 8.24].

Theorem 2.14. *Let $\alpha \in (-3, -\frac{5}{2})$ and $A = \{|\tau|_s \,; \tau \in \mathcal{F}\}$. Then $\mathcal{T}_{\mathrm{cgl}} := (A, T, G)$ is a regularity structure which contains the polynomial structure $\mathcal{T}^{\mathrm{poly}}$ and has the complex conjugate on U, the abstract integration map $\mathcal{I} : W \to T$, and the products $* : U \times U \to UU$ and $* : UU \times \overline{U} \to W$, where $UU = \langle \tau\tau' \,; \tau, \tau' \in U \rangle$.*

We introduce a class of suitable models associated with $\mathcal{T}_{\mathrm{cgl}}$. Let K be a regularizing kernel satisfying Assumption 2.10 with $r > 0$. We denote by $\mathcal{T}_{\mathrm{cgl}}^{(r)}$ the regularity structure obtained by $T_\gamma = 0$ for $\gamma > r$.

Definition 2.15. We say that a model (Π, Γ) on $\mathcal{T}_{\mathrm{cgl}}^{(r)}$ is *admissible*, if

- Π realizes K for $\mathcal{I}$, compatible with the complex conjugate, and satisfies (2.3),
- $\Gamma_{zz'} = (\Gamma_{f_z})^{-1}\Gamma_{f_{z'}}$, where $f_z \in G$ is defined by $f_z(X^k) = (-z)^k$ and

$$f_z(\mathcal{J}_k\tau) = -\partial^k K * (\Pi_z\tau)(z), \quad |\tau|_s + 2 - |k|_s > 0.$$

If the model (Π, Γ) is admissible, then the map $\Pi_z(\Gamma_{f_z})^{-1} : T \to \mathcal{S}'$ is independent to z, so that we can write $\mathbf{\Pi} = \Pi_z(\Gamma_{f_z})^{-1}$. Furthermore we have

$$(\mathbf{\Pi} X^k)(z) = z^k, \quad \mathbf{\Pi}\mathcal{I}\tau = K * (\mathbf{\Pi}\tau), \quad \mathbf{\Pi}\overline{\tau} = \overline{\mathbf{\Pi}\tau}.$$

Conversely, if a linear map $\mathbf{\Pi} : T \to \mathcal{S}'$ satisfies these conditions, and a family $\{f_z\,; z \in G\}$ satisfies $f_z(X^k) = (-z)^k$ and $f_z(\mathcal{J}_k\tau) = -\partial^k K * (\mathbf{\Pi}\Gamma_{f_z}\tau)(z)$ for $\mathcal{J}_k\tau$ with $|\tau|_s + 2 - |k|_s > 0$, then the corresponding admissible model (Π, Γ) is uniquely determined.

We assume that the model is periodic in the space direction. For $n \in \mathbf{Z}^3$ and $z = (t, x) \in \mathbf{R}^4$, we write $S_n z = (t, x + n)$.

Definition 2.16. We say that a model (Π, Γ) on $\mathcal{T}_{\mathrm{cgl}}^{(r)}$ is *periodic* if

$$(\Pi_{S_n z}\tau)(S_n z') = (\Pi_z\tau)(z'), \quad \Gamma_{(S_n z)(S_n z')} = \Gamma_{zz'},$$

for every $z, z' \in \mathbf{R}^4$ and $n \in \mathbf{Z}^3$.

2.3 Abstract solution map

In the regularity structures constructed above, we can reformulate (1.1) as a fixed point problem in the space $\mathcal{D}_P^{\gamma,\eta}$, by following [Hai14]. First, note that the fundamental solution of (2.5) is given by

$$G(t, x) = \mathbf{1}_{t>0} \frac{1}{\sqrt{4\pi(\mathsf{i} + \mu)t}^3} \exp\left(-\frac{|x|^2}{4(\mathsf{i} + \mu)t}\right).$$

Here, for $\lambda = re^{\mathsf{i}\theta} \in \mathbf{C}$ $(r > 0, \theta \in (-\frac{\pi}{2}, \frac{\pi}{2}))$, the square root is defined by $\sqrt{\lambda} = \sqrt{r}e^{\mathsf{i}\theta/2}$. Hence G has the form $G(t, x) = t^{-3/2}\hat{G}(t^{-1/2}x)$ for some $\hat{G} \in \mathcal{S}(\mathbf{R}^3, \mathbf{C})$ when $t > 0$, which satisfies the condition in [Hai14, Lemma 7.4].

Lemma 2.17 ([Hai14, Lemma 7.7])**.** *There exist a regularizing kernel K and a smooth function R with compact support such that*

$$(G * u)(z) = (K * u)(z) + (R * u)(z)$$

holds for every periodic function u supported in $\mathbf{R}_+ \times \mathbf{R}^3$ and $z \in (-\infty, 1] \times \mathbf{R}^3$. Furthermore, K and R are supported in $\mathbf{R}_+ \times \mathbf{R}^3$, and K satisfies Assumption 2.10 *with arbitrary fixed $r > 0$.*

For a periodic distribution $\xi \in \mathcal{S}'$, we define the modelled distribution

$$(R_\gamma\xi)(z) = \sum_{|k|_s < \gamma} \frac{X^k}{k!} \partial^k R * \xi(z).$$

Now we reformulate (2.4) as a fixed point problem in $\mathcal{D}_P^{\gamma,\eta}$. First, for every periodic initial condition $u_0 \in \mathcal{C}^\eta$ with $\eta < 0$ and $\eta \notin \mathbf{Z}$, the function Gu_0 is canonically lifted to an element of $\mathcal{D}_P^{\gamma,\eta}$ for every $\gamma > \eta$, by defining $(Gu_0)(z) = \sum_{|k|_s<\gamma} \frac{1}{k!} X^k \partial^k G u_0(z)$ ([Hai14, Lemma 7.5]). Second, note that by Proposition 2.9, the map $u \mapsto u^2\overline{u}$ is locally Lipschitz continuous from $\mathcal{D}_P^{\gamma,\eta}(U)$ to $\mathcal{D}_P^{\gamma+2\alpha+4,3\eta}$, if $\gamma > |2\alpha+4|$ and $\eta \le \alpha+2$. Therefore it is suitable to consider the problem

$$u = (\mathcal{K}_{\gamma+2\alpha+4} + R_\gamma \mathcal{R})(\mathbf{1}_{t>0} F(u)) + Gu_0, \quad F(u) = \Xi + \nu(1 - u\overline{u})u, \tag{2.6}$$

in $u \in \mathcal{D}_P^{\gamma,\eta}$. However, $F(u)$ takes values in the sector U of regularity $\alpha = |\Xi|_s < -\frac{5}{2} < -2$, so that Theorem 2.6 is not sufficient to define $\mathcal{R}F(u)$. In order to overcome this problem, we impose the following assumption on the distribution $\xi = \mathbf{\Pi}\Xi = \Pi_z \Xi$. (Since Ξ is G-invariant, $\Pi_z \Xi$ is independent to z.)

Assumption 2.18. *(1) For $\alpha < 0$, we denote by $\bar{\mathcal{C}}_s^\alpha$ the completion of smooth functions under the family of norms:*

$$\big|\!\big|\!\big|\xi\big|\!\big|\!\big|_{\alpha;K} := \sup_{s\in\mathbf{R}} \|\mathbf{1}_{t>s}\xi\|_{\alpha;K}$$

for all compact sets $K \subset \mathbf{R}^4$. We assume that $\xi = \mathbf{\Pi}\Xi$ belongs to $\bar{\mathcal{C}}_s^\alpha$ for $\alpha = |\Xi|_s$.

(2) $K * \xi$ belongs to the space $C(\mathbf{R}, \mathcal{C}^{\alpha+2})$.

Under the assumption $\xi \in \bar{\mathcal{C}}_s^\alpha$, we can define $\mathcal{R}(\mathbf{1}_{t>0}\Xi) := \mathbf{1}_{t>0}\xi$, so that $\mathcal{K}(\mathbf{1}_{t>0}\Xi)$ is also defined.

The following theorem is a modification of [Hai14, Theorem 7.8 and Proposition 9.8]. We denote by $O_T = (-\infty, T] \times \mathbf{R}^3$ and $|\!|\!| \cdot |\!|\!|_{\gamma,\eta;T} := |\!|\!| \cdot |\!|\!|_{\gamma,\eta;O_T}$.

Theorem 2.19. *Let $\alpha \in (-\frac{18}{7}, -\frac{5}{2})$, $\gamma > |2\alpha+4|$ and $\eta \in (-\frac{2}{3}, \alpha+2)$. Assume that the regularizing kernel K satisfies Assumption 2.10 with $r > \gamma + 2\alpha + 6 (> 2)$. Then for every admissible and periodic model $Z = (\Pi, \Gamma)$ satisfying Assumption 2.18 and every periodic $u_0 \in \mathcal{C}^\eta$, there exists $T \in (0, \infty]$ such that the fixed point problem (2.6) admits a unique solution $u \in \mathcal{D}_P^{\gamma,\eta}(U)$ on $(0, T)$. The time T can be chosen maximal in the sense that $\lim_{t\uparrow T} \|\mathcal{R}u(t,\cdot)\|_{\mathcal{C}^\eta} = \infty$ unless $T = \infty$.*

Furthermore, the solution u and the survival time T depend on $(u_0, Z, \xi, K * \xi)$ locally uniformly continuously and locally uniformly lower semi-continuously, respectively, in the topology of $\mathcal{C}^\eta \times \mathcal{M} \times \bar{\mathcal{C}}_s^\alpha \times C(\mathbf{R}, \mathcal{C}^{\alpha+2})$.

Proof. We consider Ξ and $F^+(u) = \nu(1 - u\overline{u})u$ separately. Here $F^+(u)$ takes values in the sector W^+ with regularity $|(\mathcal{I}\Xi)^2\overline{\mathcal{I}\Xi}|_s = 3(\alpha+2)$. Let $\mathcal{G}_\gamma = \mathcal{K}_{\gamma+2\alpha+4} + R_\gamma\mathcal{R}$. The modelled distribution $\mathbf{1}_{t>0}\Xi$ belongs to $\mathcal{D}_P^{\gamma,\alpha}$ for every γ. Hence under Assumption 2.18-(1), $\mathcal{G}_\gamma\Xi$ is defined as an element of $\mathcal{D}_P^{\gamma,\alpha+2}(U)$. On the other hand, $\mathcal{G}_\gamma$ maps $\mathcal{D}_P^{\gamma+2\alpha+4,3\eta}(W^+)$ into $\mathcal{D}_P^{\gamma,3\eta+2}(U)$ provided that $3\eta > -2$, as a consequence of Proposition 2.11. Furthermore, following the arguments in [Hai14, Theorem 7.1], we have the bound

$$\vertiii{\mathcal{G}_\gamma(\mathbf{1}_{t>0}(\Xi + F^+(u)))}_{\gamma,\eta;T} \lesssim T^{\frac{\alpha+2-\eta}{2}}(1 + \vertiii{F^+(u)}_{\gamma+2\alpha+4,3\eta;T})$$

for every periodic $u \in \mathcal{D}_P^{\gamma,\eta}(U)$. As in [Hai14, Theorem 7.8], this yields that there exists small $T > 0$ such that (2.6) admits a unique solution $u \in \mathcal{D}_P^{\gamma,\eta}(U)$ on $(0,T)$.

To glue local solutions up to maximal time where the solution exists, note that $\mathcal{R}u$ belongs to the space $C((0,T),\mathcal{C}^\eta)$, even though $\eta < 0$. Indeed, the solution can be written by $u = \mathcal{I}\Xi + u^+$, where u^+ takes values in the function-like sector U^+. As in Proposition 2.8, $\mathcal{R}u^+$ is Hölder continuous. By Assumption 2.18-(2), $\mathcal{R}\mathcal{I}\Xi = K * \xi$ belongs to $C(\mathbf{R},\mathcal{C}^\eta)$. For $s \in (0,T)$, we start from $u_s \in \mathcal{C}^\eta$ and consider the problem

$$u = \mathcal{G}_\gamma(\mathbf{1}_{t>s}(\Xi + F^+(u))) + Gu_s,$$

which is well-posed by defining $\mathcal{R}(\mathbf{1}_{t>s}\Xi) := \mathbf{1}_{t>s}\xi$. This can extend the time interval where the local solution exists, following [Hai14, Proposition 7.11]. The existence of maximal solution and its continuity with respect to $(u_0, Z, \xi, K*\xi)$ are obtained by standard arguments in PDE theory. □

2.4 Renormalization

For each $\epsilon > 0$, the noise ξ^ϵ defined in the beginning of Section 2 can be lifted to an admissible and periodic model $Z^\epsilon = (\Pi^\epsilon, \Gamma^\epsilon)$ on $\mathcal{T}_{\mathrm{cgl}}^{(r)}$, by defining the linear map $\mathbf{\Pi}^\epsilon : T \to C^\infty(\mathbf{R}^4)$ with the additional assumptions:

$$\mathbf{\Pi}^\epsilon\Xi = \xi^\epsilon, \quad \mathbf{\Pi}^\epsilon(\tau\tau') = (\mathbf{\Pi}^\epsilon\tau)(\mathbf{\Pi}^\epsilon\tau').$$

Furthermore, Z^ϵ has the property that $\Pi_z\tau$ is a smooth function for every $\tau \in T$ and $z \in \mathbf{R}^4$, then as a consequence, $\mathcal{R}^\epsilon f$ is also smooth and satisfies

$$(\mathcal{R}^\epsilon f)(z) = (\Pi_z^\epsilon f(z))(z)$$

for every modelled distribution f ([Hai14, Remark 3.15]).

We introduce a renormalization of Z^ϵ following [Hai14, Section 8.3]. Let $\mathcal{F}_0 \subset \mathcal{F}$ be a subset such that $\{\tau \in \mathcal{F}\,; |\tau|_s \le 0\} \subset \mathcal{F}_0$, and there exists a subset $\mathcal{F}_* \subset \mathcal{F}_0$ such that $\Delta\mathcal{F}_0 \subset \langle \mathcal{F}_0\rangle \otimes T_0^+$, where T_0^+ is the complex free commutative algebra generated by symbols

$$\mathcal{F}_0^+ = \{X^k\} \cup \{\mathcal{J}_k\tau, \overline{\mathcal{J}_k\tau}\,; \tau \in \mathcal{F}_*, |k|_s < |\tau|_s + 2\}.$$

Let $M : \langle \mathcal{F}_0\rangle \to \langle \mathcal{F}_0\rangle$ be a linear map such that

$$M\mathcal{I}\tau = \mathcal{I}M\tau, \quad M\overline{\tau} = \overline{M\tau}, \quad MX^k = X^k.$$

Then two linear maps $\hat{M} : T_0^+ \to T_0^+$ and $\Delta^M : \langle \mathcal{F}_0\rangle \to \langle \mathcal{F}_0\rangle \otimes T_0^+$ are uniquely determined by

$$\begin{aligned}
&\hat{M}X^k = X^k, \quad \hat{M}(\tau\tau') = (\hat{M}\tau)(\hat{M}\tau'),\\
&\hat{M}\mathcal{J}_k\tau = \mathcal{M}(\mathcal{J}_k \otimes \mathrm{Id})\Delta^M\tau, \quad \hat{M}\overline{\mathcal{J}_k\tau} = \overline{\hat{M}\mathcal{J}_k\tau},
\end{aligned}$$

and

$$(\mathrm{Id} \otimes \mathcal{M})(\Delta \otimes \mathrm{Id})\Delta^M\tau = (M \otimes \hat{M})\Delta\tau,$$

since $(\mathrm{Id} \otimes \mathcal{M})(\Delta \otimes \mathrm{Id})$ is invertible. Furthermore, the linear map $\hat{\Delta}^M : T_0^+ \to T_0^+ \otimes T_0^+$ is defined by

$$(\mathcal{A}\hat{M}\mathcal{A} \otimes \hat{M})\Delta^+ = (\mathrm{Id} \otimes \mathcal{M})(\Delta^+ \otimes \mathrm{Id})\hat{\Delta}^M,$$

since $(\mathrm{Id} \otimes \mathcal{M})(\Delta^+ \otimes \mathrm{Id})$ is invertible ([Hai14, Proposition 8.36]).

Theorem 2.20 ([Hai14, Theorem 8.44])**.** *Consider $\mathcal{F}_0$ and M as above. Assume that for every $\tau \in \mathcal{F}_0$ and $\hat{\tau} \in T_0^+$ we can write*

$$\Delta^M\tau = \tau \otimes \mathbf{1} + \sum_{|\tau^{(1)}|_s > |\tau|_s} \tau^{(1)} \otimes \tau^{(2)}, \quad \hat{\Delta}^M\hat{\tau} = \hat{\tau} \otimes \mathbf{1} + \sum_{|\hat{\tau}^{(1)}|_s > |\hat{\tau}|_s} \hat{\tau}^{(1)} \otimes \hat{\tau}^{(2)}.$$

Then for every admissible model $(\mathbf{\Pi}, f)$ on $\mathcal{T}_{\mathrm{cgl}}^{(r)}$, the maps $\mathbf{\Pi}^M$ and f_z^M defined by

$$\mathbf{\Pi}^M = \mathbf{\Pi}M, \quad f_z^M = f_z\hat{M}$$

are uniquely extended to an admissible model $Z^M = (\Pi^M, \Gamma^M)$ on $\mathcal{T}_{\mathrm{cgl}}^{(r)}$.

Now we give a renormalization map M in a concrete form. In order to simplify notations, we introduce a graphical notation for the element in $\mathcal{F}$. First, we draw a circle to represent Ξ. For an element $\mathcal{I}\tau$, we draw a downward black line starting at the root of τ. For a product $\tau\tau'$, we joint these trees at

their roots. The complex conjugate $\overline{\tau}$ is denoted by changing the color black and white to each other. For example,

$$\mathcal{I}\Xi = \mathbf{I}, \quad \mathcal{I}\Xi^2 = \mathbf{V}, \quad \mathcal{I}\Xi^2\overline{\mathcal{I}\Xi} = \mathbb{V}, \quad \mathcal{I}\,\mathbb{V} = \mathbb{Y}, \quad \overline{\mathbb{Y}}\,\mathbf{V} = \mathbb{W}.$$

Then we can list all of elements with negative homogeneities as follows:

Homogeneity	Symbol
α	Ξ
$3(\alpha+2)$	$\mathbb{V}$
$2(\alpha+2)$	$\mathbf{V}$, $\mathbb{V}$
$5\alpha+12$	$\mathbb{W}$, $\mathbb{W}$
$\alpha+2$	$\mathbf{I}$, $\mathbb{I}$
$4\alpha+10$	$\mathbb{Y}$, $\mathbb{Y}$, $\mathbb{Y}$, $\mathbb{Y}$, $\mathbb{W}$, $\mathbb{W}$, $\mathbb{W}$
$2\alpha+5$	$X_i\mathbf{V}$, $X_i\mathbb{V}$ $(i=1,2,3)$
0	$\mathbf{1}$

Since $\alpha > -\frac{18}{7}$, the element $\mathbb{W}$ has positive homogeneity $7\alpha+18>0$, so that it does not appear here.

Considering chaos expansions of Gaussian models as in Section 2.5, it is suitable to define the renormalization map $M = M(C_1, C_{2,1}, C_{2,2})$ by

$$\begin{aligned}
M\mathbf{V} &= \mathbf{V},\\
M\mathbb{V} &= \mathbb{V} - C_1\mathbf{1},\\
M\mathbb{Y} &= \mathbb{Y} - 2C_1\mathbf{I},\\
M\mathbb{W} &= \mathbb{Y}(\mathbb{V} - C_1\mathbf{1}) = \mathbb{W} - C_1\mathbb{Y},\\
M\mathbb{W} &= \mathbb{W} - 2C_{2,1}\mathbf{1},\\
M\mathbb{W} &= \mathbb{Y}(\mathbb{V} - C_1\mathbf{1}) - C_{2,2}\mathbf{1} = \mathbb{W} - C_1\mathbb{Y} - C_{2,2}\mathbf{1},\\
M\mathbb{W} &= \mathbb{W},\\
M\mathbb{Y} &= (\mathbb{Y} - 2C_1\mathbb{I})\mathbf{I} = \mathbb{Y} - 2C_1\mathbb{I},\\
M\mathbb{Y} &= (\mathbb{Y} - 2C_1\mathbb{I})\mathbb{I} = \mathbb{Y} - 2C_1\mathbb{I},\\
M\mathbb{Y} &= (\mathbb{Y} - 2\overline{C_1}\mathbb{I})\mathbf{I} = \mathbb{Y} - 2\overline{C_1}\mathbb{I},\\
M\mathbb{W} &= (\mathbb{Y} - 2C_1\mathbb{I})(\mathbb{V} - C_1\mathbf{1}) - 2C_{2,2}\mathbf{I}\\
&= \mathbb{W} - 2C_1\mathbb{W} - C_1(\mathbb{Y} - 2C_1\mathbb{I}) - 2C_{2,2}\mathbf{I},\\
M\mathbb{W} &= (\mathbb{Y} - 2\overline{C_1}\mathbb{I})\,\mathbf{V} - 2C_{2,1}\mathbf{I} = \mathbb{W} - 2\overline{C_1}\mathbb{W} - 2C_{2,1}\mathbf{I},\\
MX_i\mathbf{V} &= X_i\mathbf{V},\\
MX_i\mathbb{V} &= X_i(\mathbb{V} - C_1\mathbf{1}) = X_i\mathbb{V} - C_1X_i
\end{aligned}$$

for some constants $C_1, C_{2,1}$ and $C_{2,2}$. Since M must be closed in the space $\langle \mathcal{F}_0 \rangle$, we should choose $\mathcal{F}_0$ by

$$\mathcal{F}_0 = \{\Xi, \Psi, \mathbf{V}, \mathbb{V}, \Psi\!\!\Psi, \Psi\!\!\Psi\!\!\Psi, \mathbf{I}, \mathbb{I}, \Psi\!\!\mathbf{I}, \Psi\!\!\mathbb{I}, \Psi\!\!\Psi\!\!\mathbf{I}, \Psi\!\!\Psi\!\!\mathbb{I}, \Psi\!\!\Psi\!\!\Psi\!\!\mathbf{I}, \Psi\!\!\Psi\!\!\Psi\!\!\mathbb{I}, X_i\mathbf{V}, X_i\mathbb{V}, \mathbf{1},$$
$$\Psi\!\!\Psi, \Psi\!\!\Psi, \Psi\!\!\Psi, \Psi\!\!\Psi, \Psi\!\!\Psi, \Psi\!\!\Psi, \Psi\!\!\Psi, \Psi\!\!\Psi, X_i\mathbf{I}, X_i\mathbb{I}, \mathcal{I}\,; i = 1,2,3\}.$$

Then it turns out that we can take $\mathcal{F}_* = \{\mathbf{I}, \mathbb{I}, \mathbf{V}, \mathbb{V}, \Psi\}$. From now on, the subscript i of X_i runs over $\{1,2,3\}$.

Lemma 2.21. *The linear map M satisfies the conditions of Theorem* 2.20*. Furthermore, the identity*

$$(\Pi_z^M \tau)(z) = (\Pi_z M\tau)(z) \tag{2.7}$$

holds for every $\tau \in \mathcal{F}_0$ and $z \in \mathbf{R}^4$.

Proof. Calculations of $\hat{M}$, Δ^M and $\hat{\Delta}^M$ are completely parallel to those in [Hai14, Section 9.2], so here we show only the results. Indeed we have

$$\hat{M}\mathcal{J}\tau = \mathcal{J}M\tau, \quad \hat{M}\mathcal{J}_k\tau = \mathcal{J}_k\tau, \quad (|k|_s > 0)$$

and

$$\begin{aligned}
\Delta^M \Psi &= M\Psi \otimes \mathbf{1} + 2C_1 X_i \otimes \mathcal{J}_i \mathbf{I},\\
\Delta^M \Psi\!\!\mathbf{I} &= M\Psi\!\!\mathbf{I} \otimes \mathbf{1} + 2C_1 X_i \mathbf{I} \otimes \mathcal{J}_i \mathbf{I},\\
\Delta^M \Psi\!\!\mathbb{I} &= M\Psi\!\!\mathbb{I} \otimes \mathbf{1} + 2C_1 X_i \mathbb{I} \otimes \mathcal{J}_i \mathbf{I},\\
\Delta^M \Psi\!\!\Psi\!\!\mathbf{I} &= M\Psi\!\!\Psi\!\!\mathbf{I} \otimes \mathbf{1} + 2\overline{C_1} X_i \mathbf{I} \otimes \overline{\mathcal{J}_i \mathbf{I}},\\
\Delta^M \Psi\!\!\Psi\!\!\mathbb{I} &= M\Psi\!\!\Psi\!\!\mathbb{I} \otimes \mathbf{1} + 2C_1 X_i (\mathbb{V} - C_1\mathbf{1}) \otimes \mathcal{J}_i \mathbf{I},\\
\Delta^M \Psi\!\!\Psi\!\!\Psi\!\!\mathbf{I} &= M\Psi\!\!\Psi\!\!\Psi\!\!\mathbf{I} \otimes \mathbf{1} + 2\overline{C_1} X_i \mathbf{V} \otimes \overline{\mathcal{J}_i \mathbf{I}},\\
\Delta^M \tau &= M\tau \otimes \mathbf{1} \quad \text{(otherwise)}.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
\hat{\Delta}^M \mathcal{J}\Psi &= \mathcal{J}M\Psi + 2C_1(X_i \otimes \mathcal{J}_i\mathbf{I} - X_i\mathcal{J}_i\mathbf{I} \otimes \mathbf{1}),\\
\hat{\Delta}^M \tau &= \hat{M}\tau \otimes \mathbf{1} \quad \text{(otherwise)}.
\end{aligned}$$

(Here and in what follows, summation symbols over the repeated index i are omitted.) Therefore, M satisfies the conditions of Theorem 2.20. The relation (2.7) is obtained by

$$\Pi_z^M \tau = (\Pi_z \otimes f_z)\Delta^M \tau$$

([Hai14, equation (8.34)]) and the fact that $\Pi_z X_i \tau(z) = 0$ for every τ with $X_i\tau \in \mathcal{F}$. □

Proposition 2.22. *Let $Z^\epsilon = (\Pi^\epsilon, \Gamma^\epsilon)$ be a model canonically lifted from a continuous function ξ^ϵ. Let $S : (u_0, Z) \mapsto u$ be the solution map given by Theorem 2.19. Given constants $C_1, C_{2,1}$ and $C_{2,2}$, denote by $\hat{Z}^\epsilon = (\hat{\Pi}^\epsilon, \hat{\Gamma}^\epsilon)$ the renormalized model given by Theorem 2.20. Then for every periodic $u_0 \in \mathcal{C}^\eta$, $\hat{u}^\epsilon = \mathcal{R}S(u_0, \hat{Z}^\epsilon)$ solves the equation*

$$\partial_t \hat{u}^\epsilon = (\mathsf{i} + \mu)\triangle \hat{u}^\epsilon + \nu(1 - |\hat{u}^\epsilon|^2 + (2C_1 - 2\overline{\nu}C_{2,1} - 4\nu C_{2,2}))\hat{u}^\epsilon + \xi^\epsilon. \tag{2.8}$$

Proof. Since the fixed point problem (2.6) can be written by $u = \mathcal{I}F(u) + \cdots$, where $\cdots$ takes values in T^{poly}, we can find functions φ and $\{\varphi^i\}_{i=1}^3$ such that the solution $u \in \mathcal{D}_P^{\gamma,\eta}$ of (2.6) with $\gamma = 1^+$ (greater than but sufficiently close to 1) can be written by

$$u = \mathbf{I} + \varphi\mathbf{1} - \nu\,\mathbf{I}\!\mathbf{V} - 2\nu\varphi\,\mathbf{I}\!\mathbf{V} - \nu\overline{\varphi}\,\mathbf{I}\!\mathbf{V} + \varphi^i X_i.$$

In particular, since $\Pi_z \mathbf{I} = \Pi_z^M \mathbf{I} = K * \xi^\epsilon$ we have

$$\mathcal{R}u = \mathcal{R}^M u = K * \xi^\epsilon + \varphi.$$

On the other hand, by Proposition 2.11, $\hat{u}^\epsilon = \mathcal{R}^M u$ satisfies the equation

$$\hat{u}^\epsilon(t,x) = \int_0^t \int_{\mathbf{R}^4} G(t-s, x-y)(\mathcal{R}^M F(u))(s,y)dsdy + \int_{\mathbf{R}^4} G(t, x-y)u_0(y)dy.$$

Hence it suffices to show that $\mathcal{R}^M F(u)$ coincides with the driving terms of (2.8). We can expand $F(u) = \Xi + \nu(1 - u\overline{u})u$ up to homogeneity 0^+ as follows.

$$\begin{aligned} F(u) = \Xi &- \nu\,\mathbf{V}\!\mathbf{V} - 2\nu\varphi\,\mathbf{V}\!\mathbf{V} - \nu\overline{\varphi}\,\mathbf{V}\!\mathbf{V} + 2\nu^2\,\mathbf{V}\!\mathbf{V}\!\mathbf{V} + \nu\overline{\nu}\,\mathbf{V}\!\mathbf{V}\!\mathbf{V} \\ &+ \nu(1 - 2\varphi\overline{\varphi})\,\mathbf{I} - \nu\varphi^2\,\mathbb{I} + 2\nu^2\overline{\varphi}\,\mathbf{V}\!\mathbf{V} + 2\nu^2\varphi\,\mathbf{V}\!\mathbf{V} + 2\nu\overline{\nu}\varphi\,\mathbf{V}\!\mathbf{V} \\ &+ 2\nu^2\overline{\varphi}\,\mathbf{V}\!\mathbf{V} + \nu\overline{\nu}\varphi\,\mathbf{V}\!\mathbf{V} + 4\nu^2\varphi\,\mathbf{V}\!\mathbf{V} + 2\nu\overline{\nu\varphi}\,\mathbf{V}\!\mathbf{V} \\ &- \nu\overline{\varphi^i}X_i\,\mathbf{V} - 2\nu\varphi^i X_i\,\mathbf{V} + \nu(\varphi - \varphi^2\overline{\varphi})\mathbf{1}. \end{aligned}$$

Since $\mathcal{R}^M u = \mathcal{R}Mu$ follows from (2.7), we have

$$\begin{aligned} \mathcal{R}^M F(u) &= \mathcal{R}F(u) + 2\nu C_1 K * \xi^\epsilon + 2\nu C_1\varphi \\ &\quad - 4\nu^2 C_{2,2} K * \xi^\epsilon - 2\nu\overline{\nu}C_{2,1}K * \xi^\epsilon - 2\nu\overline{\nu}C_{2,1}\varphi - 4\nu^2 C_{2,2}\varphi \\ &= \xi^\epsilon + \nu(1 - \mathcal{R}u\overline{\mathcal{R}u})\mathcal{R}u + 2\nu C_1\mathcal{R}u - 2\nu\overline{\nu}C_{2,1}\mathcal{R}u - 4\nu^2 C_{2,2}\mathcal{R}u \\ &= \xi^\epsilon + \nu(1 - |\hat{u}^\epsilon|^2 + (2C_1 - 2\overline{\nu}C_{2,1} - 4\nu C_{2,2}))\hat{u}^\epsilon. \end{aligned}$$

This completes the proof. □

2.5 Convergence of Gaussian models

Our goal is to show the following renormalization result. We give its proof in the next section since it takes long.

Proposition 2.23. *If we choose $C_1^\epsilon, C_{2,1}^\epsilon$ and $C_{2,2}^\epsilon$ as in* (3.4)*, then there exists a random model $\hat{Z}$ independent of the choice of ρ, and for every $\theta \in (0, -\frac{5}{2} - \alpha)$, $\gamma < r$, $p > 1$ and every compact set $K \subset \mathbf{R}^4$, we have the bounds*

$$\boldsymbol{E}[|\!|\!|\hat{Z}^\epsilon|\!|\!|_{\gamma;K}^p] \lesssim 1, \quad \boldsymbol{E}[|\!|\!|\hat{Z}^\epsilon; \hat{Z}|\!|\!|_{\gamma;K}^p] \lesssim \epsilon^{\theta p}. \tag{2.9}$$

Furthermore, for every $T > 0$ we have

$$\boldsymbol{E}[|\!|\xi^\epsilon|\!|_{\alpha;K}^p] \lesssim 1, \quad \boldsymbol{E}[|\!|\xi^\epsilon - \xi|\!|_{\alpha;K}^p] \lesssim \epsilon^{\theta p}, \tag{2.10}$$

and

$$\boldsymbol{E}[\|K * \xi^\epsilon\|_{C([0,T],\mathcal{C}^{\alpha+2})}^p] \lesssim 1, \quad \boldsymbol{E}[\|K * \xi^\epsilon - K * \xi\|_{C([0,T],\mathcal{C}^{\alpha+2})}^p] \lesssim \epsilon^{\theta p}. \tag{2.11}$$

Combining Propositions 2.22 and 2.23, we obtain Theorem 2.1 if we choose $C^\epsilon = 2C_1^\epsilon - 2\overline{\nu}C_{2,1}^\epsilon - 4\nu C_{2,2}^\epsilon$.

3 Proof of convergence of renormalized models

In this section, we give a proof of Proposition 2.23. Since the estimates (2.10) and (2.11) are obtained in [Hai14, Proposition 9.5], we focus on the estimate (2.9). By [Hai14, Theorem 10.7], it suffices to show that there exist $\kappa, \theta > 0$ such that, for every $\tau \in \mathcal{F}$ with $|\tau|_s < 0$, every test function $\varphi \in \mathcal{B}_r$ and $z \in \mathbf{R}^4$, there exists a random variable $\langle \hat{\Pi}_z \tau, \varphi \rangle$ such that

$$\boldsymbol{E}[|\langle \hat{\Pi}_z \tau, \varphi_z^\delta \rangle|^2] \lesssim \delta^{2|\tau|_s + \kappa}, \quad \boldsymbol{E}[|\langle \hat{\Pi}_z \tau - \hat{\Pi}_z^\epsilon \tau, \varphi_z^\delta \rangle|^2] \lesssim \epsilon^{2\theta} \delta^{2|\tau|_s + \kappa}. \tag{3.1}$$

We fix $z \in \mathbf{R}^4$ throughout this section. The estimates in this section are uniform over z.

This section is organized as follows. In Section 3.1, we recall the Wiener chaos decomposition of the random variable $\hat{\Pi}_z \tau$ and introduce graphical notations to describe its kernel. In Section 3.2, we give some useful estimates to prove (3.1). In Section 3.3, we show the required estimate (3.1) for each symbol τ. In Section 3.4, we show the explicit forms of renormalization constants and their divergence orders.

3.1 Wiener chaos decomposition

The driving noise ξ is space-time white noise on $\mathbf{R} \times \mathbf{T}^3$, which is extended periodically to $\mathbf{R}^4$. In precise, we are given the complex multiple Wiener integral $\mathcal{J}_{p,q}$ on $(E, m) = (\mathbf{R} \times \mathbf{T}^3, dtdx_1dx_2dx_3)$ (see Section A) and a random distribution ξ is defined by $\langle \xi, \varphi \rangle = \mathcal{J}_{1,0}(\pi\varphi)$, where φ is a compactly supported smooth function and $\pi\varphi = \sum_{n \in \mathbf{Z}^3} S_n\varphi$ is its periodic extension, where $S_n\varphi(t,x) = \varphi(t, x+n)$. Although $\mathcal{J}_{1,0}$ is an isometry from $L^2(E)$ (not $L^2(\mathbf{R}^4)$) to $L^2(\Omega)$, when φ is supported in $\mathbf{R} \times [-\frac{1}{2}, \frac{1}{2}]^3$ (i.e. φ and $S_n\varphi$ have disjoint supports if $n \neq 0$) we have the isometry

$$\boldsymbol{E}[|\langle \xi, \varphi \rangle|^2] = \|\pi\varphi\|_E^2 = \int_{\mathbf{R}^4} |\varphi(z)|^2 dz. \tag{3.2}$$

The approximation $\xi^\epsilon(z) = \xi * \rho^\epsilon(z) = \mathcal{J}_{1,0}(\rho^\epsilon(z - \cdot))$ belongs to the first Wiener chaos. By definition and the product formula, for each $\tau \in \mathcal{F}$ we have the Wiener chaos decomposition

$$\hat{\Pi}_z^\epsilon \tau(z') = \sum_{p,q} \mathcal{J}_{p,q}(\hat{\mathcal{W}}_z^{\epsilon,(p,q)} \tau(z')),$$

where $\hat{\mathcal{W}}_z^{\epsilon,(p,q)} \tau(z') \in L^2_{p,q}$. In all these kernels mentioned below, we always assume that $\hat{\mathcal{W}}_z^{\epsilon,(p,q)} \tau(z')$ is supported in sufficiently small compact subset as a function of $(\mathbf{R}^4)^{p+q}$, so that we need not distinguish integrals on $\mathbf{R} \times \mathbf{T}^3$ and $\mathbf{R}^4$, indeed similarly to (3.2)

$$\begin{aligned}\boldsymbol{E}[|\langle \hat{\Pi}_z \tau, \varphi_z^\delta \rangle|^2] &\le p!q! \left\| \int \varphi_z^\delta(z') \hat{\mathcal{W}}_z^{\epsilon,(p,q)} \tau(z') dz' \right\|^2_{L^2_{pq}} \\ &= p!q! \iint \varphi_z^\delta(z') \varphi_z^\delta(z'') \langle \hat{\mathcal{W}}_z^{\epsilon,(p,q)} \tau(z'), \overline{\hat{\mathcal{W}}_z^{\epsilon,(p,q)} \tau(z'')} \rangle_{L^2((\mathbf{R}^4)^{p+q})} dz' dz''.\end{aligned}$$

This assumption is satisfied if we take the support of K sufficiently small.

Following [Hai14, Section 10.5], we introduce the following graphical notations to write integrated kernels. First a dot represents a variable in $\mathbf{R}^4$. A square dot ($\underset{z'}{\blacksquare}$) represents a fixed variable. A gray dot ($\color{gray}\bullet$) is a variable integrated out on $\mathbf{R}^4$, so it has no label. A variable representing the multiple Wiener integral $\mathcal{J}_{p,q}$ is written by a black dot ($\bullet$) for a variable in E^p, and a white dot ($\circ$) for a variable in E^q, respectively. Second an arrow represents a function of two variables which are represented by its vertices. We write

$K(z'-z'') =$ $z' \blacksquare \longleftarrow \blacksquare z''$, $\quad$ $K^\epsilon(z'-z'') =$ $z' \blacksquare \leftarrow\!\cdots\!\blacksquare z''$,

$\overline{K}(z'-z'') =$ $z' \blacksquare \Longleftarrow \blacksquare z''$, $\quad$ $\overline{K^\epsilon}(z'-z'') =$ $z' \blacksquare \leftarrow\!\cdots\!\blacksquare z''$,

where $K^\epsilon = K * \rho^\epsilon$. Moreover, we write

$$K(z'-z'') - K(z-z'') = \ z' \blacksquare \longleftarrow\!\!\!\!\!\!+ \blacksquare z'' \ ,$$
$$\overline{K}(z'-z'') - \overline{K}(z-z'') = \ z' \blacksquare \longleftarrow\!\!\!\!\!\!= \blacksquare z'' \ .$$

We note that $z \in \mathbf{R}^4$ is fixed. We write several kernels by combining these notations. For example,

$$\mathcal{J}_{3,1}(\ z' \blacksquare \longleftarrow\!\!\!\!\!\!+ \bullet \longleftarrow \bullet \ )$$
$$= \int \left\{ \int (K(z'-u) - K(z-u)) K^\epsilon(u-w_1) K^\epsilon(u-w_2) \overline{K^\epsilon}(u-w') du \right\}$$
$$\times K(z'-w_3) : \xi(dw_1)\xi(dw_2)\xi(dw_3)\overline{\xi}(dw') : .$$

3.2 Estimates of singularity of kernels

From the scaling property of $K = \sum_n K_n$, we can see that $|K(z)| \lesssim \|z\|_s^{-3}$. It is useful to consider the singularity of kernels like this. The notation $A(z'-z'') =$ $z' \blacksquare \boxed{-\alpha} \blacksquare z''$ implies that $A : \mathbf{R}^4 \setminus \{0\} \to \mathbf{C}$ is a smooth function supported in a ball and has the estimate $|A(z'-z'')| \lesssim \|z'-z''\|_s^{-\alpha}$. We recall some useful estimates from [Hai14, Section 10.3] and [Hos16a, Section 4.7].

Lemma 3.1 ([Hai14, Lemma 10.14])**.** *For $\alpha, \beta \in [0,5)$, we have*

$$| \ z' \blacksquare \boxed{-\alpha} \bullet \boxed{-\beta} \blacksquare z'' \ | \lesssim \begin{cases} z' \blacksquare \boxed{-\alpha-\beta+5} \blacksquare z'' \ , & \alpha+\beta > 5, \\ 1, & \alpha+\beta < 5. \end{cases}$$

Lemma 3.2 ([Hos16a, Lemma 4.31])**.** *Let $\alpha, \beta, \alpha', \beta', \gamma \in (0,5)$. If $\zeta \in (0, \alpha \wedge \beta]$ and $\eta \in (0, \alpha' \wedge \beta']$ satisfy*

$$\alpha+\beta-5 < \zeta, \quad \alpha'+\beta'-5 < \eta, \quad \alpha+\beta+\alpha'+\beta'+\gamma-10 < \zeta+\eta,$$

then we have

$$\left| \begin{array}{l} z' \blacksquare \boxed{-\alpha} \\ w' \blacksquare \boxed{-\beta} \end{array} \bullet \boxed{-\gamma} \bullet \begin{array}{l} \boxed{-\alpha'} \blacksquare z'' \\ \boxed{-\beta'} \blacksquare w'' \end{array} \right| \lesssim \begin{array}{l} z' \blacksquare \boxed{-\zeta} \blacksquare w' \\ z'' \blacksquare \boxed{-\eta} \blacksquare w'' \end{array} .$$

For $\alpha, \beta \geq 0$, we use the notation

$$Q_{\alpha,\beta}(z', z'') = \ z' \blacksquare \boxed{-\alpha} \bullet \boxed{-\beta} \bullet \boxed{-\alpha} \blacksquare z'' \ .$$

Lemma 3.3. *Let $\alpha, \beta \in [0,5)$. For $\theta \in (0, 1 \wedge (2 - \alpha - \frac{\beta}{2}))$, we have*

$$|Q_{\alpha,\beta}(z', z'')| \lesssim \|z'-z\|_s^\theta \|z''-z\|_s^\theta.$$

Proof. Since $|$  $|\lesssim \|z'-z\|_s^\theta(\|z'-u\|_s^{-3-\theta}+\|z-u\|_s^{-3-\theta})$ by [Hai14, Lemma 10.18],

$$|Q_{\alpha,\beta}(z',z'')|\lesssim \|z'-z\|_s^\theta\|z''-z\|_s^\theta(R(z',z'')+R(z,z'')+R(z',z)+R(z,z)),$$

where

$$R(u',u'')=R_{z',z''}(u',u'')=$$ 

It suffices to show that R is bounded. By the inequality $\|z'\|_s^{-\alpha}\|z''\|_s^{-\beta}\lesssim \|z'\|_s^{-\alpha-\beta}+\|z''\|_s^{-\alpha-\beta}$ for $\alpha,\beta\geq 0$, the function R is bounded by the sum of functions of the form



which is bounded by 1 since $2(-\alpha-3-\theta)-\beta+10>0$. □

3.3 Proof of L^2-estimates (3.1)

Now we prove the estimate (3.1) for every $\tau\in\mathcal{F}$ with $|\tau|_s<0$.

3.3.1 $\Xi, \mathbf{I}, \mathbb{I}, \mathbf{V}, \mathbb{V}, \mathbb{V}^{\circ}$

For $\tau=\Xi, \mathbf{I}, \mathbb{I}$, the required estimates follow from [Hai14, Proposition 9.5]. We now treat $\tau=\mathbf{V}, \mathbb{V}, \mathbb{V}^{\circ}$. By definition,

$$\hat{\Pi}_z^\epsilon \mathbf{V}=\Pi_z^\epsilon \mathbf{V}=(\Pi_z^\epsilon \mathbf{I})^2,$$
$$\hat{\Pi}_z^\epsilon \mathbb{V}=\Pi_z^\epsilon \mathbb{V}-C_1^\epsilon=(\Pi_z^\epsilon \mathbf{I})(\Pi_z^\epsilon \mathbb{I})-C_1^\epsilon.$$

By applying the product formula to

$$\Pi_z^\epsilon \mathbf{I}(z')=K*\xi^\epsilon(z')=\mathcal{J}_{1,0}(\ z'$$  $$),$$
$$\Pi_z^\epsilon \mathbb{I}(z')=\overline{K*\xi^\epsilon}(z')=\mathcal{J}_{0,1}(\ z'$$  $$),$$

we have

$$\hat{\Pi}_z^\epsilon \mathbf{V}(z')=\mathcal{J}_{2,0}(\ z'$$  $$),$$

$$\hat{\Pi}_z^\epsilon \mathbb{V}(z')=\mathcal{J}_{1,1}(\ z'$$  $$)+\ z'$$ $$-C_1^\epsilon.$$

If we choose $C_1^\epsilon=\ z'$  $=\int|K^\epsilon(z)|^2dz$, the required estimates (2.9) for $\tau=\mathbf{V}, \mathbb{V}$ easily follow. Indeed,

$$\boldsymbol{E}[|\langle\hat{\Pi}_z^\epsilon \mathbf{V},\varphi_z^\delta\rangle|^2]=2\iint\varphi_z^\delta(z')\varphi_z^\delta(z'')\ z'$$  $$z''\ dz'dz''$$

$$\lesssim \iint \varphi_z^\delta(z')\varphi_z^\delta(z'') \left| \; z' \blacksquare \; \begin{matrix} -1 \\ -1 \end{matrix} \; \blacksquare \, z'' \; \right| dz'dz''$$
$$\lesssim \iint \varphi_z^\delta(z')\varphi_z^\delta(z'')\|z'-z''\|_s^{-2}dz'dz'' \lesssim \delta^{-2}$$

and $-2 > |\mathbf{V}|_s = 4(\alpha+2)$. Moreover, if we choose

$$\hat{\Pi}_z \mathbf{V}(z') = \mathcal{J}_{2,0}(\; z' \blacksquare \; \begin{matrix} \bullet \\ \bullet \end{matrix} \;),$$

then we have

$$\boldsymbol{E}[|\langle \hat{\Pi}_z^\epsilon \mathbf{V} - \hat{\Pi}_z \mathbf{V}, \varphi_z^\delta \rangle|^2] \lesssim \epsilon^\theta \delta^{-2-\theta}$$

for small $\theta > 0$. This is obtained by similar argument, since $|(K^\epsilon - K)(z)| \lesssim \epsilon^\theta \|z\|_s^{-3-\theta}$ for $\theta \in (0,1]$ (see [Hai14, Lemma 10.17]). The case $\tau = \mathbb{V}$ is similar.

For $\tau = \mathbf{V}$, by the choice of C_1^ϵ we have

$$\hat{\Pi}_z^\epsilon \mathbf{V}(z') = (\Pi_z^\epsilon \mathbf{I})(z')^2 (\Pi_z^\epsilon \mathbb{I})(z') - 2C_1^\epsilon (\Pi_z^\epsilon \mathbf{I})(z')$$
$$= \mathcal{J}_{2,1}(\; z' \blacksquare \; \begin{matrix} \bullet \\ \bullet \\ \circ \end{matrix} \;).$$

Then the estimate (2.9) for $\tau = \mathbf{V}$ easily follows as above. Indeed,

$$\boldsymbol{E}[|\langle \hat{\Pi}_z^\epsilon \mathbf{V}, \varphi_z^\delta \rangle|^2] = 2 \iint \varphi_z^\delta(z')\varphi_z^\delta(z'') \; z' \blacksquare \; \begin{matrix} \bullet \\ \bullet \\ \bullet \end{matrix} \; \blacksquare \, z'' \; dz'dz''$$
$$\lesssim \iint \varphi_z^\delta(z')\varphi_z^\delta(z'') \left| \; z' \blacksquare \; \begin{matrix} -1 \\ -1 \\ -1 \end{matrix} \; \blacksquare \, z'' \; \right| dz'dz''$$
$$\lesssim \iint \varphi_z^\delta(z')\varphi_z^\delta(z'')\|z'-z''\|_s^{-3}dz'dz'' \lesssim \delta^{-3}$$

and $-3 > |\mathbf{V}|_s = 6(\alpha+2)$. The estimates for $\hat{\Pi}_z^\epsilon \mathbf{V} - \hat{\Pi}_z \mathbf{V}$ are similarly obtained by choosing

$$\hat{\Pi}_z^\epsilon \mathbf{V}(z') = \mathcal{J}_{2,1}(\; z' \blacksquare \; \begin{matrix} \bullet \\ \bullet \\ \circ \end{matrix} \;).$$

In the subsequent computations, the estimates of $\hat{\Pi}_z^\epsilon \tau - \hat{\Pi}_z \tau$ are obtained by similar arguments to those of $\hat{\Pi}_z^\epsilon \tau$ as above by using the bound of $K^\epsilon - K$, so we show only the uniform boundedness but not the convergence estimates explicitly. For detailed proofs, see [Hos16a, Section 4.8].

3.3.2 $X_i \mathbb{V}, X_i \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}$

For $\tau = X_i \mathbb{V}, X_i \mathbb{V}$, the corresponding estimates are easily obtained. Indeed, since $\hat{\Pi}^\epsilon_z X_i \mathbb{V}(z') = (x'_i - x_i)\hat{\Pi}^\epsilon_z \mathbb{V}(z')$ we have

$$\begin{aligned}
&\boldsymbol{E}[|\langle \hat{\Pi}^\epsilon_z X_i \mathbb{V}, \varphi^\delta_z \rangle|^2] \\
&= 2 \iint \varphi^\delta_z(z')\varphi^\delta_z(z'')(x'_i - x_i)(x''_i - x_i)\ z' \blacksquare \quad \blacksquare\ z''\ dz'dz'' \\
&\lesssim \iint \varphi^\delta_z(z')\varphi^\delta_z(z'')(x'_i - x_i)(x''_i - x_i)\|z' - z''\|_s^{-2} dz'dz'' \lesssim 1
\end{aligned}$$

and $0 > 2|X_i \mathbb{V}|_s = 2(2\alpha + 5)$. The case $\tau = X_i \mathbb{V}$ is similar.

Now we turn to $\mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}$. In particular, we consider the renormalizations of $\mathbb{V}$ and $\mathbb{V}$, since the corresponding chaos decompositions of the two other elements do not have zeroth order terms. By definition,

$$\begin{aligned}
\hat{\Pi}^\epsilon_z \mathbb{V} &= \Pi^\epsilon_z \mathbb{V} - 2C^\epsilon_{2,1} = (\Pi^\epsilon_z \mathbb{V})(\Pi^\epsilon_z \mathbb{V}) - 2C^\epsilon_{2,1}, \\
\hat{\Pi}^\epsilon_z \mathbb{V} &= \Pi^\epsilon_z \mathbb{V} - C^\epsilon_1 \Pi^\epsilon_z \mathbb{V} - C^\epsilon_{2,2} = (\Pi^\epsilon_z \mathbb{V} - C^\epsilon_1)(\Pi^\epsilon_z \mathbb{V}) - C^\epsilon_{2,2}.
\end{aligned}$$

We note that

$$\Pi^\epsilon_z \mathbb{V}(z') = \overline{K * \Pi^\epsilon_z \mathbb{V}}(z') - \overline{K * \Pi^\epsilon_z \mathbb{V}}(z) = \mathcal{J}_{0,2}(\ z' \blacksquare \longleftarrow \bullet \quad \circ \ ),$$

$$\Pi^\epsilon_z \mathbb{V}(z') = K * \Pi^\epsilon_z \mathbb{V}(z') - K * \Pi^\epsilon_z \mathbb{V}(z) = \mathcal{J}_{1,1}(\ z' \blacksquare \longleftarrow \bullet \quad \bullet \ ).$$

By applying the product formula (Theorem A.1) we have

$$\hat{\Pi}^\epsilon_z \mathbb{V}(z') = \mathcal{J}_{2,2}(\ z' \blacksquare \quad \bullet \quad \circ \ ) + 4\mathcal{J}_{1,1}(\ z' \blacksquare \quad \bullet \quad \circ \ )$$

$$+ 2(\ z' \blacksquare \quad \bullet \ - C^\epsilon_{2,1})$$

and

$$\hat{\Pi}^\epsilon_z \mathbb{V}(z') = \mathcal{J}_{2,2}(\ z' \blacksquare \quad \bullet \quad \bullet \ ) + \mathcal{J}_{1,1}(\ z' \blacksquare \quad \bullet \quad \bullet \ )$$

$$+ \mathcal{J}_{1,1}(\ z' \blacksquare \quad \bullet \quad \circ \ ) + \ z' \blacksquare \quad \bullet \ - C^\epsilon_{2,2}.$$

Hence if we choose

$$C^\epsilon_{2,1} = \ z' \blacksquare \quad \bullet \ , \quad C^\epsilon_{2,2} = \ z' \blacksquare \quad \bullet \ ,$$

we have the required bounds. Indeed, since kernels belonging to the same order chaos have the same graphs except for the difference of K and $\overline{K}$, it suffices to show the bounds for one of these kernels for each order chaos. For remaining zeroth order terms, we have the bounds



for an arbitrary small $\kappa > 0$. For the second order terms, by Lemma 3.3 we have



$$= | z' \blacksquare \blacksquare z'' | \times |Q_{1,1}(z', z'')|$$

$$\lesssim \|z' - z\|_s^{\frac{1}{2}-\kappa} \|z'' - z\|_s^{\frac{1}{2}-\kappa} \|z' - z''\|_s^{-1}$$

for small $\kappa > 0$. Similarly, for the fourth order terms, we have



$$= | z' \blacksquare \blacksquare z'' | \times |Q_{0,2}(z', z'')|$$

$$\lesssim \|z' - z\|_s^{1-\kappa} \|z'' - z\|_s^{1-\kappa} \|z' - z''\|_s^{-2}$$

for small $\kappa > 0$. As a consequence, we have

$$\begin{aligned} E|\langle \hat{\Pi}_z^\epsilon \, , \varphi_z^\delta \rangle|^2 \lesssim \iint \varphi_z^\delta(z') \varphi_z^\delta(z'') \{ & \|z' - z\|_s^{-\kappa} \|z'' - z\|_s^{-\kappa} \\ & + \|z' - z\|_s^{\frac{1}{2}-\kappa} \|z'' - z\|_s^{\frac{1}{2}-\kappa} \|z' - z''\|_s^{-1} \\ & + \|z' - z\|_s^{1-\kappa} \|z'' - z\|_s^{1-\kappa} \|z' - z''\|_s^{-2} \} dz' dz'' \lesssim \delta^{-2\kappa} \end{aligned}$$

for an arbitrary small $\kappa > 0$. The cases $\tau = \, , \, , \,$ are similar.

3.3.3 , , , ,

We treat the case $\tau = \,$. The cases $\tau = \, , \,$ are similar. By definition,

$$\begin{aligned} \hat{\Pi}_z^\epsilon \, (z') &= \Pi_z^\epsilon \, (z') - 2C_1^\epsilon (\Pi_z^\epsilon \, (z') - f_z^\epsilon(\mathcal{J}_i \,)(\Pi_z^\epsilon X_i \,)(z')) \\ &= \Pi_z^\epsilon \, (z') \{ K * \Pi_z^\epsilon \, (z') - K * \Pi_z^\epsilon \, (z) \\ &\quad - 2C_1^\epsilon (K * \Pi_z^\epsilon \, (z') - K * \Pi_z^\epsilon \, (z)) \} \\ &= \mathcal{J}_{1,0}(\; z' \blacksquare \bullet \;) \mathcal{J}_{2,1}(\; z' \blacksquare \;) \end{aligned}$$

$= \mathcal{J}_{3,1}($ $) + \mathcal{J}_{2,0}($ $).$

The summation symbol over $i = 1, 2, 3$ is omitted again. For the fourth order term, by Lemma 3.3 we have



$$= |\ z' \blacksquare \cdots \blacksquare z''\ | \times |Q_{0,3}(z', z'')|$$

$$\lesssim \|z' - z\|_s^{\frac{1}{2}-\kappa} \|z'' - z\|_s^{\frac{1}{2}-\kappa} \|z' - z''\|_s^{-1}$$

for small $\kappa > 0$. For the second order term, we decompose it as

(3.3) 

By Schwarz's inequality, it suffices to consider the bound for each term. For the first term, we have



for small $\kappa > 0$. For the second term, by Lemma 3.2 we have



$$\lesssim \|z' - z\|_s^{-\kappa} \|z'' - z\|_s^{-\kappa}$$

for small $\kappa > 0$. As a consequence, we have

$$E|\langle \hat{\Pi}_z^\epsilon \Psi, \varphi_z^\delta \rangle|^2 \lesssim \delta^{-2\kappa}.$$

Finally we treat $\tau = \Psi$. The other one is similar. By definition,

$$\begin{aligned}
\hat{\Pi}_z^\epsilon \Psi(z') &= \Pi_z^\epsilon \{ (\Psi - 2C_1^\epsilon \mathbf{I})(\mathbb{V} - C_1^\epsilon \mathbf{1}) - 2C_{2,2}^\epsilon \mathbf{I} \}(z') \\
&\quad + 2C_1^\epsilon f_z^\epsilon (\mathcal{J}_i \mathbf{I}) \Pi_z^\epsilon \{ X_i (\mathbb{V} - C_1^\epsilon \mathbf{1}) \}(z') \\
&= \Pi_z^\epsilon (\mathbb{V} - C_1^\epsilon)(z') \{ K * \Pi_z^\epsilon \mathbb{V}(z') - K * \Pi_z^\epsilon \mathbb{V}(z) \\
&\quad - 2C_1^\epsilon (K * \Pi_z^\epsilon \mathbf{I}(z') - K * \Pi_z^\epsilon \mathbf{I}(z)) \}(z') - 2C_{2,2}^\epsilon \Pi_z^\epsilon \mathbf{I}(z') \\
&= \mathcal{J}_{1,1}(\ z' \blacksquare \cdots \bullet_{\circ} \ ) \mathcal{J}_{2,1}(\ z' \blacksquare \longleftarrow \bullet \bullet_{\circ} \ )
\end{aligned}$$



For the fifth order term, we have



for small $\kappa > 0$. For the third order terms, we note that the required bounds are obtained by multiplying  to (3.3), so we have the bound

$$(\|z' - z''\|_s^{-2\kappa} + \|z' - z\|_s^{-\kappa}\|z'' - z\|_s^{-\kappa})\|z' - z''\|_s^{-1}.$$

For the first order terms, we need to introduce the renormalization



where $L^\epsilon(z' - w) =$  and $\mathfrak{R}L^\epsilon$ is the distribution defined by

$$\langle \mathfrak{R}L^\epsilon, \varphi \rangle = \int L^\epsilon(z)(\varphi(z) - \varphi(0))dz$$

for test function φ, see [Hai14, Definition 10.15]. By [Hai14, Lemma 10.16], we have the bound

$$|\mathfrak{R}L^\epsilon * K^\epsilon(z)| \lesssim \|z\|_s^{-3}.$$

For the remaining term, we have



$$\lesssim \|z' - z\|_s^{-\frac{1}{2}-\kappa} \|z'' - z\|_s^{-\frac{1}{2}-\kappa}.$$

As a consequence, we have

$$\boldsymbol{E}[|\langle \hat{\Pi}_z^\epsilon \, \varphi_z^\delta \rangle|^2] \lesssim \delta^{-1-2\kappa}.$$

3.4 Behaviors of renormalization constants

In Sections 3.3.1 and 3.3.2, we obtained renormalization constants

$$C_1^\epsilon = \int |K^\epsilon(z)|^2 dz,$$

$$C_{2,1}^\epsilon = \int \overline{K}(z)(Q^\epsilon(z))^2 dz, \tag{3.4}$$

$$C_{2,2}^\epsilon = \int K(z)|Q^\epsilon(z)|^2 dz,$$

where

$$Q^\epsilon(z) = \int K^\epsilon(z-w)\overline{K^\epsilon}(-w)dw.$$

Note that $Q^\epsilon = Q * \pi^\epsilon$, where $\pi(t,x) = \epsilon^{-5}\pi(\epsilon^{-2}t, \epsilon^{-1}x)$ and

$$Q(z) = \int K(z-w)\overline{K}(-w)dw, \quad \pi(z) = \int \rho(z-w)\rho(-w)dw,$$

Proposition 3.4. *There exist constants $C_1, C_{2,1}$ and $C_{2,2}$ independent of ϵ such that*

$$C_1^\epsilon \sim C_1\epsilon^{-1}, \quad C_{2,1}^\epsilon \sim C_{2,1}\log\epsilon^{-1}, \quad C_{2,1}^\epsilon \sim C_{2,2}\log\epsilon^{-1}$$

as $\epsilon \downarrow 0$. Here for two functions A_ϵ and B_ϵ of ϵ, we write $A_\epsilon \sim B_\epsilon$ if there exists a constant C independent of ϵ and $|A_\epsilon - B_\epsilon| \le C$ holds.

In order to prove the above estimates, we prepare some notations. For $\alpha > 0$ and a compactly supported function $A \in C^\infty(\mathbf{R}^4 \setminus \{0\}, \mathbf{C})$, we say that $A \in \mathfrak{S}_\alpha$ if there exists a function $\widetilde{A} \in C^\infty(\mathbf{R}^4 \setminus \{0\}, \mathbf{C})$ and $A_- \in L^\infty(\mathbf{R}^4, \mathbf{C})$ such that

- $\widetilde{A} = A + A_-$ on $\mathbf{R}^4 \setminus \{0\}$,
- $\widetilde{A}(\lambda^2 t, \lambda x) = \lambda^{-\alpha}\widetilde{A}(t,x)$ for every $\lambda > 0$ and $(t,x) \neq 0$,

- $|A_-(z)| \lesssim \|z\|_s^{-\alpha}$ for every $z \in \mathbf{R}^4$.

The second scaling property of $\widetilde{A}$ ensures that $|\widetilde{A}(z)| \lesssim \|z\|_s^{-\alpha}$ for every $z \in \mathbf{R}^4$ (see [Hai14, Lemma 5.5]).

Proposition 3.5. *Let $\alpha, \beta \in (0,5)$.*

(1) If $A \in \mathfrak{S}_\alpha$, $B \in \mathfrak{S}_\beta$ and $\alpha + \beta > 5$, then $A * B \in \mathfrak{S}_{\alpha+\beta-5}$.

(2) If $A \in \mathfrak{S}_\alpha$ and $B \in \mathfrak{S}_\beta$, then $AB \in \mathfrak{S}_{\alpha+\beta}$.

Proof. For (1), in the decomposition

$$A * B = \widetilde{A} * \widetilde{B} - A_- * \widetilde{B} - \widetilde{A} * B_- + A_- * B_-,$$

we see that the last three terms are bounded by $\|z\|_s^{5-\alpha-\beta}$ by using [Hos16a, Lemma 4.14]. Hence it suffices to set $\widetilde{A * B} = \widetilde{A} * \widetilde{B}$.

The assertion (2) is similarly obtained. □

Proof of Proposition 3.4. First we show the estimate of C_1^ϵ. Since $K, \overline{K} \in \mathfrak{S}_3$, we have $Q \in \mathfrak{S}_1$. Hence we have

$$\begin{aligned} C_1^\epsilon = Q^\epsilon(0) &= \int Q(-z)\pi^\epsilon(z)dz \\ &= \int \widetilde{Q}(-z)\pi^\epsilon(z)dz - \int Q_-(-z)\pi^\epsilon(z)dz \\ &= \epsilon^{-1}\int \widetilde{Q}(-z)\pi(z)dz + \mathcal{O}(1). \end{aligned}$$

The last equality follows from the scaling property of $\widetilde{Q}$ and the boundedness of Q_-.

Next we show the estimate of $C_{2,1}^\epsilon$. Note that

$$C_{2,1}^\epsilon = \int \overline{K}(z)(Q^\epsilon(z))^2 dz \sim \int_{\|z\|_s > \epsilon} \overline{K}(z)(Q(z))^2 dz.$$

Indeed, since $|Q^\epsilon(z)| \lesssim \epsilon^{-1}$ and $|Q(z) - Q^\epsilon(z)| \lesssim \epsilon^\theta \|z\|_s^{-1-\theta}$ for every $\theta \in (0,1]$ (see [Hai14, Lemma 10.17]), we have

$$\left| \int_{\|z\|_s \le \epsilon} \overline{K}(z)(Q^\epsilon(z))^2 dz \right| \lesssim \epsilon^{-2} \int_{\|z\|_s \le \epsilon} \|z\|_s^{-3} dz \lesssim 1$$

and

$$\left| \int_{\|z\|_s > \epsilon} \overline{K}(z)\{(Q^\epsilon(z))^2 - (Q(z))^2\} dz \right| \lesssim \epsilon^\theta \int_{\|z\|_s > \epsilon} \|z\|_s^{-5-\theta} dz \lesssim 1.$$

Hence it suffices to consider $\int_{\|z\|_s>\epsilon} R(z)dz$, where $R = \overline{K}Q^2 \in \mathfrak{S}_5$. However, we replace R by a function S_+ defined below. Let φ_n be a smooth and nonnegative function such that $\mathrm{supp}(\varphi) \subset \{\frac{1}{2} \le \|z\|_s \le 2\}$ and $\sum_{n=-\infty}^{\infty} \varphi(2^{2n}t, 2^n x) = 1$ for all $(t,x) \neq 0$. Define

$$S_+ = \sum_{n=0}^{\infty} \widetilde{R}\varphi_n, \quad S_- = \sum_{n=-\infty}^{-1} \widetilde{R}\varphi_n,$$

where $\varphi_n(t,x) = \varphi(2^{2n}t, 2^n x)$. Note that $R + R_- = S_+ + S_-$. Since $\mathrm{supp}(S_- - R_-)$ is compact, we have

$$\int_{\|z\|_s>\epsilon} R(z)dz = \int_{\|z\|_s>\epsilon} S_+(z)dz + \mathcal{O}(1).$$

By the scaling property of $\widetilde{R}$, we have $S_+ = \sum_{n=0}^{\infty} S_n$, where

$$S_n(t,x) = 2^{5n} S_0(2^{2n}, 2^n x), \quad S_0 = \widetilde{R}\varphi.$$

Since $S_0 \in C_0^\infty$, we have

$$\begin{aligned}\int_{\|z\|_s>\epsilon} S_+(z)dz &= \sum_{n=0}^{\infty} \int_{\|z\|_s>\epsilon} S_n(z)dz \\ &= \sum_{n=0}^{\infty} \int_{\|z\|_s>\epsilon 2^n} S_0(z)dz = \sum_{n=0}^{N(\epsilon)} \int_{\|z\|_s>\epsilon 2^n} S_0(z)dz,\end{aligned}$$

where $N(\epsilon) \in \mathbf{N}$ is the largest number such that $\{\|z\|_s > \epsilon 2^{N(\epsilon)}\} \cap \mathrm{supp}(S_0) \neq \emptyset$, so there exists a constant $C > 0$ such that $N(\epsilon) \sim C \log \epsilon^{-1}$. Since

$$\begin{aligned}&\sum_{n=0}^{N(\epsilon)} \left| \int_{\|z\|_s>\epsilon 2^n} S_0(z)dz - \int S_0(z)dz \right| \\ &\lesssim \sum_{n=0}^{N(\epsilon)} \int_{\|z\|_\epsilon \le \epsilon 2^n} dz \lesssim \sum_{n=0}^{N(\epsilon)} \epsilon^5 2^{5n} \lesssim \epsilon^5 2^{5N(\epsilon)} \lesssim 1,\end{aligned}$$

we have the estimate

$$C_{2,1}^\epsilon = N(\epsilon) \int S_0(z)dz + \mathcal{O}(1) \sim C \int S_0(z)dz \log \epsilon^{-1}.$$

The estimate of $C_{2,2}^\epsilon$ is similar. □

4 CGL by the theory of paracontrolled distributions

In Sections 4 and 5, we study well-posedness of CGL (1.1) by using the paracontrolled distribution theory introduced by [GIP15]. In that paper, they stud-

ied some problems such as differential equations driven by fractional Brownian motion, a Burgers-type stochastic PDE, and a nonlinear version of the parabolic Anderson model. After that Catellier-Chouk [CC13] showed local well-posedness of the three-dimensional stochastic quantization equation (the dynamic Φ^4_3 model), which is an $\mathbf{R}$-valued version of CGL.

Our proof of the local well-posedness of CGL consists of two parts: a deterministic and a probabilistic part.

In Section 4, we deal with a deterministic version of CGL and construct a solution map from a space of driving vectors to a space of solutions. We also see that the solution map is continuous. In this section, ξ is a deterministic distribution which takes values in the Hölder-Besov space $\mathcal{C}^{-\frac{5}{2}-\kappa}$ for any $\kappa > 0$ small enough. To construct the solution map, we rely on the method introduced by Mourrat-Weber [MW16]. We state the precise assertion concerning the well-posedness in Theorem 4.26. In Theorem 4.29, we see that the solution obtained in Theorem 4.26 solves the renormalized equation (1.2) in the usual sense.

Section 5 is the probabilistic part and devoted to constructing a driving vector X associated to the space-time white noise ξ defined on $\mathbf{R} \times \mathbf{T}^3$. We follow the approach as in [GP17] and obtain the driving vector in Theorem 5.9. Here we explain how to mollify the white noise ξ. Let χ be a smooth function defined on $\mathbf{R}^3$ such that (1) $\operatorname{supp} \chi \subset B(0,1)$, where $B(x,r)$ denotes the open ball of radius $r > 0$ and canter $x \in \mathbf{R}^3$, (2) $\chi(0) = 1$. We set $\chi^\epsilon(k) = \chi(\epsilon k)$ for every $k \in \mathbf{Z}^3$. Define $\mathbf{e}_k(x) = e^{-2\pi \mathrm{i} k \cdot x}$ for every $k \in \mathbf{Z}^3$ and $x \in \mathbf{T}^3$. Here, the dot $\cdot$ denotes the usual inner product. We define ξ^ϵ by

$$\xi^\epsilon = \sum_{k \in \mathbf{Z}^3} \chi^\epsilon(k) \hat{\xi}(k) \mathbf{e}_k. \tag{4.1}$$

Here, $\{\hat{\xi}(k)\}_{k \in \mathbf{Z}^3}$ denotes the Fourier transform of ξ and it has the same law with independent copies of the complex white noise on $\mathbf{R}$. We see that $\xi^\epsilon \to \xi$ in an appropriate topology. For the smeared noise ξ^ϵ, we define a family of processes $\{X^\epsilon\}_{0<\epsilon<1}$. In this definition of X^ϵ, we will use the dyadic partition of unity $\{\rho_m\}_{m=-1}^\infty$ via the resonant $\odot$ and renormalization constants $\mathfrak{c}_1^\epsilon$, $\mathfrak{c}_{2,1}^\epsilon$ and $\mathfrak{c}_{2,2}^\epsilon$; see (5.3). We obtain the driving vector X as a limit of $\{X^\epsilon\}_{0<\epsilon<1}$. By setting $\mathfrak{c}^\epsilon = 2(\mathfrak{c}_1^\epsilon - \overline{\nu \mathfrak{c}_{2,1}^\epsilon} - 2\nu \mathfrak{c}_{2,2}^\epsilon)$, we have $|\mathfrak{c}^\epsilon| \to \infty$ as $\epsilon \to 0$.

By combining Theorems 4.26, 4.29 and 5.9, we obtain the following main theorem in Sections 4 and 5:

Theorem 4.1. *Let $0 < \kappa' < 1/18$ and $u_0 \in \mathcal{C}^{-\frac{2}{3}+\kappa'}$. Consider the renormalized equation (1.2) with $C^\epsilon = \mathfrak{c}^\epsilon$. Then, for every $0 < \epsilon < 1$, there exist a unique process u^ϵ and a random time $T_*^\epsilon \in (0,1]$ such that*

- u^ϵ *solves* (1.2) *on* $[0, T_*^\epsilon] \times \mathbf{T}^3$,
- T_*^ϵ *converges to some a.s. positive random time* T_* *in probability,*
- u^ϵ *converges to some process* u *defined on* $[0, T_*) \times \mathbf{T}^3$ *in the following sense:*

$$\lim_{\varepsilon \downarrow 0} \sup_{0 \le s \le T_*/2} \|u_s^\epsilon - u_s\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} = 0$$

in probability. Here, we set $\sup_{0 \le s \le T_*/2} \|u_s^\epsilon - u_s\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} = \infty$ *on the event* $\{T_*^\epsilon < T_*/2\}$. *Furthermore,* u *is independent of the choice of* $\{\rho_m\}_{m=-1}^\infty$ *and* χ.

Here, we will make comments on this theorem. Note that the process u^ϵ and u are obtained by substituting X^ϵ and X into the solution map, respectively. Since X^ϵ converges to X and the solution map is continuous, we see that u^ϵ converges to u. In addition, u^ϵ solves (1.2) in the usual sense, hence we see the theorem. We need to pay attention to the assertion that u is independent of the choice of ξ^ϵ. Recall that X^ϵ depends on $\{\rho_m\}_{m=-1}^\infty$. Hence u^ϵ may, too. However, we see that u^ϵ does not. In fact, we obtain an expression of the renormalization constant $\mathfrak{c}^\epsilon$ which does not depend on $\{\rho_m\}_{m=-1}^\infty$ in Proposition 5.20. Hence, (1.2) is independent of $\{\rho_m\}_{m=-1}^\infty$ and so is the solution u^ϵ. As a consequence, the limit u is independent of $\{\rho_m\}_{m=-1}^\infty$. In addition, the limit u is independent of χ because the driving vector X is independent of χ (Theorem 5.9). Hence we see the solution does not depend on $\{\rho_m\}_{m=-1}^\infty$ or χ

Sections 4 and 5 is independent of Sections 2 and 3. We do not use the symbols introduced in Sections 2 and 3.

4.1 Besov-Hölder spaces and paradifferential calculus

In this section, we introduce the Besov-Hölder spaces and paradifferential calculus. The results in this section can be found in [GIP15, BCD11] or follow from them easily.

4.1.1 Besov spaces

We introduce the Besov spaces and recall their basic properties. Let $\mathcal{D} \equiv \mathcal{D}(\mathbf{T}^3, \mathbf{C})$ be the space of all smooth $\mathbf{C}$-valued functions on $\mathbf{T}^3$ and $\mathcal{D}'$ its dual of $\mathcal{D}$. We set $\mathbf{e}_k(x) = e^{2\pi \mathrm{i} k \cdot x}$ for every $k \in \mathbf{Z}^3$ and $x \in \mathbf{T}^3$. The Fourier transform $\mathcal{F}f$ for $f \in \mathcal{D}$ is defined by $\mathcal{F}f(k) = \int_{\mathbf{T}^3} \mathbf{e}_{-k}(x) f(x)\, dx$ and its inverse $\mathcal{F}^{-1}g$ for a rapidly decreasing sequence $\{g(k)\}_{k \in \mathbf{Z}^3}$ is defined by

$\mathcal{F}^{-1}g(x) = \sum_{k\in\mathbf{Z}^3} g(k)\mathbf{e}_k(x)$. For every rapidly decreasing smooth function ϕ, we set $\phi(D)f = \mathcal{F}^{-1}\phi\mathcal{F}f = \sum_{k\in\mathbf{Z}^3}\phi(k)\hat{f}(k)\mathbf{e}_k$.

We denote by $\{\rho_m\}_{m=-1}^{\infty}$ a dyadic partition of unity, that is, it satisfies the following: (1) $\rho_m\colon \mathbf{R} \to [0,1]$ is radial and smooth, (2) $\mathrm{supp}(\rho_{-1}) \subset B(0,\frac{4}{3})$, $\mathrm{supp}(\rho_0) \subset B(0,\frac{8}{3}) \setminus B(0,\frac{3}{4})$, (3) $\rho_m(\cdot) = \rho_0(2^{-m}\cdot)$ for $m \geq 0$, (4) $\sum_{m=-1}^{\infty}\rho_m(\cdot) = 1$. Here $B(0,r) = \{x \in \mathbf{R}^3; |x| < r\}$. The Littlewood-Paley blocks $\{\triangle_m\}_{m=-1}^{\infty}$ are defined by $\triangle_m = \rho_m(D)$.

We are ready to define Besov space $\mathcal{C}^\alpha = \mathcal{C}^\alpha(\mathbf{T}^3, \mathbf{C})$ for $\alpha \in \mathbf{R}$. It is defined as the completion of $\mathcal{D}$ under the norm

$$\|f\|_{\mathcal{C}^\alpha} = \sup_{m\geq -1} 2^{m\alpha}\|\triangle_m f\|_{L^\infty}.$$

The next is frequently used results on Besov spaces:

Proposition 4.2 ([BCD11, Theorem 2.80])**.** *We have the following:*

- $\|f\|_{\mathcal{C}^\alpha} \lesssim \|f\|_{\mathcal{C}^\beta}$ *if* $\alpha < \beta$.
- *Let* $\alpha, \alpha_1, \alpha_2 \in \mathbf{R}$ *satisfy* $\alpha = (1-\theta)\alpha_1 + \theta\alpha_2$ *for some* $0 < \theta < 1$. *Then, we have*

$$\|f\|_{\mathcal{C}^\alpha} \leq \|f\|_{\mathcal{C}^{\alpha_1}}^{1-\theta}\|f\|_{\mathcal{C}^{\alpha_2}}^{\theta}.$$

4.1.2 Paraproducts and Commutator estimates

For every $f \in \mathcal{C}^\alpha$, $g \in \mathcal{C}^\beta$, we set

$$f \varolessthan g = \sum_{m_1 \geq m_2+2} \triangle_{m_1} f \triangle_{m_2} g, \quad f \varogreaterthan g = \sum_{m_1+2\leq m_2} \triangle_{m_1} f \triangle_{m_2} g,$$
$$f \odot g = \sum_{|m_1-m_2|\leq 1} \triangle_{m_1} f \triangle_{m_2} g.$$

The following are properties of paraproduct.

Proposition 4.3 (Paraproduct and resonant estimate [BCD11, Theorem 2.82 and 2.85])**.** *We have the following:*

(1) For every $\beta \in \mathbf{R}$, $\|f \varogreaterthan g\|_{\mathcal{C}^\beta} \lesssim \|f\|_{L^\infty}\|g\|_{\mathcal{C}^\beta}$.

(2) For every $\alpha < 0$ *and* $\beta \in \mathbf{R}$, $\|f \varogreaterthan g\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|f\|_{\mathcal{C}^\alpha}\|g\|_{\mathcal{C}^\beta}$.

(3) If $\alpha + \beta > 0$, *then* $\|f \odot g\|_{\mathcal{C}^{\alpha+\beta}} \lesssim \|f\|_{\mathcal{C}^\alpha}\|g\|_{\mathcal{C}^\alpha}$.

Remark 4.4. Let $u \in \mathcal{C}^\alpha$ and $g \in \mathcal{C}^\beta$ for $\alpha < 0$ and $\beta > 0$ with $\alpha + \beta > 0$. Then the product uv is well-defined as an element in $\mathcal{C}^\alpha$.

Proposition 4.5 (Commutator estimates, [GIP15, Lemma 2.4])**.** *Let* $0 < \alpha < 1$*,* $\beta, \gamma \in \mathbf{R}$ *satisfy* $\beta + \gamma < 0$ *and* $\alpha + \beta + \gamma > 0$*. Define the map* R *by*

$$R(f, g, h) = (f \varolessthan g) \odot h - f(g \odot h)$$

for $f, g, h \in C^\infty(\mathbf{T}^3, \mathbf{C})$*. Then* R *is uniquely extended to a continuous trilinear map from* $\mathcal{C}^\alpha \times \mathcal{C}^\beta \times \mathcal{C}^\gamma$ *to* $\mathcal{C}^{\alpha+\beta+\gamma}$*.*

4.1.3 Regularity of $\mathcal{C}^\alpha$-valued functions

Here we consider $\mathcal{C}^\alpha$-valued functions. We introduce several classes of them as follows:

- $C_T\mathcal{C}^\alpha$ is the space of all continuous functions from $[0, T]$ to $\mathcal{C}^\alpha$ which is equipped with the supremum norm

$$\|u\|_{C_T\mathcal{C}^\alpha} = \sup_{0 \le t \le T} \|u_t\|_{\mathcal{C}^\alpha},$$

- $C_T^\delta\mathcal{C}^\alpha$ is the space of all δ-Hölder continuous functions from $[0, T]$ to $\mathcal{C}^\alpha$ which is equipped with the seminorm

$$\|u\|_{C_T^\delta\mathcal{C}^\alpha} = \sup_{0 \le s < t \le T} \frac{\|u_t - u_s\|_{\mathcal{C}^\alpha}}{|t - s|^\delta},$$

- $\mathcal{E}_T^\eta\mathcal{C}^\alpha = \{u \in C((0, T], \mathcal{C}^\alpha); \|u\|_{\mathcal{E}_T^\eta\mathcal{C}^\alpha} < \infty\}$, where

$$\|u\|_{\mathcal{E}_T^\eta\mathcal{C}^\alpha} = \sup_{0 < t \le T} t^\eta \|u_t\|_{\mathcal{C}^\alpha},$$

- $\mathcal{E}_T^{\eta,\delta}\mathcal{C}^\alpha = \{u \in C((0, T], \mathcal{C}^\alpha); \|u\|_{\mathcal{E}_T^{\eta,\delta}\mathcal{C}^\alpha} < \infty\}$, where

$$\|u\|_{\mathcal{E}_T^{\eta,\delta}\mathcal{C}^\alpha} = \sup_{0 < s < t \le T} s^\eta \frac{\|u_t - u_s\|_{\mathcal{C}^\alpha}}{|t - s|^\delta},$$

- $\mathcal{L}_T^{\alpha,\delta} = C_T\mathcal{C}^\alpha \cap C_T^\delta\mathcal{C}^{\alpha-2\delta}$,
- $\mathcal{L}_T^{\eta,\alpha,\delta} = \mathcal{E}_T^\eta\mathcal{C}^\alpha \cap C_T\mathcal{C}^{\alpha-2\eta} \cap \mathcal{E}_T^{\eta,\delta}\mathcal{C}^{\alpha-2\delta}$.

Remark 4.6. We introduced the norms on the spaces $\mathcal{E}_T^\eta\mathcal{C}^\alpha$ and $\mathcal{E}_T^{\eta,\delta}\mathcal{C}^\alpha$ in order to control explosion at $t = 0$. The definition of $\mathcal{L}_T^{\alpha,\delta}$ is natural from the time-space scaling of CGL.

For $\mu > 0$, we set $\mathcal{L}^1 = \partial_t - \{(\mathrm{i} + \mu)\triangle - 1\}$, $P_t^1 = e^{t\{(\mathrm{i}+\mu)\triangle - 1\}}$. We present results on smoothing effects of semigroup $\{P_t^1\}_{t \ge 0}$.

Proposition 4.7 (Effects of heat semigroup)**.** *Let $\alpha \in \mathbf{R}$.*

(1) For every $\delta > 0$, $\|P_t^1 u\|_{\mathcal{C}^{\alpha+2\delta}} \lesssim t^{-\delta}\|u\|_{\mathcal{C}^\alpha}$ uniformly in $t > 0$.

(2) For every $\delta \in [0,1]$, $\|(P_t^1 - 1)u\|_{\mathcal{C}^{\alpha-2\delta}} \lesssim t^{\delta}\|u\|_{\mathcal{C}^\alpha}$ uniformly in $t > 0$.

Proposition 4.8 (Schauder estimates)**.** *Let $T > 0$. We see the following:*

(1) Let $u \in \mathcal{C}^\alpha$. For every $\beta \geq \alpha$ and $\delta \in [0,1]$, we have

$$\|(t \mapsto P_t^1 u)_{t\geq 0}\|_{\mathcal{L}_T^{\frac{\beta-\alpha}{2},\beta,\delta}} \lesssim \|u\|_{\mathcal{C}^\alpha}.$$

(2) Let $\alpha \neq \beta$. Let $u \in \mathcal{E}_T^\eta \mathcal{C}^\alpha$ for $\eta \in [0,1)$ and set

$$U_t = \int_0^t P_{t-s}^1 u_s \, ds.$$

Then for every $\gamma \in [\alpha, \alpha - 2\eta + 2)$, $\beta \in [\gamma, \alpha + 2)$ and $\delta \in (0, \frac{\beta-\alpha}{2}]$, we have

$$\|U\|_{\mathcal{L}_T^{\frac{\beta-\gamma}{2},\beta,\delta}} \lesssim T^{\frac{\alpha-2\eta+2-\gamma}{2}} \|u\|_{\mathcal{E}_T^\eta \mathcal{C}^\alpha}.$$

Proposition 4.9 (Commutation between paraproduct and heat semigroup)**.** *Let $\alpha < 1$, $\beta \in \mathbf{R}$, $\delta \geq 0$. Define*

$$[P_t^1, u\varotimes]v = P_t^1(u \varotimes v) - u \varotimes P_t^1 v.$$

Then we have

$$\|[P_t^1, u\varotimes]v\|_{\mathcal{C}^{\alpha+\beta+2\delta}} \lesssim t^{-\delta}\|u\|_{\mathcal{C}^\alpha}\|v\|_{\mathcal{C}^\beta}$$

uniformly over $t > 0$.

We can show the above results in a similar way as [Hos16b, Corollary 2.6, Proposition 2.8] and [MW16, Propsosition A.15], because $\mu > 0$.

4.2 Definitions of driving vectors and solutions

First of all, we give the definition of a driving vector. We set

$$I(u_0, v)_t = P_t^1 u_0 + \int_0^t P_{t-s}^1 v_s \, ds, \tag{4.2}$$

$$I(v)_t = \int_{-\infty}^t P_{t-s}^1 v_s \, ds, \tag{4.3}$$

whenever they are well-defined. Note that if we can choose $u_0 = \int_{-\infty}^0 P_{-s}^1 v_s \, ds$, then $I(u_0, v) = I(v)$.

Let $0 < \kappa < \kappa' < 1/18$ and $T > 0$. The following is the definition of a driving vector.

Definition 4.10. We call a vector of space-time distributions

$$X = (X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet}, X^{\bullet})$$
$$\in C_T\mathcal{C}^{-\frac{1}{2}-\kappa}\times(C_T\mathcal{C}^{-1-\kappa})^2\times(C_T\mathcal{C}^{1-\kappa})^2\times\mathcal{L}_T^{\frac{1}{2}-\kappa,\frac{1}{4}-\frac{1}{2}\kappa}\times(C_T\mathcal{C}^{-\kappa})^6\times(C_T\mathcal{C}^{-\frac{1}{2}-\kappa})^2$$

which satisfies $I(X_0^{\bullet}, X^{\bullet}) = X^{\bullet}$ and $I(X_0^{\bullet}, X^{\bullet}) = X^{\bullet}$ a *driving vector* of CGL. We denote by $\mathcal{X}_T^\kappa$ the set of all driving vectors. We define the norm $\|\cdot\|_{\mathcal{X}_T^\kappa}$ by the sum of the norm of each component.

Note that we assume that the component $X^{\bullet}$ has Hölder continuity and it belongs to $\mathcal{L}_T^{\frac{1}{2}-\kappa,\frac{1}{4}-\frac{1}{2}\kappa}$. We easily see that the space $\mathcal{X}_T^\kappa$ is a closed set of the product Banach spaces. Next we define the space of solutions and give the notion of a solution.

Definition 4.11. We set

$$\mathcal{D}_T^{\kappa,\kappa'} = \mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}\times\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}.$$

Next, we fix $X\in\mathcal{X}_T^\kappa$ and set $Z = X^{\bullet}$ and $W = X^{\bullet}$. Define F and G on $\mathcal{D}_T^{\kappa,\kappa'}$ by

$$F(v,w) = -\nu\{2(-\nu X^{\bullet} + v + w)\circledcirc X^{\bullet} + (-\overline{\nu}\overline{X^{\bullet}} + \overline{v} + \overline{w})\circledcirc X^{\bullet}\}, \tag{4.4}$$
$$G(v,w) = G_1(v,w) + \cdots + G_8(v,w). \tag{4.5}$$

Here $G_1(v,w),\dots,G_8(v,w)$ will be defined shortly.

Since $Z_t\in\mathcal{C}^{-\frac{1}{2}-\kappa}$ and $W_t\in\mathcal{C}^{\frac{1}{2}-\kappa}$, the product W_tZ_t and $W_t\overline{Z}_t$ are not defined a priori. We define them by

$$WZ = W(\circledcirc+\circledcirc)Z + X^{\bullet},\qquad W\overline{Z} = W(\circledcirc+\circledcirc)Z + X^{\bullet}.$$

The products $W^2\overline{Z}$ and $W\overline{W}Z$ are also defined by

$$W^2\overline{Z} = 2WX^{\bullet} + R(W,\overline{Z},W) + (W\circledcirc\overline{Z})(\circledcirc+\circledcirc)W + W(W\circledcirc\overline{Z}),$$
$$W\overline{W}Z = \overline{W}X^{\bullet} + W\overline{X^{\bullet}} + R(\overline{W},Z,W) + (\overline{W}\circledcirc Z)(\circledcirc+\circledcirc)W + W(\overline{W}\circledcirc Z).$$

It follows from Proposition 4.3 that $(WZ)_t, (W\overline{Z})_t, (W^2\overline{Z})_t, (W\overline{W}Z)_t\in\mathcal{C}^{-\frac{1}{2}-\kappa}$ hold.

In order to define $G_6(v,w)$, we use $\mathrm{com}(v,w)$ defined as follows. For every $v_0 \in \mathcal{C}^{-\frac{2}{3}+\kappa'}$ and $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$, we set

$$\hat{v}_t = P_t^1 v_0 + \int_0^t P_{t-s}^1 [F(v,w)(s)]\, ds.$$

Define

$$\mathrm{com}(v,w) = \hat{v} + \nu\{2(-\nu W + v + w) \circledcirc X^{\Upsilon^{\cdot}} + (-\nu\overline{W} + \overline{v} + \overline{w}) \circledcirc X^{\Upsilon} \}. \tag{4.6}$$

From Lemma 4.22, we see that $\mathrm{com}(v,w) \odot X^{\Psi^{\cdot}}$ and $\overline{\mathrm{com}(v,w)} \odot X^{\Psi}$ are well-defined. Roughly speaking, $\mathrm{com}(v,w)_t$ is something like

$$[[I(\cdot), -2\nu(-\nu W + v + w) \circledcirc] X^{\Psi^{\cdot}}]_t + [[I(\cdot), -\nu(-\nu\overline{W} + \overline{v} + \overline{w}) \circledcirc] X^{\Psi}]_t.$$

Here, $[[I(\cdot), u \circledcirc] v]_t = I(u \circledcirc v)_t - u_t \circledcirc I(v)_t$.

We are in a position to define $G_1, \ldots, G_8$. We write $u_2 = v + w$ and set

$$\begin{aligned}
G_1(v,w) &= -\nu u_2^2 \overline{u_2}, \\
G_2(v,w) &= -\nu\{u_2^2(\overline{Z} - \overline{\nu}\overline{W}) + 2u_2\overline{u_2}(Z - \nu W)\}, \\
G_3(v,w) &= -\nu\big\{u_2(2\nu\overline{\nu}W\overline{W} - 2\nu W\overline{Z} - 2\overline{\nu}\overline{W}Z - 4\nu X^{\Psi^{\cdot}} - \overline{\nu}\overline{X^{\Psi^{\cdot}}}) \\
&\qquad + \overline{u_2}(\nu^2 W^2 - 2\nu W Z - 2\overline{\nu}\overline{X^{\Psi^{\cdot}}} - 2\nu X^{\Psi^{\cdot}})\big\} \\
&\quad + (\nu+1)u_2, \\
G_4(v,w) &= -\nu\big\{ - \nu^2\overline{\nu}W^2\overline{W} + \nu^2 W^2\overline{Z} + 2\nu\overline{\nu}W\overline{W}Z \\
&\qquad + 4\nu^2 W X^{\Psi^{\cdot}} + 4\nu^2 R(W, X^{\Upsilon^{\cdot}}, X^{\Psi^{\cdot}}) \\
&\qquad + 2\overline{\nu}^2\overline{W}\overline{X^{\Psi^{\cdot}}} + 2\overline{\nu}^2 R(\overline{W}, \overline{X^{\Upsilon^{\cdot}}}, X^{\Psi}) \\
&\qquad + 2\nu\overline{\nu}\overline{W}X^{\Psi^{\cdot}} + 2\nu\overline{\nu}R(\overline{W}, X^{\Upsilon}, X^{\Psi^{\cdot}}) \\
&\qquad + \nu\overline{\nu}W\overline{X^{\Psi^{\cdot}}} + \nu\overline{\nu}R(W, \overline{X^{\Upsilon}}, X^{\Psi}) \\
&\qquad - 2\nu X^{\Psi^{\cdot}} - 2\nu W \circledcirc X^{\Psi^{\cdot}} \\
&\qquad - \overline{\nu}\overline{X^{\Psi^{\cdot}}} - \overline{\nu}\overline{W} \circledcirc X^{\Psi}\big\} \\
&\quad + (\nu+1)(Z - \nu W), \\
G_5(v,w) &= -\nu\big\{ - 4\nu R(u_2, X^{\Upsilon^{\cdot}}, X^{\Psi^{\cdot}}) - 2\nu R(\overline{u_2}, X^{\Upsilon}, X^{\Psi^{\cdot}}) \\
&\qquad - 2\overline{\nu}R(\overline{u_2}, \overline{X^{\Upsilon^{\cdot}}}, X^{\Psi}) - \overline{\nu}R(u_2, \overline{X^{\Upsilon}}, X^{\Psi})\big\}, \\
G_6(v,w) &= -\nu\{2\,\mathrm{com}(v,w) \odot X^{\Psi^{\cdot}} + \overline{\mathrm{com}(v,w)} \odot X^{\Psi}\}, \\
G_7(v,w) &= -\nu\{2w \odot X^{\Psi^{\cdot}} + \overline{w} \odot X^{\Psi}\},
\end{aligned}$$

$$G_8(v,w) = -\nu\{2u_2 \varcirc X^{\Vdots} + \overline{u_2} \varcirc X^{\Vdots}\}.$$

The map $\mathcal{M} = (\mathcal{M}^1, \mathcal{M}^2)$ is defined on $\mathcal{D}_T^{\kappa,\kappa'}$ by

$$[\mathcal{M}^1(v,w)](t) = P_t^1 v_0 + \int_0^t P_{t-s}^1 F(v,w)(s)\,ds, \tag{4.7}$$

$$[\mathcal{M}^2(v,w)](t) = P_t^1 w_0 + \int_0^t P_{t-s}^1 G(v,w)(s)\,ds \tag{4.8}$$

for every $(v_0, w_0) \in \mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa}$. We will use Proposition 4.8 to check that the map is well-defined map from $\mathcal{D}_T^{\kappa,\kappa'}$ to itself and has good property.

Definition 4.12. For every $(v_0, w_0) \in \mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa}$ and $X \in \mathcal{X}_T^\kappa$, we consider the system

$$\begin{cases} v_t = [\mathcal{M}^1(v,w)](t), \\ w_t = [\mathcal{M}^2(v,w)](t). \end{cases} \tag{4.9}$$

If there exists $(v,w) \in \mathcal{D}_{T_*}^{\kappa,\kappa'}$ for some $T_* > 0$, then we call (v,w) the solution to (1.1) on $[0, T_*]$.

We interpret (4.9) as a fixed point problem $\mathcal{M} : (v,w) \mapsto (\mathcal{M}^1(v,w), \mathcal{M}^2(v,w))$. We show that the map $\mathcal{M}$ is well-defined and a contraction in Section 4.3. Section 4.4 is devoted to the construction and the uniqueness of the solution. We show that the solution to (1.1) satisfies a renormalized equation in Section 4.5. In that section, we see the validity of the notion of the solution to CGL.

Before starting our discussion, we will remark on the function spaces we have just introduced.

Remark 4.13. We make several comments on $\mathcal{L}_T^{\eta,\alpha}$ and $\mathcal{L}_T^{\eta,\alpha,\delta}$.

- The inclusion $\mathcal{L}_T^{\alpha,\delta} \subset \mathcal{L}_T^{\alpha,\delta'}$ holds for every $0 < \delta' < \delta$. To prove this assertion, we use Proposition 4.2. Set $\theta = \delta'/\delta$. Then $\alpha - 2\delta' = (\alpha - 2\delta)\theta + \alpha(1-\theta)$. For every $W \in \mathcal{L}_T^{\alpha,\delta}$, we see

$$\begin{aligned} \|W_t - W_s\|_{\mathcal{C}^{\alpha-2\delta'}} &\le \|W_t - W_s\|_{\mathcal{C}^{\alpha-2\delta}}^{\theta} \|W_t - W_s\|_{\mathcal{C}^{\alpha}}^{1-\theta} \\ &\lesssim \{(t-s)^{\delta} \|W\|_{C_T^\delta \mathcal{C}^{\alpha-2\delta}}\}^{\theta} \|W\|_{C_T\mathcal{C}^{\alpha}}^{1-\theta} \\ &\lesssim (t-s)^{\delta'} \|W\|_{\mathcal{L}_T^{\alpha,\delta}}. \end{aligned}$$

- The inclusion $\mathcal{L}_T^{\eta,\alpha,\delta} \subset \mathcal{L}_T^{\eta,\alpha,\delta'}$ holds for $0 < \delta' < \delta$. Indeed, for every $v \in \mathcal{L}_T^{\eta,\alpha,\delta}$ and $0 < s < t \le T$, we have

$$\|v_t - v_s\|_{\mathcal{C}^{\alpha-2\delta}} \le s^{-\eta} |t-s|^{\delta} \|v\|_{\mathcal{E}_T^{\eta,\delta}\mathcal{C}^{\alpha}},$$

$$\|v_t - v_s\|_{\mathcal{C}^\alpha} \le \|v_t\|_{\mathcal{C}^\alpha} + \|v_s\|_{\mathcal{C}^\alpha} \le t^{-\eta}\|v\|_{\mathcal{E}_T^\eta \mathcal{C}^\alpha} + s^{-\eta}\|v\|_{\mathcal{E}_T^\eta \mathcal{C}^\alpha}.$$

Hence, for $0 < \theta < 1$, we see

$$\begin{aligned}\|v_t - v_s\|_{\mathcal{C}^{\alpha-2\delta'}} &\le \|v_t - v_s\|_{\mathcal{C}^{\alpha-2\delta}}^{\theta} \|v_t - v_s\|_{\mathcal{C}^\alpha}^{1-\theta} \\ &\lesssim \{s^{-\eta}|t-s|^\delta \|v\|_{\mathcal{E}_T^{\eta,\delta}\mathcal{C}^\alpha}\}^\theta \{s^{-\eta}\|v\|_{\mathcal{E}_T^\eta \mathcal{C}^\alpha}\}^{1-\theta} \\ &\lesssim s^{-\eta}|t-s|^{\delta'} \|v\|_{\mathcal{L}_T^{\eta,\alpha,\delta}},\end{aligned}$$

which implies $v \in \mathcal{L}_T^{\eta,\alpha,\delta'}$.

- For every $v \in \mathcal{L}_T^{\eta,\alpha,\delta}$ and $\alpha - 2\eta \le \gamma \le \alpha$, we have $v_t \in \mathcal{C}^\gamma$ and

$$\|v_t\|_{\mathcal{C}^\gamma} \le t^{-\frac{1}{2}(\gamma-(\alpha-2\eta))} \|v\|_{\mathcal{L}_T^{\eta,\alpha,\delta}}$$

for any $0 < t \le T$. Since $v_t \in \mathcal{C}^{\gamma-2\eta} \cap \mathcal{C}^\gamma$, we take θ such that $\gamma = (\alpha - 2\eta)(1-\theta) + \alpha\theta$ and use Proposition 4.2 to obtain

$$\|v_t\|_{\mathcal{C}^\gamma} \le \|v_t\|_{\mathcal{C}^{\alpha-2\eta}}^{1-\theta} \|v_t\|_{\mathcal{C}^\alpha}^{\theta} \le \|v\|_{C_T\mathcal{C}^{\alpha-2\eta}}^{1-\theta} \{t^{-\eta}\|v\|_{\mathcal{E}_T^\eta \mathcal{C}^\alpha}\}^\theta \le t^{-\eta\theta}\|v\|_{\mathcal{L}_T^{\eta,\alpha,\delta}}.$$

Combining this with $\eta\theta = -\frac{1}{2}(\gamma - (\alpha - 2\eta))$, we see the assertion.

Remark 4.14. We make several comments on $\mathcal{L}_T^{\frac{1}{2}-\kappa,\frac{1}{4}-\frac{1}{2}\kappa}$ and $\mathcal{D}_T^{\kappa,\kappa'}$. Recall that

$$\begin{aligned}\mathcal{L}_T^{\frac{1}{2}-\kappa,\frac{1}{4}-\frac{1}{2}\kappa} &= C_T\mathcal{C}^{\frac{1}{2}-\kappa} \cap C_T^{\frac{1}{4}-\frac{1}{2}\kappa}\mathcal{C}^0, \\ \mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'} &= \mathcal{E}_T^{\frac{5}{6}-\kappa'}\mathcal{C}^{1-\kappa'} \cap C_T\mathcal{C}^{-\frac{2}{3}+\kappa'} \cap \mathcal{E}_T^{\frac{5}{6}-\kappa',1-\frac{1}{2}\kappa'}\mathcal{C}^{-1}, \\ \mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'} &= \mathcal{E}_T^{1-\kappa'+\kappa}\mathcal{C}^{\frac{3}{2}-2\kappa'} \cap C_T\mathcal{C}^{-\frac{1}{2}-2\kappa} \cap \mathcal{E}_T^{1-\kappa'+\kappa,1-\kappa'}\mathcal{C}^{-\frac{1}{2}}.\end{aligned}$$

- For every $0 < \kappa < \kappa'$, we have $\mathcal{L}_T^{\frac{1}{2}-\kappa,\frac{1}{4}-\frac{1}{2}\kappa} \subset \mathcal{L}_T^{\frac{1}{2}-\kappa,\frac{1}{4}-\frac{1}{2}\kappa'} = C_T\mathcal{C}^{\frac{1}{2}-\kappa} \cap C_T^{\frac{1}{4}-\frac{1}{2}\kappa}\mathcal{C}^{\kappa'-\kappa}$. This inclusion implies

$$\|X_t^{\Psi} - X_s^{\Psi}\|_{\mathcal{C}^{\kappa'-\kappa}} \lesssim (t-s)^{\frac{1}{4}-\frac{1}{2}\kappa} \|X^{\Psi}\|_{C_T^{\frac{1}{4}-\frac{1}{2}\kappa}\mathcal{C}^{\kappa'-\kappa}}.$$

- Note that

$$\begin{aligned}\mathcal{D}_T^{\kappa,\kappa'} &= \mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'} \times \mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'} \\ &\subset \mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',\frac{1}{2}-\kappa'} \times \mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',\frac{1}{2}-\kappa'}\end{aligned}$$

holds and, for every $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have

$$\begin{aligned}\|v_t - v_s\|_{L^\infty} \lesssim \|v_t - v_s\|_{\mathcal{C}^{\kappa'}} &\lesssim s^{-\left(\frac{5}{6}-\kappa'\right)}(t-s)^{\frac{1}{2}-\kappa'} \|v\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}}, \\ \|w_t - w_s\|_{L^\infty} \lesssim \|w_t - w_s\|_{\mathcal{C}^{\kappa'}} &\lesssim s^{-(1-\kappa'+\kappa)}(t-s)^{\frac{1}{2}-\kappa'} \|w\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}}.\end{aligned}$$

- For every $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have

$$\|v_t\|_{\mathcal{C}^\alpha} \le t^{-\frac{1}{2}\left(\alpha+\frac{2}{3}-\kappa'\right)} \|v\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}},$$
$$\|w_t\|_{\mathcal{C}^\beta} \le t^{-\frac{1}{2}\left(\beta+\frac{1}{2}+2\kappa\right)} \|w\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}},$$

  where α and β satisfy $-\frac{2}{3}+\kappa' \le \alpha \le 1-\kappa'$ and $-\frac{1}{2}-2\kappa \le \beta \le \frac{3}{2}-2\kappa'$. In particular, for $\alpha = \beta = \kappa' - \kappa$, we have

$$\|v_t\|_{L^\infty} \le \|v_t\|_{\mathcal{C}^{\kappa'-\kappa}} \le t^{-\frac{2-3\kappa}{6}} \|v\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}},$$
$$\|w_t\|_{L^\infty} \le \|w_t\|_{\mathcal{C}^{\kappa'-\kappa}} \le t^{-\frac{1}{2}\left(\frac{1}{2}+\kappa'+\kappa\right)} \|w\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}},$$
$$\|v_t + w_t\|_{L^\infty} \le \|v_t\|_{L^\infty} + \|w_t\|_{L^\infty} \lesssim t^{-\frac{2-3\kappa}{6}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}.$$

  In the last estimate, we used $0 < \kappa < \kappa' < 1/18$ and $0 < t \le T$. We also see

$$\|v_t + w_t\|_{\mathcal{C}^\gamma} \lesssim t^{-\frac{1}{2}\left(\gamma+\frac{2}{3}-\kappa'\right)} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}$$

  if $-\frac{1}{2}-2\kappa \le \gamma \le 1-\kappa'$.

4.3 Properties of the integration map

Let $0 < \kappa < \kappa' < 1/18$ and $0 < T \le 1$. We fix $X, X^{(1)}, X^{(2)} \in \mathcal{X}_1^\kappa$ and set $Z = X^{\wr}$, $W = X^{\Psi}$, $Z^{(i)} = X^{\wr,(i)}$ and $W^{(i)} = X^{(i),\Psi}$ for $i = 1, 2$. We sometimes use the symbol F_X, G_X and $\mathcal{M}_{(v_0,w_0),X} = (\mathcal{M}^1_{(v_0,w_0),X}, \mathcal{M}^2_{(v_0,w_0),X})$ to indicate the dependence on the driving vector X and the initial data (v_0, w_0).

4.3.1 Properties of $\mathcal{M}^1$

Let us start our discussion with F.

Lemma 4.15. *For any $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$ and $0 < t \le T$, we have $F(v,w)(t) \in \mathcal{C}^{-1-\kappa}$ and*

$$\|F(v,w)(t)\|_{\mathcal{C}^{-1-\kappa}} \le C(\|X\|_{\mathcal{X}_1^\kappa} + t^{-\frac{2-3\kappa}{6}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}) \|X\|_{\mathcal{X}_1^\kappa},$$

where C is a positive constant depending only on κ, κ', μ and ν.

Proof. Set $\Phi = -\nu W + v + w$. From $W_t \in \mathcal{C}^{\frac{1}{2}-\kappa}$, $v_t \in \mathcal{C}^{1-\kappa'}$ and $w_t \in \mathcal{C}^{\frac{3}{2}-2\kappa'}$, we see that $\Phi_t \in \mathcal{C}^{\frac{1}{2}-\kappa}$ and $\|\Phi_t\|_{L^\infty} \le \|W_t\|_{L^\infty} + \|v_t\|_{L^\infty} + \|w_t\|_{L^\infty}$ hold for

every $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$. Note that $\|W_t\|_{L^\infty} \lesssim \|W_t\|_{\mathcal{C}^{\frac{1}{2}-\kappa}} \le \|X\|_{\mathcal{X}_T^\kappa}$ holds from Proposition 4.2. From Remark 4.14, we see

$$\|v_t\|_{L^\infty} \lesssim t^{-\frac{2-3\kappa}{6}} \|v\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}} \le t^{-\frac{2-3\kappa}{6}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}},$$
$$\|w_t\|_{L^\infty} \lesssim t^{-\frac{1}{2}\left(\frac{1}{2}+\kappa'+\kappa\right)} \|w\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}} \le t^{-\frac{2-3\kappa}{6}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}.$$

Combining this with $X_t^{\mathbb{V}} \in \mathcal{C}^{-1-\kappa}$ and using Proposition 4.3, we see $\Phi_t \circledcirc X_t^{\mathbb{V}} \in \mathcal{C}^{-1-\kappa}$ and

$$\begin{aligned}\|\Phi_t \circledcirc X_t^{\mathbb{V}}\|_{\mathcal{C}^{-1-\kappa}} &\lesssim \|\Phi_t\|_{L^\infty} \|X_t^{\mathbb{V}}\|_{\mathcal{C}^{-1-\kappa}} \\ &\le (\|X\|_{\mathcal{X}_1^\kappa} + t^{-\frac{2-3\kappa}{6}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}) \|X\|_{\mathcal{X}_1^\kappa}.\end{aligned}$$

The term $\overline{\Phi_t} \circledcirc X_t^{\mathbb{V}}$ also has a similar bound. From the defintion of $F(v,w)$, we see the assertion. □

Proposition 4.16. *The map $\mathcal{M}^1 : \mathcal{D}_T^{\kappa,\kappa'} \to \mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}$ is well-defined and, for any $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have*

$$\|\mathcal{M}^1(v,w)\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}} \le C_1(1 + \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}) + C_2 T^{\frac{1-\kappa'}{2}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}.$$

Here, C_1 and C_2 are positive constants depending only on κ, κ', μ, ν and $\|X\|_{\mathcal{X}_1^\kappa}$. In particular, they are given by at most second-order polynomials in $\|X\|_{\mathcal{X}_1^\kappa}$.

Proof. Applying the first assertion of Proposition 4.8 with $\alpha = -\frac{2}{3} + \kappa'$, $\beta = 1-\kappa'$ and $\delta = 1 - \frac{1}{2}\kappa'$ to $v_0 \in \mathcal{C}^{-\frac{2}{3}+\kappa'}$, we see

$$\|(t \mapsto P_t^1 v_0)_{t\ge 0}\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}} \lesssim \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}.$$

From Lemma 4.15, we see $F(v,w) \in \mathcal{E}_T^{\frac{2-3\kappa}{6}} \mathcal{C}^{-1-\kappa}$ and its norm has an upper bound $C_1(1 + \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}})$. Here, C_1 is positive and is given by a polynomial in $\|X\|_{\mathcal{X}_1^\kappa}$. Applying the second assertion in Proposition 4.8 with $\alpha = -1-\kappa$, $\beta = 1-\kappa'$, $\gamma = -\frac{2}{3} + \kappa'$ $\delta = 1 - \frac{1}{2}\kappa'$ and $\eta = \frac{2-3\kappa}{6}$ to $F(v,w) \in \mathcal{E}_T^{\frac{2-3\kappa}{6}} \mathcal{C}^{-1-\kappa}$, we see

$$\begin{aligned}&\left\| (t \mapsto \int_0^t P_{t-s}^1 F(v,w)(s)\, ds)_{t\ge 0} \right\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}} \\ &\qquad \lesssim T^{\frac{1-\kappa'}{2}} \|F(v,w)\|_{\mathcal{E}_T^{\frac{2-3\kappa}{6}} \mathcal{C}^{-1-\kappa}} \le T^{\frac{1-\kappa'}{2}} C_2(1 + \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}).\end{aligned}$$

The proof is completed. □

Next we show that $\mathcal{M}^1$ is Lipschitz continuous.

Lemma 4.17. *For any $(v^{(1)}, w^{(1)}), (v^{(2)}, w^{(2)}) \in \mathcal{D}_T^{\kappa,\kappa'}$ and $0 < t \le T$, we have*

$$\begin{aligned}&\|F_{X^{(1)}}(v^{(1)}, w^{(1)})(t) - F_{X^{(2)}}(v^{(2)}, w^{(2)})(t)\|_{\mathcal{C}^{-1-\kappa}} \\ &\quad \le C(1 + t^{-\frac{2-3\kappa}{6}}) \left\{ \|X^{(1)} - X^{(2)}\|_{\mathcal{X}_1^\kappa} + \|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \right\}.\end{aligned}$$

Here, C is a positive constant depending only on κ, κ', μ, ν, $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)}, w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$. In particular, it is given by a first-order polynomial in $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)}, w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$.

Proof. Set $\Phi^{(i)} = -\nu W^{(i)} + v^{(i)} + w^{(i)}$ for $i = 1, 2$. Then

$$\begin{aligned}&\|\Phi_t^{(1)} \circledcirc X_t^{(1),\mathbb{V}} - \Phi_t^{(2)} \circledcirc X_t^{(2),\mathbb{V}}\|_{\mathcal{C}^{-1-\kappa}} \\ &\qquad = \frac{1}{2}\|(\Phi_t^{(1)} + \Phi_t^{(2)}) \circledcirc (X_t^{(1),\mathbb{V}} - X_t^{(2),\mathbb{V}}) \\ &\qquad\qquad + (\Phi_t^{(1)} - \Phi_t^{(2)}) \circledcirc (X_t^{(1),\mathbb{V}} + X_t^{(2),\mathbb{V}})\|_{\mathcal{C}^{-1-\kappa}} \\ &\qquad \lesssim \|\Phi_t^{(1)} + \Phi_t^{(2)}\|_{L^\infty} \|X_t^{(1),\mathbb{V}} - X_t^{(2),\mathbb{V}}\|_{\mathcal{C}^{-1-\kappa}} \\ &\qquad\qquad + \|\Phi_t^{(1)} - \Phi_t^{(2)}\|_{L^\infty} \|X_t^{(1),\mathbb{V}} + X_t^{(2),\mathbb{V}}\|_{\mathcal{C}^{-1-\kappa}}.\end{aligned}$$

The term $\|\Phi_t^{(1)} + \Phi_t^{(2)}\|_{L^\infty}$ is dominated as follows:

$$\begin{aligned}\|\Phi_t^{(1)} + \Phi_t^{(2)}\|_{L^\infty} &\lesssim \|\Phi_t^{(1)}\|_{L^\infty} + \|\Phi_t^{(2)}\|_{L^\infty} \\ &\lesssim \sum_{i=1,2} (\|X^{(i)}\|_{\mathcal{X}_T^\kappa} + t^{-\frac{2-3\kappa}{6}} \|(v, w)^{(i)}\|_{\mathcal{D}_T^{\kappa,\kappa'}}) \|X^{(i)}\|_{\mathcal{X}_T^\kappa} \\ &= C_1 + t^{-\frac{2-3\kappa}{6}} C_2,\end{aligned}$$

where C_1 and C_2 are positive constants given by

$$\begin{aligned}C_1 &= \|X^{(1)}\|_{\mathcal{X}_1^\kappa}^2 + \|X^{(2)}\|_{\mathcal{X}_1^\kappa}^2, \\ C_2 &= \|(v^{(1)}, w^{(1)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \|X^{(1)}\|_{\mathcal{X}_1^\kappa} + \|(v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \|X^{(2)}\|_{\mathcal{X}_1^\kappa}.\end{aligned}$$

The term $\|\Phi_t^{(1)} - \Phi_t^{(2)}\|_{L^\infty}$ is dominated as follows:

$$\|\Phi_t^{(1)} - \Phi_t^{(2)}\|_{L^\infty} \lesssim \|X^{(1)} - X^{(2)}\|_{\mathcal{X}_T^\kappa} + t^{-\frac{2-3\kappa}{6}} \|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}.$$

Setting $C_3 = \|X^{(1)}\|_{\mathcal{X}_1^\kappa} + \|X^{(2)}\|_{\mathcal{X}_1^\kappa}$, we see

$$\|\Phi_t^{(1)} \circledcirc X_t^{(1),\mathbb{V}} - \Phi_t^{(2)} \circledcirc X_t^{(2),\mathbb{V}}\|_{\mathcal{C}^{-1-\kappa}}$$

$$
\begin{aligned}
&\lesssim (C_1 + t^{-\frac{2-3\kappa}{6}} C_2) \|X^{(1)} - X^{(2)}\|_{\mathcal{X}_T^\kappa} \\
&\quad + \left\{ \|X^{(1)} - X^{(2)}\|_{\mathcal{X}_T^\kappa} + t^{-\frac{2-3\kappa}{6}} \|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \right\} C_3 \\
&\lesssim (C_1 + C_2 + C_3)(1 + t^{-\frac{2-3\kappa}{6}}) \\
&\quad \times \left\{ \|X^{(1)} - X^{(2)}\|_{\mathcal{X}_T^\kappa} + \|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \right\},
\end{aligned}
$$

which implies the conclusion. □

Proposition 4.18. *For any $(v^{(1)}, w^{(1)}), (v^{(2)}, w^{(2)}) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have*

$$
\begin{aligned}
&\|\mathcal{M}^1_{(v_0^{(1)}, w_0^{(1)}), X^{(1)}}(v^{(1)}, w^{(1)}) - \mathcal{M}^1_{(v_0^{(2)}, w_0^{(2)}), X^{(2)}}(v^{(2)}, w^{(2)})\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa', 1-\kappa', 1-\frac{1}{2}\kappa'}} \\
&\qquad \le C_3 \|v_0^{(1)} - v_0^{(2)}\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} \\
&\qquad + C_4 T^{\frac{1-\kappa'}{2}} \left(\|X^{(1)} - X^{(2)}\|_{\mathcal{X}_1^\kappa} + \|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \right)
\end{aligned}
$$

Here, C_3 and C_4 are positive constants depending only on κ, κ', μ, ν, $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)}, w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$. In particular, they are given by at most first-order polynomials in $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)}, w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$.

Proof. The assertion follows from Lemma 4.17 and the fact that

$$
\begin{aligned}
&\mathcal{M}^1_{X^{(1)}}(v^{(1)}, w^{(1)}) - \mathcal{M}^1_{X^{(2)}}(v^{(2)}, w^{(2)}) \\
&= P_t^1 (v_0^{(1)} - v_0^{(2)}) + \int_0^t P_{t-s}^1 \{ F_{X^{(1)}}(v^{(1)}, w^{(1)})(s) - F_{X^{(2)}}(v^{(2)}, w^{(2)})(s) \} \, ds.
\end{aligned}
$$

By a similar argument to the proof of Proposition 4.16, we see the conclusion. □

4.3.2 Properties of $\mathcal{M}^2$

Here, we consider properties of $\mathcal{M}^2$. Let $0 < T \le 1$. We fix $X \in \mathcal{X}_1^\kappa$ and write $Z = X^{\text{𝍐}}$ and $W = X^{\text{𝍖}}$. We denote by δ_{st} the difference operator, that is, $\delta_{st} f = f(t) - f(s)$.

First of all, we study $\mathrm{com}(v, w)$ defined by (4.6). Let $v_0 \in \mathcal{C}^{-\frac{2}{3}+\kappa'}$ and $(v, w) \in \mathcal{D}_T^{\kappa,\kappa'}$. For notational simplicity, we set $\Phi^1 = -2\nu(-\nu W + v + w)$, $\Phi^2 = -\nu(-\overline{\nu}\overline{W} + \overline{v} + \overline{w})$, $\Psi^1 = X^{\text{𝍗}}$, $\Psi^2 = X^{\text{𝍘}}$ and

$$
U_t = \int_0^t P_{t-s}^1 [F(v, w)(s)] \, ds - \Phi_t^1 \circledcirc X_t^{\text{𝍙}} - \Phi_t^2 \circledcirc X_t^{\text{𝍚}}.
$$

Remark 4.19. The implicit constants which will appear in Lemma 4.20, 4.21 and 4.22 depend only on κ, κ', μ, ν and $\|X\|_{\mathcal{X}_1^\kappa}$. In particular, the constants are given by an at most first-order polynomials in $\|X\|_{\mathcal{X}_1^\kappa}$.

Lemma 4.20. *For every $0 < t \le T$, we have the following:*

(1) We have

$$U_t = -\Phi_t^1 \circledcirc P_t^1 X_0^{\Upsilon^{\bullet}} - \Phi_t^2 \circledcirc P_t^2 X_0^{\Upsilon} + \sum_{i=1,2} \left\{ \int_0^t \delta_{st}\Phi^i \circledcirc P_{t-s}^1 \Psi_s^i \, ds - \int_0^t [P_{t-s}^1, \Phi_s^i \circledcirc] \Psi_s^i \, ds \right\}.$$

(2) We have $U_t \in \mathcal{C}^{1+\kappa'}$

$$\begin{aligned} \|U_t\|_{\mathcal{C}^{1+\kappa'}} &\lesssim 1 + t^{-\kappa'} \{1 + \|v_t\|_{L^\infty} + \|w_t\|_{L^\infty}\} \\ &\quad + \int_0^t (t-s)^{-\frac{3+2\kappa}{4}} \|v_s\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \, ds + \int_0^t (t-s)^{-\frac{1+2\kappa'}{2}} \|w_s\|_{\mathcal{C}^{1+2\kappa'}} \, ds \\ &\quad + \int_0^t (t-s)^{-\frac{2+\kappa+\kappa'}{2}} \{\|\delta_{st} v\|_{L^p} + \|\delta_{st} w\|_{L^p}\} \, ds. \end{aligned}$$

Proof. We show the first assertion. For $(\Phi, \Psi) = (\Phi^1, \Psi^1)$, Proposition 4.9 implies

$$\begin{aligned} P_{t-s}^1[\Phi \circledcirc \Psi](s) &= P_{t-s}^1[\Phi_s \circledcirc \Psi_s] \\ &= \Phi_s \circledcirc P_{t-s}^1 \Psi_s + [P_{t-s}^1, \Phi_s \circledcirc] \Psi_s \\ &= \Phi_t \circledcirc P_{t-s}^1 \Psi_s - \delta_{st}\Phi \circledcirc P_{t-s}^1 \Psi_s + [P_{t-s}^1, \Phi_s \circledcirc] \Psi_s. \end{aligned}$$

Hence

$$\begin{aligned} &\int_0^t P_{t-s}^1[\Phi \circledcirc \Psi](s) \, ds \\ &\quad = \Phi_t \circledcirc \int_0^t P_{t-s}^1 \Psi_s \, ds - \int_0^t \delta_{st}\Phi \circledcirc P_{t-s}^1 \Psi_s \, ds + \int_0^t [P_{t-s}^1, \Phi_s \circledcirc] \Psi_s \, ds. \end{aligned}$$

Substituting $\int_0^t P_{t-s}^1 \Psi_s \, ds = X_t^{\Upsilon^{\bullet}} - P_t^1 X_0^{\Upsilon^{\bullet}}$ to the first term in the above, we see

$$\begin{aligned} &\int_0^t P_{t-s}^1[\Phi \circledcirc \Psi](s) \, ds - \Phi_t \circledcirc X_t^{\Upsilon^{\bullet}} \\ &\quad = -\Phi_t \circledcirc P_t^1 X_0^{\Upsilon^{\bullet}} - \int_0^t \delta_{st}\Phi \circledcirc P_{t-s}^1 \Psi_s \, ds + \int_0^t [P_{t-s}^1, \Phi_s \circledcirc] \Psi_s \, ds. \end{aligned} \tag{4.10}$$

Since a similar equality holds for $(\Phi, \Psi) = (\Phi^2, \Psi^2)$, we have verified the first assertion.

For the second assertion, we estimate the terms in (4.10). For the first term in (4.10), we use $1 + \kappa' = 1 - \kappa + 2 \cdot \frac{\kappa + \kappa'}{2}$ and obtain

$$\|\Phi_t^1 \circledcirc P_t^1 X_0^{\Upsilon^{\bullet}}\|_{\mathcal{C}^{1+\kappa'}} \lesssim \|\Phi_t^1\|_{L^\infty} \|P_t^1 X_0^{\Upsilon^{\bullet}}\|_{\mathcal{C}^{1+\kappa'}}$$

$$\lesssim \| - \nu W_t + v_t + w_t\|_{L^\infty} \cdot t^{-\frac{\kappa+\kappa'}{2}} \|X_0^{\blacktriangledown}\|_{\mathcal{C}^{1-\kappa}}$$
$$\lesssim t^{-\kappa'}\{1 + \|v_t\|_{L^p} + \|w_t\|_{L^p}\}.$$

We estimate the second term in (4.10). From $1 + \kappa' = -1 - \kappa + 2 \cdot \frac{2+\kappa+\kappa'}{2}$, we have

$$\begin{aligned}\|\delta_{st}\Phi \circledcirc P^1_{t-s}\Psi_s\|_{\mathcal{C}^{1+\kappa'}} &\lesssim \|\delta_{st}\Phi\|_{L^\infty}\|P^1_{t-s}\Psi_s\|_{\mathcal{C}^{1+\kappa'}} \\ &\lesssim \|\delta_{st}\Phi\|_{L^\infty} \cdot (t-s)^{-\frac{2+\kappa+\kappa'}{2}}\|\Psi_s\|_{\mathcal{C}^{-1-\kappa}} \\ &\lesssim (t-s)^{-\frac{2+\kappa+\kappa'}{2}}\|\delta_{st}\Phi\|_{L^p}.\end{aligned}$$

Note

$$\|\delta_{st}\Phi\|_{L^\infty} \lesssim \|\delta_{st}v\|_{L^\infty} + \|\delta_{st}w\|_{L^\infty} + \|\delta_{st}W\|_{L^\infty},$$
$$\|\delta_{st}W\|_{L^\infty} \lesssim \|\delta_{st}W\|_{\mathcal{C}^{\kappa'-\kappa}} \lesssim (t-s)^{\frac{1}{4}-\frac{1}{2}\kappa'}\|W\|_{C_T^{\frac{1}{4}-\frac{1}{2}\kappa'}\mathcal{C}^{\kappa'-\kappa}}.$$

For the latter estimate, see Remark 4.14. From them, we have

$$\begin{aligned}&\left\|\int_0^t \delta_{st}\Phi \circledcirc P^1_{t-s}\Psi_s\, ds\right\|_{\mathcal{C}^{1+\kappa'}} \\ &\qquad\lesssim \int_0^t (t-s)^{-\frac{2+\kappa+\kappa'}{2}}\{\|\delta_{st}v\|_{L^\infty} + \|\delta_{st}w\|_{L^\infty} + (t-s)^{\frac{1}{4}-\frac{1}{2}\kappa'}\}\, ds \\ &\qquad\lesssim \int_0^t (t-s)^{-\frac{2+\kappa+\kappa'}{2}}\{\|\delta_{st}v\|_{L^\infty} + \|\delta_{st}w\|_{L^\infty}\}\, ds + 1.\end{aligned}$$

The last inequality follows from $\int_0^t (t-s)^{-1-\frac{\kappa+\kappa'}{2}+\frac{1}{4}-\frac{1}{2}\kappa'}\, ds < \infty$. The estimate of the second term has finished.

Lastly, we estimate the third term. We consider the contribution of W, v and w separably. In the proof, we use Proposition 4.9. Note

$$\begin{aligned}1 + \kappa' &= \left(\frac{1}{2} - \kappa\right) + (-1-\kappa) + 2 \cdot \frac{3 + 4\kappa + 2\kappa'}{4} \\ &= \left(\frac{1}{2} + \kappa'\right) + (-1-\kappa) + 2 \cdot \frac{3 + 2\kappa}{4} \\ &= (1 - \kappa' + \kappa) + (-1-\kappa) + 2 \cdot \frac{1 + 2\kappa'}{2}.\end{aligned}$$

We also use $\|\Psi_s\|_{\mathcal{C}^{-1-\kappa}} \lesssim 1$. For W, we see

$$\begin{aligned}\left\|\int_0^t [P^1_{t-s}, W_s\circledcirc]\Psi_s\, ds\right\|_{\mathcal{C}^{1+\kappa'}} &\lesssim \int_0^t (t-s)^{-\frac{3+4\kappa+2\kappa'}{4}}\|W_s\|_{\mathcal{C}^{\frac{1}{2}-\kappa}}\|\Psi_s\|_{\mathcal{C}^{-1-\kappa}}\, ds \\ &\lesssim 1.\end{aligned}$$

For v and w, we have

$$\begin{aligned}\left\|\int_0^t [P^1_{t-s}, v_s\circledcirc]\Psi_s\,ds\right\|_{\mathcal{C}^{1+\kappa'}} &\lesssim \int_0^t (t-s)^{-\frac{3+2\kappa}{4}}\|v_s\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}}\,ds,\\ \left\|\int_0^t [P^1_{t-s}, w_s\circledcirc]\Psi_s\,ds\right\|_{\mathcal{C}^{1+\kappa'}} &\lesssim \int_0^t (t-s)^{-\frac{1+2\kappa'}{2}}\|w_s\|_{\mathcal{C}^{1-\kappa'+\kappa}}\,ds\\ &\lesssim \int_0^t (t-s)^{-\frac{1+2\kappa'}{2}}\|w_s\|_{\mathcal{C}^{1+2\kappa'}}\,ds.\end{aligned}$$

The proof is completed. □

Lemma 4.21. *For any $(v,w)\in\mathcal{D}_T^{\kappa,\kappa'}$ and $0<t\le T$, we have $\operatorname{com}(v,w)(t)\in\mathcal{C}^{1+\kappa'}$ and*

$$\begin{aligned}&\|\operatorname{com}(v,w)(t)\|_{\mathcal{C}^{1+\kappa'}}\\ &\quad\lesssim 1+t^{-\frac{5}{6}}\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+t^{-\kappa'}(1+\|v_t\|_{L^\infty}+\|w_t\|_{L^\infty})\\ &\qquad+\int_0^t (t-s)^{-\frac{3+2\kappa'}{4}}\|v_s\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}}\,ds+\int_0^t (t-s)^{-\frac{1+2\kappa'}{2}}\|w_s\|_{\mathcal{C}^{1+2\kappa'}}\,ds\\ &\qquad+\int_0^t (t-s)^{-1-\frac{\kappa+\kappa'}{2}}(\|\delta_{st}v\|_{L^\infty}+\|\delta_{st}w\|_{L^\infty})\,ds.\end{aligned}$$

Proof. From definition (4.6), we have $\operatorname{com}(v,w)(t)=P^1_tv_0+U_t$.

Noting $1+\kappa'=\left(-\frac{2}{3}+\kappa'\right)+2\cdot\frac{5}{6}=\frac{1}{2}+\kappa'+2\cdot\frac{1}{4}$ and using Proposition 4.7, we see

$$\begin{aligned}\|P^1_tv_0\|_{\mathcal{C}^{1+\kappa'}}&\lesssim t^{-\frac{5}{6}}\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}},\\ \left\|\int_0^t P^1_{t-s}v_s\,ds\right\|_{\mathcal{C}^{1+\kappa'}}&\le\int_0^t\|P^1_{t-s}v_s\|_{\mathcal{C}^{1+\kappa'}}\,ds\lesssim\int_0^t(t-s)^{-\frac{1}{4}}\|v_s\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}}\,ds.\end{aligned}$$

Note that the last term smaller than or equal to $\int_0^t(t-s)^{-\frac{3+2\kappa}{4}}\|v_s\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}}\,ds$. Combining these and Lemma 4.20, we see the conclusion. □

Lemma 4.22. *For any $(v,w)\in\mathcal{D}_T^{\kappa,\kappa'}$ and $0<t\le T$, we have*

$$\|\operatorname{com}(v,w)(t)\|_{\mathcal{C}^{1+\kappa'}}\lesssim C(1+t^{-\frac{5}{6}}\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+t^{-\frac{1+2\kappa+2\kappa'}{2}}\|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}).$$

Proof. We estimate each term in the upper bound of $\|\operatorname{com}(v,w)(t)\|_{\mathcal{C}^{1+\kappa'}}$ in Lemma 4.21 by using Remark 4.14. The first three terms are estimated as follows:

$$\begin{aligned}&1+t^{-\frac{5}{6}}\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+t^{-\kappa'}(1+\|v(t)\|_{L^\infty}+\|w(t)\|_{L^\infty})\\ &\qquad\lesssim 1+t^{-\frac{5}{6}}\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+t^{-\left(\kappa'+\frac{2-3\kappa}{6}\right)}(1+\|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}).\end{aligned}$$

To estimate other terms, we use the fact that the inequality

$$\int_0^t (t-s)^{-\theta_1} s^{-\theta_2}\, ds \lesssim t^{1-\theta_1-\theta_2}$$

holds for $0 < \theta_1, \theta_2 < 1$ and $t > 0$. From Remark 4.14, we see

$$\begin{aligned}
&\int_0^t (t-s)^{-\frac{3+2\kappa'}{4}} \|v_s\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}}\, ds + \int_0^t (t-s)^{-\frac{1+2\kappa'}{2}} \|w_s\|_{\mathcal{C}^{1+2\kappa'}}\, ds \\
&\qquad \lesssim t^{-\left(\frac{1}{3}+\frac{1}{2}\kappa'\right)} \|v\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}} + t^{-\left(\frac{1}{4}+2\kappa'+\kappa\right)} \|w\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}}
\end{aligned}$$

and

$$\begin{aligned}
&\int_0^t (t-s)^{-\left(1+\frac{\kappa+\kappa'}{2}\right)} (\|\delta_{st} v\|_{L^\infty} + \|\delta_{st} w\|_{L^\infty})\, ds \\
&\qquad \lesssim t^{-\frac{2+3\kappa+3\kappa'}{6}} \|v\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}} + t^{-\frac{1+3\kappa+\kappa'}{2}} \|w\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}}.
\end{aligned}$$

Combining them, we see the estimate of $\|\operatorname{com}(v,w)(t)\|_{\mathcal{C}^{1+\kappa'}}$. □

Lemma 4.23. *For any $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$ and $0 < t \le T$, we have*

$$\begin{aligned}
&\|G(v,w)(t)\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}} \\
&\qquad \le C\Big(1 + t^{-\frac{5}{6}} \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + t^{-\frac{2-3\kappa}{2}} \Big(\|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^3 + \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}\Big)\Big).
\end{aligned}$$

Here, C is a positive constant depending only on κ, κ', μ, ν and $\|X\|_{\mathcal{X}_1^\kappa}$ and it is given by a third-order polynomial in $\|X\|_{\mathcal{X}_1^\kappa}$.

Proof. We write $u_2 = v + w$. It follows from Remark 4.14 that

$$\|G_1(v,w)(t)\|_{L^\infty} \lesssim \|v_t + w_t\|_{L^\infty}^3 \lesssim t^{-\frac{2-3\kappa}{2}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^3.$$

To estimate $G_2(v,w)(t)$, we use the Bony decomposition. Applying the decomposition to $u_2(t) \in \mathcal{C}^{\frac{1}{2}+\kappa'}$, we see

$$\begin{aligned}
\|u_2(t)^2\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} &= \|u_2(t) \odot u_2(t) + 2u_2(t) \oslash u_2(t)\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \\
&\lesssim \|u_2(t)\|_{\mathcal{C}^{\frac{1}{2}\left(\frac{1}{2}+\kappa'\right)}}^2 + 2\|u_2(t)\|_{L^\infty} \|u_2(t)\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \\
&\lesssim t^{-\frac{11-6\kappa'}{12}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^2 + 2t^{-\frac{2-3\kappa}{6}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}} \cdot t^{-\frac{7}{12}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}} \\
&= t^{-\frac{11-6\kappa'}{12}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^2 + 2t^{-\frac{11-6\kappa}{12}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^2 \\
&\lesssim t^{-\frac{11-6\kappa}{12}} \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^2.
\end{aligned}$$

In these estimate, we used $0 < \kappa < \kappa' < 1/18$ (see Remark 4.14). The term $\|u_2(t)\overline{u_2(t)}\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}}$ has the same bound. Since $G_2(v,w)(t) = a_1 u_2(t)^2 + a_2 u_2(t)\overline{u_2(t)}$ for some $a_1, a_2 \in \mathcal{C}^{-\frac{1}{2}-\kappa}$ and $u_2(t) \in \mathcal{C}^{\frac{1}{2}+\kappa'}$, we have

$$\begin{aligned}\|G_2(v,w)(t)\|_{\mathcal{C}^{-\frac{1}{2}-\kappa}} &\lesssim \|u_2(t)^2\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} + \|u_2(t)\overline{u_2(t)}\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \\ &\lesssim t^{-\frac{11-6\kappa}{12}} \|(v,w)\|^2_{\mathcal{D}^{\kappa,\kappa'}_T}.\end{aligned}$$

Noting $\|(v,w)\|^2_{\mathcal{D}^{\kappa,\kappa'}_T} \le \|(v,w)\|^3_{\mathcal{D}^{\kappa,\kappa'}_T} + \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}$, we see the estimate.

Since $G_3(v,w)(t) = b_1 u_2(t) + b_2\overline{u_2(t)} + (\nu+1)u_2(t)$ for some $b_1, b_2 \in \mathcal{C}^{-\frac{1}{2}-\kappa}$, we have

$$\|G_3(v,w)(t)\|_{\mathcal{C}^{-\frac{1}{2}-\kappa}} \lesssim \|u_2(t)\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \lesssim t^{-\frac{7}{12}} \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}.$$

The estimates of $G_4(v,w)(t)$ and $G_5(v,w)(t)$ are obtained easily. The terms which admits the lowest regularity in the defintions of $G_4(v,w)(t)$ are $W_t \circledcirc X^{\mathbb{V}}$ and $\overline{W_t} \circledcirc X^{\mathbb{V}}$ and their regularity is $-\frac{1}{2}-2\kappa$. Therefore we obtain $\|G_4(v,w)(t)\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}} \lesssim 1$. From Proposition 4.5, we see

$$\|G_5(v,w)(t)\|_{\mathcal{C}^{\frac{1}{2}+\kappa'-2\kappa}} \lesssim \|u_2\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \lesssim t^{-\frac{7}{12}} \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}.$$

From the definition of $G_6(v,w)(t)$, we have

$$\begin{aligned}\|G_6(v,w)(t)\|_{\mathcal{C}^{\kappa'-\kappa}} &\lesssim \|\operatorname{com}(v,w)(t) \odot X^{\mathbb{V}}_t\|_{\mathcal{C}^{1+\kappa'+(-1-\kappa)}} \\ &\qquad + \|\overline{\operatorname{com}(v,w)(t)} \odot X^{\mathbb{V}}_t\|_{\mathcal{C}^{1+\kappa'+(-1-\kappa)}} \\ &\lesssim \|\operatorname{com}(v,w)(t)\|_{\mathcal{C}^{1+\kappa'}} \\ &\lesssim C(1 + t^{-\frac{5}{6}}\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + t^{-\frac{1+2\kappa+2\kappa'}{2}}\|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}).\end{aligned}$$

In the last line, we used Lemma 4.22.

For $\tau = \mathbb{V}, \mathbb{V}$, we see

$$\begin{aligned}\|w_t \odot X^\tau_t\|_{\mathcal{C}^{(1+\kappa')+(-1-\kappa)}} &\lesssim \|w_t\|_{\mathcal{C}^{1+\kappa'}} \|X^\tau_t\|_{\mathcal{C}^{-1-\kappa}} \lesssim t^{-\frac{3+2\kappa+\kappa'}{4}} \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}, \\ \|u_2(t) \circledcirc X^\tau_t\|_{\mathcal{C}^{(\frac{1}{2}+\kappa')+(-1-\kappa)}} &\lesssim \|u_2(t)\|_{\mathcal{C}^{\frac{1}{2}+\kappa'}} \|X^\tau_t\|_{\mathcal{C}^{-1-\kappa}} \lesssim t^{-\frac{7}{12}} \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}.\end{aligned}$$

In these estimates, we used Remark 4.14 . We obtain

$$\begin{aligned}\|G_7(v,w)(t)\|_{\mathcal{C}^{\kappa'-\kappa}} &\lesssim t^{-\frac{3+4\kappa+2\kappa'}{4}} \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}, \\ \|G_8(v,w)(t)\|_{\mathcal{C}^{-\frac{1}{2}+\kappa'-\kappa}} &\lesssim t^{-\frac{7}{12}} \|(v,w)\|_{\mathcal{D}^{\kappa,\kappa'}_T}.\end{aligned}$$

The proof is completed. □

Proposition 4.24. *The map $\mathcal{M}^2 : \mathcal{D}_T^{\kappa,\kappa'} \to \mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}$ is well-defined and, for any $(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have*

$$\begin{aligned}\|\mathcal{M}^2(v,w)\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}} &\le C_1(1+\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+\|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}})\\ &\quad + C_2T^{\frac{3}{2}\kappa}\left(\|(v,w)\|^3_{\mathcal{D}_T^{\kappa,\kappa'}}+\|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}\right).\end{aligned}$$

Here, C_1 and C_2 are positive constants depending only on κ, κ', μ, ν and $\|X\|_{\mathcal{X}_1^\kappa}$. They are given by at most third-order polynomials in $\|X\|_{\mathcal{X}_1^\kappa}$.

Proof. Recall (4.8). It follows from Proposition 4.8 that

$$\|(t\mapsto P_t^1 w_0)_{t\ge0}\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}} \lesssim \|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}}.$$

Lemma 4.23 implies $G(v,w) \in \mathcal{E}_T^{\frac{2-3\kappa}{2}}\mathcal{C}^{-\frac{1}{2}-2\kappa}$ and

$$\|G(v,w)\|_{\mathcal{E}_T^{\frac{2-3\kappa}{2}}\mathcal{C}^{-\frac{1}{2}-2\kappa}} \le C\Big(1+\|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+\|(v,w)\|^3_{\mathcal{D}_T^{\kappa,\kappa'}}+\|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}\Big).$$

Proposition 4.8 implies

$$\begin{aligned}\left\|(t\mapsto\int_0^t P_{t-s}^1 G(v,w)(s)\,ds)_{t\ge0}\right\|_{\mathcal{L}_T^{1-\kappa'+\kappa,\frac{3}{2}-2\kappa',1-\kappa'}}&\\ \lesssim T^{\frac{3}{2}\kappa}\|G(v,w)\|_{\mathcal{E}_T^{\frac{2-3\kappa}{2}}\mathcal{C}^{-\frac{1}{2}-2\kappa}}.&\end{aligned}$$

Combining these, we have shown the assertion. □

We also have local Lipschitz continuity of $\mathcal{M}_2$.

Proposition 4.25. *For any $(v^{(1)},w^{(1)}),(v^{(2)},w^{(2)}) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have*

$$\begin{aligned}\|\mathcal{M}^2_{(v_0^{(1)},w_0^{(1)}),X^{(1)}}(v^{(1)},w^{(1)})-\mathcal{M}^2_{(v_0^{(2)},w_0^{(2)}),X^{(2)}}(v^{(2)},w^{(2)})\|_{\mathcal{L}_T^{\frac{5}{6}-\kappa',1-\kappa',1-\frac{1}{2}\kappa'}}&\\ \le C_3\left(\|v_0^{(1)}-v_0^{(2)}\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+\|w_0^{(1)}-w_0^{(2)}\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}}\right)&\\ + C_4T^{\frac{3}{2}\kappa}\Big(\|X^{(1)}-X^{(2)}\|_{\mathcal{X}_T^\kappa}+\|(v^{(1)},w^{(1)})-(v^{(2)},w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}\Big)&\end{aligned}$$

Here, C_3 and C_4 are positive constants depending only on κ, κ', μ, ν, $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)},w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$. In particular, they are given by at most second-order polynomials in $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)},w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$.

Proof. We can show the assertion by a similar way as Proposition 4.24. □

4.4 Local existence and uniqueness

We show local well-posedness of CGL (1.1). This is the most important theorem in this section.

Theorem 4.26. *Let $0 < \kappa < \kappa' < 1/18$. There exists a continuous function $\tilde{T}_* : \mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa} \times \mathcal{X}_1^\kappa \to (0,1]$ such that the following (1) and (2) hold:*

(1) For every $(v_0, w_0) \in \mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa}$ and $X \in \mathcal{X}_1^\kappa$, set $T_* = \tilde{T}_*(v_0, w_0, X)$. Then, the system (4.9) admits a unique solution $(v, w) \in \mathcal{D}_{T_*}^{\kappa,\kappa'}$ and there is a positive constant C depending only on μ, ν, κ, κ', T_* and $\|X\|_{\mathcal{X}_1^\kappa}$ such that

$$\|(v,w)\|_{\mathcal{D}_{T_*}^{\kappa,\kappa'}} \le C\left(1 + \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}}\right).$$

(2) Let $\{(v_0^{(n)}, w_0^{(n)})\}_{n=1}^\infty$ and $\{X^{(n)}\}_{n=1}^\infty$ converge to (v_0, w_0) in $\mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa}$ and X in $\mathcal{X}_1^\kappa$, respectively. Set $T_*^{(n)} = \tilde{T}_*(v_0^{(n)}, w_0^{(n)}, X^{(n)})$ and let $(v^{(n)}, w^{(n)})$ be a unique solution on $[0, T_*^{(n)}]$ to the system (4.9) with the initial condition $(v_0^{(n)}, w_0^{(n)})$ driven by $X^{(n)}$. Then, for every $0 < t < T_*$, we have

$$\lim_{n\to\infty} \|(v^{(n)}, w^{(n)}) - (v,w)\|_{\mathcal{D}_t^{\kappa,\kappa'}} = 0.$$

In the proof the function $\tilde{T}_*$ is concretely given by $\tilde{T}_*(v_0, w_0, X) = T_*$, where T_* is defined by (4.12) and (4.13). We prove the theorem by using the properties of $\mathcal{M}$ we have just shown.

For every $0 < T \le 1$ and $M > 0$, we define

$$\mathbf{B}_{T,M} = \{(v,w) \in \mathcal{D}_T^{\kappa,\kappa'}; \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}} \le M\}.$$

Propositions 4.16 and 4.24 imply

$$\begin{aligned}\|\mathcal{M}(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}} \le C_1\big(1 + \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}}\big) \\ + C_2 T^{\frac{3}{2}\kappa}(\|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}^3 + \|(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}}).\end{aligned}$$

Here, C_1 and C_2 are positive constants depending only on κ, κ', μ, ν and $\|X\|_{\mathcal{X}_1^\kappa}$. In particular, they are given by at most third-order polynomials in $\|X\|_{\mathcal{X}_1^\kappa}$. Propositions 4.18 and 4.25 imply

$$\begin{aligned}(4.11)\quad &\|\mathcal{M}_{(v_0^{(1)}, w_0^{(1)}), X^{(1)}}(v^{(1)}, w^{(1)}) - \mathcal{M}_{(v_0^{(2)}, w_0^{(2)}), X^{(2)}}(v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \\ &\le C_3(\|v_0^{(1)} - v_0^{(2)}\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w_0^{(1)} - w_0^{(2)}\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}})\end{aligned}$$

$$+ C_4 T^{\frac{3}{2}\kappa}(\|X^{(1)} - X^{(2)}\|_{\mathcal{X}_T^\kappa} + \|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}).$$

Here, C_3 and C_4 are positive constants depending only on κ, κ', μ, ν, $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)}, w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$. In particular, they are given by at most second-order polynomials in $\|X^{(i)}\|_{\mathcal{X}_1^\kappa}$ and $\|(v^{(i)}, w^{(i)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}$.

Proof of Theorem 4.26. For the proof of existence, we use Propositions 4.16 and 4.24. We will show the map $\mathcal{M}$ is contraction from $\mathbf{B}_{T,M}$ to itself for small $T > 0$ and suitable $M > 0$ and obtain the existence of solution by the fixed point theorem. Let $M \geq 1$. For any $(v, w) \in \mathcal{D}_T^{\kappa,\kappa'}$, we have

$$\begin{aligned}\|\mathcal{M}(v,w)\|_{\mathcal{D}_T^{\kappa,\kappa'}} &\leq (C_1 + C_2)(1 + \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}} + T^{\frac{3}{2}\kappa}M^3) \\ &\leq (C_1 + C_2)(1 + \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}})(1 + T^{\frac{3}{2}\kappa}M^3).\end{aligned}$$

In a similar way, we see

$$\begin{aligned}&\|\mathcal{M}(v^{(1)}, w^{(1)}) - \mathcal{M}(v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}} \\ &\qquad \leq C_{\|X\|_{\mathcal{X}_1^\kappa}} T^{\frac{3}{2}\kappa}(1 + M^2)\|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}.\end{aligned}$$

Here $C_{\|X\|_{\mathcal{X}_1^\kappa}} > 0$ is given by a second-order polynomial with respect to $\|X\|_{\mathcal{X}_T^\kappa}$. Set

$$M_* = 2(C_1 + C_2)(1 + \|v_0\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w_0\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}}) \vee 1, \tag{4.12}$$

$$T_*^{\frac{3}{2}\kappa} = M_*^{-3} \wedge \{2C_{\|X\|_{\mathcal{X}_1^\kappa}}(1 + M_*^2)\}^{-1} \wedge 1. \tag{4.13}$$

Then

$$\|\mathcal{M}(v,w)\|_{\mathcal{D}_{T_*}^{\kappa,\kappa'}} \leq M_*, \tag{4.14}$$

(4.15)

$$\|\mathcal{M}(v^{(1)}, w^{(1)}) - \mathcal{M}(v^{(2)}, w^{(2)})\|_{\mathcal{D}_{T_*}^{\kappa,\kappa'}} \leq \frac{1}{2}\|(v^{(1)}, w^{(1)}) - (v^{(2)}, w^{(2)})\|_{\mathcal{D}_{T_*}^{\kappa,\kappa'}}.$$

We see that the map $\mathcal{M}$ is contraction on $\mathbf{B}_{T_*,M_*}$ Therefore there exists a unique fixed point $(v, w) \in \mathbf{B}_{T_*,M_*}$ of $\mathcal{M}$, which is a solution on $[0, T_*]$.

Next we show that the solution on $[0, T_*]$ is unique. Let $(v^{(1)}, w^{(1)}), (v^{(2)}, w^{(2)}) \in \mathcal{D}_{T_*}^{\kappa,\kappa'}$ are solutions with a common initial condition (v_0, w_0). We show that $(v^{(1)}, w^{(1)}) = (v^{(2)}, w^{(2)})$. Taking $M > 0$ such that

$$\|(v^{(1)}, w^{(1)})\|_{\mathcal{D}_{T_*}^{\kappa,\kappa'}} \vee \|(v^{(2)}, w^{(2)})\|_{\mathcal{D}_{T_*}^{\kappa,\kappa'}} \leq M,$$

the similar arguments as above ensure that $\mathcal{M}$ is a contraction on $\mathbf{B}_{T_{**},M}$, where $T_{**}(\le T_*)$ depends on M. Hence $(v^{(1)},w^{(1)})$ and $(v^{(2)},w^{(2)})$ coincide on $[0,T_{**}]$. We can continue this procedure on $[T_{**},2T_{**}]$, $[2T_{**},3T_{**}],\dots$. However, in these steps, we need to check that $(\tilde{v}^{(i)},\tilde{w}^{(i)})(t)=(v^{(i)},w^{(i)})(t+T_{**})$ satisfies

$$\|(\tilde{v}^{(i)},\tilde{w}^{(i)})\|_{\mathcal{D}^{\kappa,\kappa'}_{T_*-T_{**}}}\le M,$$

since for example

$$\begin{aligned}\|\tilde{v}^{(i)}\|_{\mathcal{E}^{\frac{5}{6}-\kappa'}_{T_*-T_{**}}\mathcal{C}^{1-\kappa'}}&=\sup_{T_{**}<t\le T_*}(t-T_{**})^{\frac{5}{6}-\kappa'}\|v^{(i)}(t)\|_{\mathcal{C}^{1-\kappa'}}\\&\le\sup_{0<t\le T_*}t^{\frac{5}{6}-\kappa'}\|v^{(i)}(t)\|_{\mathcal{C}^{1-\kappa'}}\\&=\|v^{(i)}\|_{\mathcal{E}^{\frac{5}{6}-\kappa'}_{T_*}\mathcal{C}^{1-\kappa'}}.\end{aligned}$$

Obviously $(\tilde{v}^{(i)},\tilde{w}^{(i)})$ is a solution with the initial condition $(v^{(1)},w^{(1)})(T_{**})=(v^{(2)},w^{(2)})(T_{**})$. Therefore we can iterate the above arguments on $[kT_*,(k+1)T_{**}\wedge T]$ for $k=1,2,\dots$ and thus $(v^{(1)},w^{(1)})$ and $(v^{(2)},w^{(2)})$ coincide on $[0,T_*]$.

We show the last assertion. From (4.12) and (4.13), we see that T_* continuously depends on the initial condition (v_0,w_0) and the driving vector X. Since C_2 depends on the driving vector X continuously, M_* is a continuous map from (v_0,w_0) and X. From this fact and the continuity of $C_{\|X\|_{\mathcal{X}_1^\kappa}}$, we see the continuity of T_*. Hence we have $T_*^{(n)}\to T_*$. Without loss of generality, for fixed $t<T_*$, we assume that $T_*^{(n)}>t$ for every n. From (4.14) and the continuity of M_* with respect to (v_0,w_0) and X, we see $\sup_n\|(v^{(n)},w^{(n)})\|_{\mathcal{D}_T^{\kappa,\kappa'}}<\infty$. From this fact and (4.11), we can choose C_3' and C_4' such that

$$\begin{aligned}\|(v,w)-(v^{(n)},w^{(n)})\|_{\mathcal{D}_t^{\kappa,\kappa'}}\le{}&C_3'\Big(\|v_0-v_0^{(n)}\|_{\mathcal{C}^{-\frac{2}{3}+\kappa'}}+\|w_0-w_0^{(n)}\|_{\mathcal{C}^{-\frac{1}{2}-2\kappa}}\Big)\\&+C_4't^{\frac{3}{2}\kappa}\Big(\|X-X^{(n)}\|_{\mathcal{X}_1^\kappa}+\|(v,w)-(v^{(n)},w^{(n)})\|_{\mathcal{D}_t^{\kappa,\kappa'}}\Big).\end{aligned}$$

Hence we have $(v^{(n)},w^{(n)})\to(v,w)$ in $[0,t_*]$ for some $t_*\le t$ depending on C_3' and C_4'. Iterating this argument, we have the convergence in $[0,t]$. The proof is completed. □

Remark 4.27. If $(v_0,w_0)\in\mathcal{C}^{1-\kappa'}\times\mathcal{C}^{\frac{3}{2}-2\kappa'}$, we obtain the local well-posedness on the space $\mathcal{L}_T^{1-\kappa',1-\frac{\kappa'}{2}}\times\mathcal{L}_T^{\frac{3}{2}-\kappa',1-\kappa'}$ without explosion at $t=0$ by similar arguments.

Proposition 4.28. *For every $(v_0,w_0)\in\mathcal{C}^{-\frac{2}{3}+\kappa'}\times\mathcal{C}^{-\frac{1}{2}-2\kappa}$ and $X\in\mathcal{X}_T^\kappa$, there exists $T_{\mathrm{sur}}\in(0,\infty]$ such that the system (4.9) has a unique solution $(v,w)\in$*

$\mathcal{D}_t^{\kappa,\kappa'}$ *for every* $t < T_{\mathrm{sur}}$*, and*

$$\lim_{t\uparrow T_{\mathrm{sur}}} (\|v\|_{C_t\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w\|_{C_t\mathcal{C}^{-\frac{1}{2}-2\kappa}}) = \infty$$

unless $T_{\mathrm{sur}} = \infty$*. Furthermore, the mapping from* (v_0, w_0, X) *to the maximal solution* (v, w) *is continuous in the sense that, for a sequence* $\{(v_0^{(n)}, w_0^{(n)}, X^{(n)})\}$ *which converges to* (v_0, w_0, X)*, we have* $T_{\mathrm{sur}} \le \liminf_{n\to\infty} T_{\mathrm{sur}}^{(n)}$ *and*

$$\|(v^{(n)}, w^{(n)}) - (v, w)\|_{\mathcal{D}_t^{\kappa,\kappa'}} \to 0$$

for every $t < T_{\mathrm{sur}}$*.*

Proof. Let $(v, w) \in \mathcal{D}_{T_*}^{\kappa,\kappa'}$ be a unique solution on $[0, T_*]$ shown in Theorem 4.26 Because of Remark 4.27, we can start from $(v, w)(T_*) \in \mathcal{C}^{1-\kappa'} \times \mathcal{C}^{\frac{3}{2}-2\kappa'}$ and construct a solution $(\bar{v}, \bar{w}) \in \mathcal{D}_{T_{**}}^{\kappa,\kappa'}$ with $(\bar{v}, \bar{w})(0) = (v, w)(T_*)$. Obviously the extended function

$$(\hat{v}, \hat{w})(t) = \begin{cases} (v, w)(t) & t \in [0, T_*] \\ (\bar{v}, \bar{w})(t - T_*) & t \in [T_*, T_* + T_{**}] \end{cases}$$

belongs to $\mathcal{D}_{T_*+T_{**}}^{\kappa,\kappa'}$ and solves the system (4.12). Uniqueness on $[0, T_*+T_{**}]$ also holds. We can iterate this argument until the time T_{sur}, which is a supremum up to when the existence and uniqueness hold.

The lower semi-continuity of T_{sur} follows from the continuity of T_*. Let $(v_0^{(n)}, w_0^{(n)}, X^{(n)}) \to (v_0, w_0, X)$. For any fixed $t < T_{\mathrm{sur}}$, we can construct a unique solution in $[0, t]$ by gluing finite number of local solutions as above. In this procedure, each of length of time interval converges, so that the solution $(v^{(n)}, w^{(n)})$ exists in $[0, t]$ for sufficiently large n. This implies $t < \liminf_{n\to\infty} T_{\mathrm{sur}}^{(n)}$.

Now assume that $T_{\mathrm{sur}} < \infty$. If

$$\lim_{t\uparrow T_{\mathrm{sur}}} (\|v\|_{C_t\mathcal{C}^{-\frac{2}{3}+\kappa'}} + \|w\|_{C_t\mathcal{C}^{-\frac{1}{2}-2\kappa}}) < \infty,$$

we can start from $(v, w)(T_{\mathrm{sur}} - \delta) \in \mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa}$ for small $\delta > 0$ and construct a solution on $[0, T_*]$, where T_* is uniform over δ. This implies that for sufficiently small $\delta > 0$, we can construct a solution on $[T_{\mathrm{sur}} - \frac{\delta}{2}, T_{\mathrm{sur}} + \frac{\delta}{2}]$ without explosion at the starting time. This is a contradiction, so we obtain the existence and uniqueness up to survival time with respect to the weaker norms. □

4.5 Renormalized equation

In this subsection, we show that a solution in the sense of Theorem 4.26 to the equation with a driving vector constructed from a driving force $\xi \in C_T\mathcal{C}^\beta$ for $\beta > -2$ and renormalization constants solves the renormalized equation.

We fix complex constants $\mathfrak{c}_1$, $\mathfrak{c}_{2,1}$ and $\mathfrak{c}_{2,2}$ and define functions X^τ as in Table 1 for every graphical symbols τ and construct the driving vector $X = (X^{\tau_1}, \dots, X^{\tau_{\#\mathcal{T}}})$. The Y/N in the Driver column in Table 1 indicates whether the term X^τ is included in the definition of a driving vector or not. The term X^τ with Driver column N is going to be used to define other terms. For the definition of $I(*, \bullet)$, see (4.2). Note that we can interpret the product in Table 1 in the usual sense because X^{τ_1} is a $\mathbf{C}$-valued continuous function by the assumption $\xi \in C_T\mathcal{C}^\beta$ for $\beta > -2$. The number in Regularity column denotes the exponent α_τ of the Hölder-Besov space $\mathcal{C}^{\alpha_\tau}$ where the term X^τ lives in. Precisely, α_τ means $\alpha_\tau - \kappa$ for any $\kappa > 0$ small enough.

Table 1: Definition of a driving vectors

Driver	Symbol	Definition	Regularity α_τ
Y	$X^{\tau_1}(= Z)$	$I(X_0^{\tau_1}, \xi)$	$-1/2$
N	X^{τ_2}	$\overline{X^{\tau_1}}$	$-1/2$
Y	X^{τ_3}	$(X^{\tau_1})^2$	-1
Y	X^{τ_4}	$X^{\tau_1}X^{\tau_2} - \mathfrak{c}_1$	-1
N	X^{τ_5}	$(X^{\tau_2})^2$	-1
N	X^{τ_6}	$X^{\tau_3}X^{\tau_2} - 2\mathfrak{c}_1 X^{\tau_1}$	$-3/2$
Y	X^{τ_7}	$I(X_0^{\tau_7}, X^{\tau_3})$	$+1$
Y	X^{τ_8}	$I(X_0^{\tau_8}, X^{\tau_4})$	$+1$
Y	$X^{\tau_9}(= W)$	$I(X_0^{\tau_9}, X^{\tau_6})$	$+1/2$
Y	$X^{\tau_{10}}$	$X^{\tau_9} \odot X^{\tau_1}$	0
Y	$X^{\tau_{11}}$	$X^{\tau_9} \odot X^{\tau_2}$	0
Y	$X^{\tau_{12}}$	$X^{\tau_7} \odot X^{\tau_4}$	0
Y	$X^{\tau_{13}}$	$X^{\tau_7} \odot X^{\tau_5} - 2\mathfrak{c}_{2,1}$	0
Y	$X^{\tau_{14}}$	$X^{\tau_8} \odot X^{\tau_4} - \mathfrak{c}_{2,2}$	0
Y	$X^{\tau_{15}}$	$X^{\tau_8} \odot X^{\tau_5}$	0
Y	$X^{\tau_{16}}$	$X^{\tau_9} \odot X^{\tau_4} - 2\mathfrak{c}_{2,2} X^{\tau_1}$	$-1/2$
Y	$X^{\tau_{17}}$	$X^{\tau_9} \odot X^{\tau_5} - 2\mathfrak{c}_{2,1} X^{\tau_2}$	$-1/2$

The next theorem is about the renormalized equation.

Theorem 4.29. *Let $0 < \kappa < \kappa' < 1/18$. Let $\xi \in C_T\mathcal{C}^\beta$ and $X_0^{\text{ı}}, X_0^{\Upsilon}, X_0^{\Upsilon'}, X_0^{\Upsilon''} \in \mathcal{C}^{\beta+2}$ for $\beta > -2$. Construct $X \in \mathcal{X}_1^\kappa$ as in Table 1. Let $(v, w) \in \mathcal{D}_T^{\kappa,\kappa'}$ be the solution to (4.9) with the initial condition $(v_0, w_0) \in \mathcal{C}^{-\frac{2}{3}+\kappa'} \times \mathcal{C}^{-\frac{1}{2}-2\kappa}$ for the driving vector X. Set $u = Z - \nu W + v + w$ and $\mathfrak{c} = 2(\mathfrak{c}_1 - \overline{\nu\mathfrak{c}_{2,1}} - 2\nu\mathfrak{c}_{2,2})$. Then u solves*

$$\partial_t u = (\mathsf{i} + \mu)\triangle u + \nu(1 - |u|^2)u + \nu\mathfrak{c}u + \xi, \qquad t > 0, \quad x \in \mathbf{T}^3.$$

with the initial condition $u_0 = Z_0 - \nu W_0 + v_0 + w_0$ in the usual mild sense.

The next lemma plays a key role to prove Theorem 4.29.

Lemma 4.30. *Let (v, w) be the solution to (4.9). Set $u_2 = v + w$. Then, we have*

$$\begin{aligned}
(4.16)\quad & F(v,w) + G(v,w) \\
&= -\nu\big\{(-\nu W + u_2)^2(\overline{Z} - \overline{\nu}\overline{W} + \overline{u_2}) + 2(-\nu W + u_2)(-\overline{\nu}\overline{W} + \overline{u_2})Z \\
&\qquad + 2(-\nu W + u_2)X^{\text{ı}} + (-\overline{\nu}\overline{W} + \overline{u_2})X^{\Upsilon} \\
&\qquad + 2(\overline{\nu\mathfrak{c}_{2,1}} + 2\nu\mathfrak{c}_{2,2})(Z - \nu W + u_2)\big\} \\
&\quad + (\nu + 1)(Z - \nu W + u_2).
\end{aligned}$$

Proof. It follows from the definition that

$$(4.17)\quad G_1(v,w) + G_2(v,w) = -\nu\big\{u_2^2(\overline{Z} - \overline{\nu}\overline{W}) + \overline{u_2}(u_2^2 + 2u_2(Z - \nu W))\big\}.$$

We will show

$$\begin{aligned}
(4.18)\quad & G_3(v,w) + G_4(v,w) + G_5(v,w) \\
&= -\nu\big\{u_2(2\nu\overline{\nu}W\overline{W} - 2\nu W\overline{Z} - 2\overline{\nu}\overline{W}Z) + \overline{u_2}(\nu^2W^2 - 2\nu WZ) \\
&\qquad - \nu^2\overline{\nu}W^2\overline{W} + \nu^2W^2\overline{Z} + 2\nu\overline{\nu}W\overline{W}Z \\
&\qquad - 4\nu((-\nu W + u_2) \circledcirc X^{\Upsilon'}) \odot X^{\text{ı}} \\
&\qquad - 2\overline{\nu}((-\overline{\nu}\overline{W} + \overline{u_2}) \circledcirc \overline{X^{\Upsilon'}}) \odot X^{\Upsilon} \\
&\qquad - 2\nu((-\overline{\nu}\overline{W} + \overline{u_2}) \circledcirc X^{\Upsilon}) \odot X^{\text{ı}} \\
&\qquad - \overline{\nu}((-\nu W + u_2) \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} \\
&\qquad - 2\nu W(\odot + \circledcirc)X^{\text{ı}} - \overline{\nu}\overline{W}(\odot + \circledcirc)X^{\Upsilon} \\
&\qquad + 2(\overline{\nu\mathfrak{c}_{2,1}} + 2\nu\mathfrak{c}_{2,2})(Z - \nu W + u_2)\big\}
\end{aligned}$$

$$
\begin{aligned}
&\quad + (\nu+1)(Z-\nu W+u_2),\\
(4.19)\qquad &G_6(v,w)+G_7(v,w)+G_8(v,w)\\
&\quad = -\nu\big\{2u_2(\odot+\circledcirc)X^{\blacktriangledown\blacktriangledown}+\overline{u_2}(\odot+\circledcirc)X^{\blacktriangledown}\\
&\qquad\quad + 4\nu((-\nu W+u_2)\circledcirc X^{\blacktriangledown\blacktriangledown})\odot X^{\blacktriangledown\blacktriangledown}\\
&\qquad\quad + 2\overline{\nu}((-\overline{\nu}\overline{W}+\overline{u_2})\circledcirc \overline{X^{\blacktriangledown\blacktriangledown}})\odot X^{\blacktriangledown}\\
&\qquad\quad + 2\nu((-\overline{\nu}\overline{W}+\overline{u_2})\circledcirc X^{\blacktriangledown})\odot X^{\blacktriangledown\blacktriangledown}\\
&\qquad\quad + \overline{\nu}((-\nu W+u_2)\circledcirc \overline{X^{\blacktriangledown}})\odot X^{\blacktriangledown}\big\}.
\end{aligned}
$$

Summing them up, we obtain

$$
\begin{aligned}
&G_1(v,w)+\cdots+G_8(v,w)\\
&\quad = -\nu\big\{\big\{u_2^2+2u_2(Z-\nu W)+\nu^2W^2-2\nu WZ\big\}\overline{u_2}\\
&\qquad + \big\{u_2^2-2\nu Wu_2+2Zu_2+\nu^2W^2-2\nu WZ\big\}(-\overline{\nu}\overline{W})\\
&\qquad + \big\{u_2^2-2\nu Wu_2+\nu^2W^2\big\}\overline{Z}\\
&\qquad + 2(-\nu W+u_2)(\odot+\circledcirc)X^{\blacktriangledown\blacktriangledown}+(-\overline{\nu}\overline{W}+\overline{u_2})(\odot+\circledcirc)X^{\blacktriangledown}\\
&\qquad + 2(\overline{\nu\mathfrak{c}_{2,1}}+2\nu\mathfrak{c}_{2,2})(Z-\nu W+u_2)\big\}\\
&\quad\quad + (\nu+1)(Z-\nu W+u_2)+cv-w\\
&\quad = -\nu\big\{(-\nu W+u_2)^2(\overline{Z}-\overline{\nu}\overline{W}+\overline{u_2})+2(-\nu W+u_2)(-\overline{\nu}\overline{W}+\overline{u_2})Z\\
&\qquad + 2(-\nu W+u_2)(\odot+\circledcirc)X^{\blacktriangledown\blacktriangledown}+(-\overline{\nu}\overline{W}+\overline{u_2})(\odot+\circledcirc)X^{\blacktriangledown}\\
&\qquad + 2(\overline{\nu\mathfrak{c}_{2,1}}+2\nu\mathfrak{c}_{2,2})(Z-\nu W+u_2)\big\}\\
&\quad\quad + (\nu+1)(Z-\nu W+u_2),
\end{aligned}
$$

which implies the conclusion.

For the rest of this proof, we prove (4.18) and (4.19). To show (4.18), we use the definition of $X^{\blacktriangledown\blacktriangledown}$ and Proposition 4.5. From them, we see

$$
\begin{aligned}
&WX^{\blacktriangledown\blacktriangledown}+4\nu R(W,X^{\blacktriangledown\blacktriangledown},X^{\blacktriangledown})\\
&\qquad = W\{(X^{\blacktriangledown\blacktriangledown}\odot X^{\blacktriangledown})-\mathfrak{c}_{2,2}\}+(W\circledcirc X^{\blacktriangledown\blacktriangledown})\odot X^{\blacktriangledown}-W(X^{\blacktriangledown\blacktriangledown}\odot X^{\blacktriangledown})\\
&\qquad = (W\circledcirc X^{\blacktriangledown\blacktriangledown})\odot X^{\blacktriangledown}-\mathfrak{c}_{2,2}W
\end{aligned}
$$

A similar argument implies

$$
\begin{aligned}
&\overline{W}\overline{X^{\blacktriangledown\blacktriangledown}}+R(\overline{W},\overline{X^{\blacktriangledown\blacktriangledown}},X^{\blacktriangledown})=(\overline{W}\circledcirc\overline{X^{\blacktriangledown\blacktriangledown}})\odot X^{\blacktriangledown}\\
&\overline{W}X^{\blacktriangledown\blacktriangledown}+R(\overline{W},X^{\blacktriangledown\blacktriangledown},X^{\blacktriangledown})=(\overline{W}\circledcirc X^{\blacktriangledown\blacktriangledown})\odot X^{\blacktriangledown}
\end{aligned}
$$

$$W\overline{X^{\Upsilon}} + R(W, \overline{X^{\Upsilon}}, X^{\Upsilon}) = (W \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} - 2\overline{\mathfrak{c}_{2,1}}W.$$

Applying these identities and the definitions of X^{Ψ} and X^{Ψ}, we obtain

$$\begin{aligned}
G_4(v,w) = -\nu\big\{ & -\nu^2\overline{\nu}W^2\overline{W} + \nu^2W^2\overline{Z} + 2\nu\overline{\nu}W\overline{W}Z \\
& + 4\nu^2\{(W \circledcirc X^{\Upsilon}) \odot X^{\Psi} - \mathfrak{c}_{2,2}W\} + 2\overline{\nu}^2(\overline{W} \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} \\
& + 2\nu\overline{\nu}(\overline{W} \circledcirc X^{\Upsilon}) \odot X^{\Psi} + \nu\overline{\nu}\{(W \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} - 2\overline{\mathfrak{c}_{2,1}}W\} \\
& - 2\nu(W \odot X^{\Psi} - 2\mathfrak{c}_{2,2}X^{\mathsf{I}}) - 2\nu W \circledcirc X^{\Psi} \\
& - \overline{\nu}(\overline{W \odot X^{\Psi}} - 2\overline{\mathfrak{c}_{2,1}X^{\Psi}}) - \overline{\nu}\overline{W} \circledcirc X^{\Upsilon}\big\} \\
& + (\nu+1)(Z - \nu W) \\
= -\nu\big\{ & -\nu^2\overline{\nu}W^2\overline{W} + \nu^2W^2\overline{Z} + 2\nu\overline{\nu}W\overline{W}Z \\
& + 4\nu(\nu W \circledcirc X^{\Upsilon}) \odot X^{\Psi} + 2\overline{\nu}(\overline{\nu}\overline{W} \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} \\
& + 2\nu(\overline{\nu}\overline{W} \circledcirc X^{\Upsilon}) \odot X^{\Psi} + \overline{\nu}(\nu W \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} \\
& - 2\nu W(\odot + \circledcirc)X^{\Psi} - \overline{\nu}\overline{W}(\odot + \circledcirc)X^{\Upsilon} \\
& + 2(\overline{\nu}\overline{\mathfrak{c}_{2,1}} + 2\nu\mathfrak{c}_{2,2})(Z - \nu W)\big\} \\
& + (\nu+1)(Z - \nu W).
\end{aligned}$$

We use the similar argument to obtain

$$\begin{aligned}
& G_3(v,w) + G_5(v,w) \\
& \quad = -\nu\big\{u_2(2\nu\overline{\nu}W\overline{W} - 2\nu W\overline{Z} - 2\overline{\nu}\overline{W}Z) + \overline{u_2}(\nu^2W^2 - 2\nu WZ) \\
& \qquad - 4\nu(u_2 \circledcirc X^{\Upsilon}) \odot X^{\Psi} - 2\overline{\nu}(\overline{u_2} \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} \\
& \qquad - 2\nu(\overline{u_2} \circledcirc X^{\Upsilon}) \odot X^{\Psi} - \overline{\nu}(u_2 \circledcirc \overline{X^{\Upsilon}}) \odot X^{\Upsilon} \\
& \qquad + 2(\overline{\nu}\overline{\mathfrak{c}_{2,1}} + 2\nu\mathfrak{c}_{2,2})u_2\big\} \\
& \quad + (\nu+1)u_2.
\end{aligned}$$

Combining them, we see (4.18). From the definition of $\mathrm{com}(v,w)$, we obtain (4.19). The proof is completed. □

Proof of Theorem 4.29. Set $u_2 = v + w$, $u_1 = -\nu W + u_2$ and $u = Z + u_1$. Note that u_2 solves $\mathcal{L}^1 u_2 = F(v,w) + G(v,w)$. Substituting $X^{\Psi} = Z\overline{Z} - \mathfrak{c}$ and $X^{\Upsilon} = Z^2$ to (4.16), we have

$$\begin{aligned}
F(v,w) + G(v,w) = & -\nu(Z+u_1)^2(\overline{Z} + \overline{u_1}) + \nu(Z+u_1) + 2\nu\mathfrak{c}(Z+u_1) \\
& + \nu(Z^2\overline{Z} - 2\mathfrak{c}_1 Z) + Z + u_1
\end{aligned}$$

$$= -\nu u^2\overline{u} + \nu u + 2\nu \mathfrak{c} u + \nu(Z^2\overline{Z} - 2\mathfrak{c}_1 Z) + u$$

where $\mathfrak{c} = \mathfrak{c}_1 - \overline{\nu\mathfrak{c}_{2,1}} + 2\nu\mathfrak{c}_{2,2}$. Hence

$$\begin{aligned}\{\partial_t - (\mathrm{i}+\mu)\triangle\}u &= \mathcal{L}^1 u - u \\ &= \mathcal{L}^1(Z - \nu W + u_2) - u \\ &= \xi - \nu(Z^2\overline{Z} - 2\mathfrak{c}_1 Z) + \{F(v,w) + G(v,w)\} - u \\ &= -\nu u^2\overline{u} + \nu u + 2\nu\mathfrak{c} u + \xi.\end{aligned}$$

The proof is completed. □

5 Proof of convergence of driving vectors

This section is a probabilistic part of proof of Theorem 4.1. In this section, we construct a driving vector $X \in \mathcal{X}_T^\kappa$ associated to the white noise ξ (Theorem 5.9). After that we derive the expression of renormalization constants $\mathfrak{c}_1^\epsilon$, $\mathfrak{c}_{2,1}^\epsilon$ and $\mathfrak{c}_{2,2}^\epsilon$ used in the construction of X (Proposition 5.20) and obtain the divergence rate of them (Proposition 5.21).

First of all, we define Ornstein-Uhlenbeck like process $Z = Z(t,x)$, which is is a seed of the driving vector. The process Z is defined as a stationary solution to the following equation:

$$\partial_t Z = \{(\mathrm{i}+\mu)\triangle - 1\}Z + \xi.$$

The solution has a formal expression

$$Z_t = I(\xi)_t = \int_{-\infty}^t P_{t-s}^1 \xi_s \, ds = \sum_{k\in\mathbf{Z}^3} \left(\int_{-\infty}^t P_{t-s}^1 \mathbf{e}_k \hat{\xi}_s(k) \, ds \right).$$

Here, I is defined by (4.3). Since Z is a distribution-valued process, we cannot define processes such as Z^2 and $Z^2\overline{Z}$ a priori. To define such processes, we consider an approximation $\{Z^\epsilon\}_{0<\epsilon<1}$ of Z and define Z^2 and $Z^2\overline{Z}$ as renormalized limits of $(Z^\epsilon)^2$ and $Z^2\overline{Z}$ in an appreciate topology, respectively. To this end, we recall the smeared noise $\{\xi^\epsilon\}_{0<\epsilon<1}$ defined by (4.1) approximates the white noise ξ. Using the approximation, we define

$$Z_t^\epsilon = \int_{-\infty}^t P_{t-s}^1 \xi_s^\epsilon \, ds = \sum_{k\in\mathbf{Z}^3} \chi^\epsilon(k) \left(\int_{-\infty}^t P_{t-s}^1 \mathbf{e}_k \hat{\xi}_s(k) \, ds \right). \tag{5.1}$$

We recall that the Fourier transform $\{\hat{\xi}(k)\}_{k\in\mathbf{Z}^3}$ of ξ has the same law of the white noise associated to $(E, \mathcal{B}, d\mathfrak{m})$. Here, $E = \mathbf{R}\times\mathbf{Z}^3$, $\mathcal{B}$ is the product σ-field

of $\mathcal{B}(\mathbf{R})$ and $2^{\mathbf{Z}^3}$ and $d\mathfrak{m} = dsdk$, where ds and dk are the Lebesgue measure on $\mathbf{R}$ and the counting measure $\mathbf{Z}^3$, respectively. Note that $d\mathfrak{m}$ is given by

$$\mathfrak{m}(A) = \int_E \mathtt{1}_A(s,k)\, dsdk = \sum_{k\in\mathbf{Z}^3} \int_{\mathbf{R}} \mathtt{1}_A(s,k)\, ds.$$

We denote by $\mathcal{B}^*$ the set of all elements $A \in \mathcal{B}$ such that $\mathfrak{m}(A) < \infty$. Let $M(A) = \sum_{k\in\mathbf{Z}^3} \int_{\mathbf{R}} \mathtt{1}_A(s,k)\hat{\xi}_s(k)\, ds$ for $A \in \mathcal{B}^*$. Since $\{M(A); A \in \mathcal{B}^*\}$ is a jointly isotropic complex normal such that $\boldsymbol{E}[M(A)\overline{M(B)}] = \mathfrak{m}(A \cap B)$, we can define complex multiple Itô-Wiener integrals $\mathcal{J}_{p,q}$ to calculate $(Z^\epsilon)^2$ and $(Z^\epsilon)^2\overline{Z^\epsilon}$; see Section A. By using them, we show their convergence after renormalization and construct the driving vector X.

Throughout this section, we use the notations in Section A and the following:

- We use $m = (s,k)$, $n = (t,l)$, $\mu = (\sigma,k)$ and $\nu = (\tau,l)$ to denote a generic element in E.
- For $\nu_i = (\tau_i, l_i)$, we write $\nu_{-i} = (\tau_i, -l_i)$.
- For $p_1,\dots,p_n \in \mathbf{Z}\backslash\{0\}$, we write $k_{p_1,\dots,p_n} = (k_{p_1},\dots,k_{p_n})$ and $k_{[p_1,\dots,p_n]} = k_{p_1} + \cdots + k_{p_n}$ for shorthand. We use the same abbreviation for s, t, l, m, n, σ, τ, μ and ν.
- We define $|k|_* = 1 + |k| = 1 + \sqrt{k_1^2 + k_2^2 + k_3^2}$ for $k = (k_1,k_2,k_3) \in \mathbf{Z}^3$ and $|m|_* = |(s,k)|_* = 1 + |s|^{1/2} + |k|$. The same notations are used for l, n, μ and ν.

Let $f : E^{p+q} \to \mathbf{C}$ satisfy

$$\int_{\mathbf{R}^{p+q}} |f((s,k)_{1,\dots,p},(t,l)_{1,\dots,q})|^2\, ds_1\cdots ds_p dt_1\cdots dt_q < \infty$$

for every $k_1,\dots,k_p,l_1,\dots,l_q \in \mathbf{Z}^3$. For such f, we can define the Fourier transform $\mathcal{F}_{\mathrm{time}}f$ with respect to time parameters. In particular, if f is integrable and square-integrable with respect to the time parameters, then $\mathcal{F}_{\mathrm{time}}f$ is given by

$$\begin{aligned}[\mathcal{F}_{\mathrm{time}}f]((\sigma,k)_{1,\dots,p},(\tau,l)_{1,\dots,q}) = \int_{\mathbf{R}^{p+q}} & ds_1\cdots ds_p dt_1\cdots dt_q \\ & \times e^{-2\pi\mathrm{i}(\sigma_1 s_1+\cdots+\sigma_p s_p+\tau_1 t_1+\cdots+\tau_q t_q)} f((s,k)_{1,\dots,p},(t,l)_{1,\dots,q}).\end{aligned}$$

5.1 Convergence criteria

In this subsection, we establish convergence criteria of Itô-Wiener integrals.

5.1.1 $\mathcal{C}^\alpha$-valued random variables

We want to define a random field of the form

$$X(x) = \mathcal{J}_{p,q}(f(x))$$

for a kernel $f \in C(\mathbf{T}^3, L^\infty_{p,q})$ even if $f(x) \notin L^2_{p,q}$. Here $L^\infty_{p,q}$ is the space of the essentially bounded measurable functions defined on E^{p+q}. Assume now that $\langle f, \phi\rangle = \int_{\mathbf{T}^3} f(x)\phi(x)\,dx \in L^2_{p,q}$ for every $\phi \in \mathcal{D}$ and define the family of random variables

$$X(\phi) = \mathcal{J}_{p,q}(\langle f, \phi\rangle).$$

If there exists a $\mathcal{D}'$-valued random variable $\tilde{X}$ such that

$$\langle \tilde{X}, \phi\rangle = X(\phi),$$

then we write $\tilde{X}(x) = \mathcal{J}_{p,q}(f(x))$.

Now we define $X_j(x) = X((\mathcal{F}^{-1}\rho_j)(x - \cdot))$. If $X = \sum_{j\geq -1} X_j$ converges in $\mathcal{D}'$, it satisfies $\langle X, \phi\rangle = X(\phi)$ for every $\phi \in \mathcal{D}$, so we can write $X(x) = \mathcal{J}_{p,q}(f(x))$.

Proposition 5.1. *Let $\alpha \in \mathbf{R}$ and $p \in (1, \infty)$. If*

$$C_{\alpha,p} = \sum_{j\geq -1} 2^{(2\alpha p+1)j} \left(\sup_{x\in\mathbf{T}^3} \boldsymbol{E}[|X_j(x)|^2]\right)^p < \infty,$$

then $X = \sum_{j\geq -1} X_j$ converges in $L^{2p}(\Omega, \mathcal{C}^\alpha)$ and we have

$$\boldsymbol{E}[\|X\|^{2p}_{\mathcal{C}^\alpha}] \lesssim C_{\alpha,p}.$$

Proof. Since $\langle f, (\mathcal{F}^{-1}\rho_j)(x - \cdot)\rangle = \triangle_j f(x)$, we have $\mathcal{F}X_j(k) = \mathcal{J}_{p,q}(\rho_j \mathcal{F}f(k))$, which implies that the support of $\mathcal{F}X_j(k)$ contained in an annulus. Hence we can apply [BCD11, Lemma 2.69] to X. By a similar argument as [Hos16b, Lemma 5.3], we see the assertion. □

5.1.2 Good kernels

We consider a random field of the form

$$X(t, x) = \mathcal{J}_{p,q}(f_{(t,x)})$$

for a kernel $f(t, \cdot) \in C(\mathbf{T}^3, L^\infty_{p,q})$ which satisfies the conditions as above for each fixed t. We are interested in the case that f satisfies the following good conditions.

Definition 5.2. We say that a family $\{f_{(t,x)}\}_{t\geq 0, x\in \mathbf{T}^3}$ is good if it has the form

$$f_{(t,x)}(m_{1,\dots,p}, n_{1,\dots,q}) = \mathbf{e}_{k_{[1\dots p]}-l_{[1\dots q]}}(x) H_t(m_{1,\dots,p}, n_{1,\dots,q})$$

for some $H_t \in L^\infty_{p,q}$ which is in L^2 with respect to $(s_{1,\dots,p}, t_{1,\dots,q})$ for each fixed $(k_{1,\dots,p}, l_{1,\dots,q})$ and $Q_t = \mathcal{F}_{\mathrm{time}} H_t$ satisfies

$$Q_t(\mu_{1,\dots,p}, \nu_{1,\dots,q}) = e^{-2\pi \mathrm{i}(\sigma_{[1\dots p]}+\tau_{[1\dots q]})t} Q_0(\mu_{1,\dots,p}, \nu_{1,\dots,q}).$$

For a function $f : E^{p+q} \to \mathbf{C}$, we set

$$\tilde{\rho_j} f(m_{1,\dots,p}, n_{1,\dots,q}) = \rho_j(k_{[1\dots p]} - l_{[1\dots q]}) f(m_{1,\dots,p}, n_{1,\dots,q}).$$

We define

$$R(\sigma_{1,\dots,p}, \tau_{1,\dots,q}) = \sigma_{[1\dots p]} + \tau_{[1\dots q]}.$$

In order to estimate Besov norm of X, is is enough to estimate Q_0.

Proposition 5.3. *Let $\{f_{(t,x)}\}_{t\geq 0, x\in \mathbf{T}^3}$ be a good kernel. Assume that there exist $\beta \in \mathbf{R}$, $\theta_0 \in (0,1)$ and $C > 0$ such that*

$$\| |R|^{\frac{\theta}{2}} \tilde{\rho_j} Q_0 \|_{L^2_{p,q}} \leq C 2^{(\beta+\theta)j}$$

for every $j \geq -1$ and $\theta \in [0, \theta_0]$. Then we have

$$\boldsymbol{E}[\|X\|^{2p}_{C^\kappa_T \mathcal{C}^{\alpha-2\kappa}}] \lesssim C^{2p},$$

for every $p \in (1,\infty)$, $\alpha < -\beta$ and $\kappa \in [0, \frac{\theta_0}{2})$. Here $C^0_T \mathcal{C}^\alpha = C_T \mathcal{C}^\alpha$.

Proof. Let $1 < p < \infty$ satisfy $2(\alpha+\beta)p + 1 < 0$. For every $t \in [0,\infty)$, we have $X_t \in C_T \mathcal{C}^\alpha$ from

$$\boldsymbol{E}[\|X_t\|^{2p}_{\mathcal{C}^\alpha}] < \infty.$$

We will show this inequality. Set $X_j(t,x) = \mathcal{J}_{p,q}(\langle f_{(t,\cdot)}, (\mathcal{F}^{-1}\rho_j)(x-\cdot)\rangle)$. Since $\langle f(t,\cdot), (\mathcal{F}^{-1}\rho_j)(x-\cdot)\rangle = [\triangle_j f_{(t,\cdot)}](x) = \tilde{\rho_j} f_{(t,x)}$, we have

$$\begin{aligned}\boldsymbol{E}[|X_j(t,x)|^2] &= \|\tilde{\rho_j} f_{(t,x)}\|^2_{L^2_{p,q}} \\ &= \|\mathcal{F}_{\mathrm{time}} \tilde{\rho_j} f_{(t,x)}\|^2_{L^2_{p,q}} = \|\tilde{\rho_j} e^{-2\pi \mathrm{i} R t} Q_0\|^2_{L^2_{p,q}} = \|\tilde{\rho_j} Q_0\|^2_{L^2_{p,q}}.\end{aligned}$$

Using the assumption with $\theta = 0$, we obtain $\boldsymbol{E}[|X_j(t,x)|^2] \leq (C 2^{\beta j})^2$. Hence

$$C_{\alpha,p} = \sum_{j=-1}^{\infty} 2^{(2\alpha p+1)j} (C 2^{\beta j})^{2p} \leq C^{2p} \sum_{j=-1}^{\infty} 2^{(2(\alpha+\beta)p+1)j} < \infty.$$

From Proposition 5.1, we see the inequality.

Set $\alpha' = \alpha - 2\kappa$ and take $0 < \theta < \theta_0$ such that $\alpha' + \beta + \theta < 0$. For any $1 < p < \infty$ such that $2(\alpha' + \beta + \theta)p + 1 < 0$ and $p\theta > 1$, we can show that

$$\boldsymbol{E}[\|X_t - X_s\|^{2p}_{\mathcal{C}^{\alpha'}}] \le C|t-s|^{p\theta},$$

where C is a positive constant independent of s, t. This estimate and the Kolmogorov continuity theorem implies $X \in C^{\kappa}_T\mathcal{C}^{\alpha-2\kappa}$. □

For a function $f : E^{p+q} \to \mathbf{C}$ and $\mu = (\sigma, k)$, we write

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} f(\mu_{1,\dots,p}, \nu_{1,\dots,q})$$

for the integration over the "hyperplane" $\{\mu_{[1\dots p]} + \nu_{[(-1)\dots(-q)]} = \mu\}$.

Proposition 5.4. *Let $\{f_{(t,x)}\}_{t\ge 0, x\in\mathbf{T}^3}$ be a good kernel. Assume that there exist $\gamma > 1$, $\delta \ge 0$ and $C > 0$ such that*

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0(\mu_{1,\dots,p}, \nu_{1,\dots,q})|^2 \le C|\mu|_*^{-2\gamma}|k|_*^{-2\delta}. \tag{5.2}$$

Then we have

$$\||R|^{\theta/2}\tilde{\rho}_j Q_0\|_{L^2_{p,q}} \lesssim C2^{(\frac{5}{2}-\gamma-\delta+\theta)j}$$

for every $\theta \in [0, \gamma - 1)$. As a consequence, we have

$$\boldsymbol{E}[\|X\|^{2p}_{C^{\kappa}_T\mathcal{C}^{\alpha-2\kappa}}] \lesssim C^{2p},$$

for every $p \in (1, \infty)$, $\alpha < -\frac{5}{2} + \gamma + \delta$ and $\kappa \in [0, \frac{\gamma-1}{2})$.

Proof. Since

$$\begin{aligned}
&\||R|^{\theta/2}\tilde{\rho}_j Q_0\|^2_{L^2_{p,q}} \\
&\quad= \sum_{k\in\mathbf{Z}^3} \rho_j(k)^2 \int_{\mathbf{R}} |\sigma|^{\theta} \left(\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0(\mu_{1,\dots,p}, \nu_{1,\dots,q})|^2 \right) d\sigma \\
&\quad\le C \sum_{k\in\mathbf{Z}^3} \rho_j(k)^2 |k|_*^{-2\delta} \int_{\mathbf{R}} |\mu|_*^{-2(\gamma-\theta)}\, d\sigma
\end{aligned}$$

if $\gamma - \theta > 1$, we have

$$\begin{aligned}
\||R|^{\theta/2}\tilde{\rho}_j Q_0\|^2_{L^2_{p,q}} &\lesssim C \sum_{k\in\mathbf{Z}^3} \rho_j(k)^2 |k|_*^{-2\delta} |k|_*^{2(1-\gamma+\theta)} \\
&\lesssim C(2^j)^3 (2^j)^{2(1-\gamma+\theta-\delta)} \\
&= C2^{(5-2\gamma-2\delta+2\theta)j}.
\end{aligned}$$

The proof is completed. □

5.2 Definitions of driving vectors

Since Z is a distribution-valued process, we cannot define a process such as Z^2, $Z\overline{Z}$ and $Z^2\overline{Z}$ a priori. To define such processes, we consider an approximation $\{Z^\epsilon\}_{0<\epsilon<1}$ of Z and define Z^2, $Z\overline{Z}$ and $Z^2\overline{Z}$ as renormalized limits of $(Z^\epsilon)^2$, $Z^\epsilon\overline{Z^\epsilon}$ and $(Z^\epsilon)^2\overline{Z^\epsilon}$.

5.2.1 Ornstein-Uhlenbeck like process and its approximations

We give an expression of Z^ϵ defined by (5.1) in terms of Itô-Wiener integral. Since we have $P_s^1\mathbf{e}_k = h(s,k)\mathbf{e}_k$, where $h(s,k) = e^{-\{4\pi^2(\mathsf{i}+\mu)|k|^2+1\}s}$, we see

$$Z_t^\epsilon = \sum_{k\in\mathbf{Z}^3} \chi^\epsilon(k) \int_{-\infty}^t h(t-s,k)\mathbf{e}_k \hat{\xi}_s(k)\, ds.$$

Hence, we can write $Z^\epsilon_{(t,x)} = \mathcal{J}_{1,0}(f^\epsilon_{(t,x)})$ with

$$f^\epsilon_{(t,x)}(s,k) = \mathbf{e}_k(x)H_t^\epsilon(s,k), \quad H_t^\epsilon(s,k) = \chi^\epsilon(k)H_t(s,k),$$
$$H_t(s,k) = \mathbf{1}_{[0,\infty)}(t-s)h(t-s,k).$$

Note that $Q_t = \mathcal{F}_{\mathrm{time}}H_t$ is given by

$$Q_t(\sigma,k) = \frac{e^{-2\pi\mathsf{i}\sigma t}}{-2\pi\mathsf{i}\sigma + 4\pi^2(\mathsf{i}+\mu)|k|^2+1}.$$

In particular, we see $Q_t(\mu) = e^{-2\pi\mathsf{i}t}Q_0(\mu)$. We simply write $Q_0 = Q$.

5.2.2 Definition of driving vectors

For every $0<\epsilon<1$ and graphical symbols τ, we define distributions $X^{\epsilon,\tau}$ as in Table 2. The operator I is defined by (4.3) and the constants $\mathfrak{c}_1^\epsilon$, $\mathfrak{c}_{2,1}^\epsilon$ and $\mathfrak{c}_{2,2}^\epsilon$ in Table 2 are defined by

(5.3)
$$\mathfrak{c}_1^\epsilon = \boldsymbol{E}[X^{\epsilon,\mathbin{|}}_{(t,x)}X^{\epsilon,\vdots}_{(t,x)}], \quad \mathfrak{c}_{2,1}^\epsilon = \frac{1}{2}\boldsymbol{E}[X^{\epsilon,\curlyvee}_{(t,x)} \odot X^{\epsilon,\vdots}_{(t,x)}], \quad \mathfrak{c}_{2,2}^\epsilon = \boldsymbol{E}[X^{\epsilon,\curlyvee}_{(t,x)} \odot X^{\epsilon,\curlyvee}_{(t,x)}].$$

The other symbols and regularities have the same meanings as in Table 1. We set

$$X^\epsilon = (X^{\epsilon,\mathbin{|}}, X^{\epsilon,\mathsf{V}}, X^{\epsilon,\mathsf{V}:}, X^{\epsilon,\curlyvee}, X^{\epsilon,\curlyvee:}, X^{\epsilon,\mathsf{V}:\mathsf{V}:},$$
$$X^{\epsilon,\mathsf{V}:\curlyvee}, X^{\epsilon,\curlyvee:\mathsf{V}:}, X^{\epsilon,\curlyvee:\curlyvee}, X^{\epsilon,\mathsf{V}:\mathsf{V}:\mathsf{V}:}, X^{\epsilon,\mathsf{V}:\curlyvee:}, X^{\epsilon,\curlyvee:\mathsf{V}:\mathsf{V}:}, X^{\epsilon,\curlyvee:\curlyvee:}).$$

The constants $\mathfrak{c}_1^\epsilon$, $\mathfrak{c}_{2,1}^\epsilon$ and $\mathfrak{c}_{2,2}^\epsilon$ look dependent on (t,x) and the dyadic partition $\{\rho_m\}_{m=-1}^\infty$ of unity. However, we will show that they are not in Proposition 5.20.

Table 2: List of distributions

Driver	Distribution $X^{\epsilon,\tau}$	Definition	Regularity α_τ
Y	$X^{\epsilon,\bullet}$	Z^ϵ	$-1/2$
N	$X^{\epsilon,\bullet\bullet}$	$\overline{X^{\epsilon,\bullet}}$	$-1/2$
Y	$X^{\epsilon,\bullet\bullet\bullet}$	$(X^{\epsilon,\bullet})^2$	-1
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet}X^{\epsilon,\bullet\bullet} - \mathfrak{c}_1^\epsilon$	-1
N	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet}$	$(X^{\epsilon,\bullet\bullet})^2$	-1
N	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet}X^{\epsilon,\bullet\bullet} - 2\mathfrak{c}_1^\epsilon X^{\epsilon,\bullet}$	$-3/2$
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$I(X^{\epsilon,\bullet\bullet\bullet})$	$+1$
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$I(X^{\epsilon,\bullet\bullet\bullet\bullet})$	$+1$
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$I(X^{\epsilon,\bullet\bullet\bullet\bullet\bullet})$	$+1/2$
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet}$	0
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet}$	0
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet\bullet}$	0
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet\bullet\bullet} - 2\mathfrak{c}_{2,1}^\epsilon$	0
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet\bullet\bullet} - \mathfrak{c}_{2,2}^\epsilon$	0
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet\bullet\bullet\bullet}$	0
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet\bullet\bullet} - 2\mathfrak{c}_{2,2}^\epsilon X^{\epsilon,\bullet}$	$-1/2$
Y	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet}$	$X^{\epsilon,\bullet\bullet\bullet\bullet\bullet\bullet} \odot X^{\epsilon,\bullet\bullet\bullet\bullet\bullet} - 2\mathfrak{c}_{2,1}^\epsilon X^{\epsilon,\bullet\bullet}$	$-1/2$

5.2.3 Itô-Wiener integral expressions of driving vectors

We give expressions of $X^{\epsilon,\tau}$ by Itô-Wiener integrals.

We start to discuss with $\tau = \bullet, \bullet\bullet, \bullet\bullet\bullet, \bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet$. We denote by $p(\tau)$ and $q(\tau)$ the number of circles and squares in τ, respectively. We write

$$\chi^\epsilon(k_{1,\dots,p}, l_{1,\dots,q}) = \prod_{i=1}^{p} \chi^\epsilon(k_i) \prod_{j=1}^{q} \chi^\epsilon(l_j).$$

Proposition 5.5. *Let $\tau = \bullet, \bullet\bullet, \bullet\bullet\bullet, \bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet, \bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet\bullet$, $p = p(\tau)$ and $q = q(\tau)$. Then $X_{(t,x)}^{\epsilon,\tau} = \mathcal{J}_{p,q}(f_{(t,x)}^{\epsilon,\tau})$, where $f_{(t,x)}^{\epsilon,\tau} = f_{(t,x)}^{\epsilon,\tau}(m_{1,\dots,p}, n_{1,\dots,q})$ is a good kernel with functions $H_t^{\epsilon,\tau}$ and $Q_0^{\epsilon,\tau}$ defined as follows.*

(1) We have $H_t^{\epsilon,\tau}(m_{1,\dots,p}, n_{1,\dots,q}) = \chi^\epsilon(k_{1,\dots,p}, l_{1,\dots,q}) H_t^\tau(m_{1,\dots,p}, n_{1,\dots,q})$, where $\{H_t^\tau\}_{t\geq 0} \in L_{p,q}^2$ is given as follows.

- *$H_t^\bullet(m_1) = H_t(m_1)$ and $H_t^{\bullet\bullet}(n_1) = \overline{H_t(n_1)}$.*

- *For* $\tau = \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}$,

$$H_t^\tau(m_{1,\dots,p}, n_{1,\dots,q}) = \prod_{i=1}^{p} H_t^{\mathbb{I}}(m_i) \prod_{j=1}^{q} H_t^{\mathbb{I}}(n_j).$$

- *Let* $\tau_0 = \mathbb{V}, \mathbb{V}, \mathbb{V}$ *for* $\tau = \mathbb{Y}, \mathbb{Y}, \mathbb{Y}$, *respectively.*

$$H_t^\tau(m_{1,\dots,p}, n_{1,\dots,q}) = \int_{\mathbf{R}} H_t(u, k_{[1\dots p]} - l_{[1\dots q]}) H_u^{\tau_0}(m_{1,\dots,p}, n_{1,\dots,q})\, du.$$

(2) We have $Q_0^{\epsilon,\tau}(\mu_{1,\dots,p}, \nu_{1,\dots,q}) = \chi^\epsilon(k_{1,\dots,p}, l_{1,\dots,q}) Q_0^\tau(\mu_{1,\dots,p}, \nu_{1,\dots,q})$, *where* $Q_0^\tau \in L^2_{p,q}$ *is given as follows.*

- $Q_0^{\mathbb{I}}(\mu_1) = Q(\mu_1)$ *and* $Q_0^{\mathbb{I}}(\nu_1) = \overline{Q_0(-\nu_{-1})}$.
- *For* $\tau = \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}$,

$$Q_0^\tau(\mu_{1,\dots,p}, \nu_{1,\dots,q}) = \prod_{i=1}^{p} Q_0^{\mathbb{I}}(\mu_i) \prod_{j=1}^{q} Q_0^{\mathbb{I}}(\nu_j).$$

- *Let* $\tau_0 = \mathbb{V}, \mathbb{V}, \mathbb{V}$ *for* $\tau = \mathbb{Y}, \mathbb{Y}, \mathbb{Y}$, *respectively.*

$$Q_0^\tau(\mu_{1,\dots,p}, \nu_{1,\dots,q}) = Q(\mu_{[1\dots p]} + \nu_{[(-1)\dots(-q)]}) Q_0^{\tau_0}(\mu_{1,\dots,p}, \nu_{1,\dots,q}).$$

In the above, we regard H_t^τ *as a function with respect to* $n_{1,\dots,q}$ *and* $m_{1,\dots,p}$ *for* $p = 0$ *and* $q = 0$, *respectively. In particular,* H_t^τ *is a constant for* $p = q = 0$. *We use the same convention for* Q_0^τ.

Proof. The assertion follows from Proposition A.1. □

From this proposition, we can guess the limit X^τ of $\{X^{\epsilon,\tau}\}_{0<\epsilon<1}$ as follows:

Definition 5.6. Let $\tau \in \{\mathbb{I}, \mathbb{I}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{Y}, \mathbb{Y}, \mathbb{Y}\}$, $p = p(\tau)$ and $q = q(\tau)$. We define

$$f^\tau_{(t,x)}(m_{1,\dots,p}, n_{1,\dots,q}) = \mathbf{e}_{k_{[1\dots p]} - l_{[1\dots q]}} H_t^\tau(m_{1,\dots,p}, n_{1,\dots,q})$$

and $X^\tau_{(t,x)} = \mathcal{J}_{p,q}(f^\tau_{(t,x)})$.

Next, we consider $X^{\epsilon,\tau}$ for $\tau = \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}, \mathbb{V}$. For these τ, we define $(\tau_1, \tau_2) = (\mathbb{V}, \mathbb{I})$, $(\mathbb{V}, \mathbb{I})$, $(\mathbb{Y}, \mathbb{V})$, $(\mathbb{Y}, \mathbb{V})$, $(\mathbb{Y}, \mathbb{V})$, $(\mathbb{Y}, \mathbb{V})$, $(\mathbb{V}, \mathbb{V})$, $(\mathbb{V}, \mathbb{V})$, respectively. We simply write $p_i = p(\tau_i)$ and $q_i = q(\tau_i)$ for $i = 1, 2$. We set $p = p_1 + p_2$ and $q = q_1 + q_2$.

We define the function $\psi_\circ : \mathbf{Z}^3 \times \mathbf{Z}^3 \to \mathbf{R}$ by

$$\psi_\circ(k, l) = \sum_{|i-j| \le 1} \rho_i(k) \rho_j(l). \tag{5.4}$$

Proposition 5.7. *For above (τ_1, τ_2), it holds that*

$$X_t^{\epsilon,\tau_1} \odot X_t^{\epsilon,\tau_2}(x) = \sum_g \mathcal{J}_{p-\#g,q-\#g}(f_{(t,x)}^{\epsilon,(\tau_1,\tau_2,g)}),$$

where g runs over all of the graphs consisting of disjoint edges

$$(i,j) \in \{1,\dots,p_1\} \times \{q_1+1,\dots,q\} \cup \{p_1+1,\dots,p\} \times \{1,\dots,q_1\},$$

and $f_{(t,x)}^{\epsilon,(\tau_1,\tau_2,g)}$ is a good kernel with functions $H_t^{\epsilon,(\tau_1,\tau_2,g)}$ and $Q_0^{\epsilon,(\tau_1,\tau_2,g)}$ defined as follows.

(1) $H_t^{\epsilon,(\tau_1,\tau_2,\emptyset)}$ is given by

$$\begin{aligned} H_t^{\epsilon,(\tau_1,\tau_2,\emptyset)}(m_{1,\dots,p}, n_{1,\dots,q}) &= \psi_\circ(k_{[1\dots p_1]} - l_{[1\dots q_1]}, k_{[(p_1+1)\dots p]} - l_{[(q_1+1)\dots q]}) \\ &\quad \times H_t^{\epsilon,\tau_1}(m_{1,\dots,p_1}, n_{1,\dots,q_1}) H_t^{\epsilon,\tau_2}(m_{(p_1+1),\dots,p}, n_{(q_1+1),\dots,q}). \end{aligned}$$

For general g, $H_t^{\epsilon,(\tau_1,\tau_2,g)}$ is given by

$$H_t^{\epsilon,(\tau_1,\tau_2,g)}(m_{1,\dots,p}, n_{1,\dots,q} \setminus g) = \int_{E^{2\#g}} H_t^{\epsilon,(\tau_1,\tau_2,\emptyset)}(m_{1,\dots,p}, n_{1,\dots,q})\, d(m,n)_g,$$

where $(m_{1,\dots,p}, n_{1,\dots,q} \setminus g)$ means that variables (m_i, n_j) are removed for all $(i,j) \in g$ and $d(m,n)_g = \prod_{(i,j)\in g} \delta(m_i - n_j) dm_i dn_j$.

(2) $Q_0^{\epsilon,(\tau_1,\tau_2,\emptyset)}$ is given by

$$\begin{aligned} Q_0^{\epsilon,(\tau_1,\tau_2,\emptyset)}(\mu_{1,\dots,p}, \nu_{1,\dots,q}) &= \psi_\circ(k_{[1\dots p_1]} - l_{[1\dots q_1]}, k_{[(p_1+1)\dots p]} - l_{[(q_1+1)\dots q]}) \\ &\quad \times Q_0^{\epsilon,\tau_1}(\mu_{1,\dots,p_1}, \nu_{1,\dots,q_1}) Q_0^{\epsilon,\tau_1}(\mu_{(p_1+1),\dots,p}, \nu_{(q_1+1),\dots,q}). \end{aligned}$$

For general g, $Q_0^{\epsilon,(\tau_1,\tau_2,g)}$ is given by

$$Q_0^{\epsilon,(\tau_1,\tau_2,g)}(\mu_{1,\dots,p}, \nu_{1,\dots,q} \setminus g) = \int_{E^{2\#g}} Q_0^{\epsilon,(\tau_1,\tau_2,\emptyset)}(\mu_{1,\dots,p}, \nu_{1,\dots,q})\, d(\mu,\nu)_g,$$

where $(\mu_{1,\dots,p}, \nu_{1,\dots,q} \setminus g)$ means that variables (μ_i, ν_j) are removed for all $(i,j) \in g$ and $d(\mu,\nu)_g = \prod_{(i,j)\in g} \delta(\mu_i + \nu_{-j}) d\mu_i d\nu_j$.

For example,

$$\begin{aligned} Q_0^{\epsilon,(\Yvcentermath1\begin{smallmatrix}\cdot&\cdot\\\cdot\end{smallmatrix},\mathbb{I},\emptyset)}(\mu_{1,2,3}, \nu_1) &= \psi_\circ(k_{[12]} - l_1, k_3)\chi^\epsilon(k_1,k_2,k_3,l_1) \\ &\quad \times Q(\mu_{[12]} + \nu_{-1}) Q(\mu_1) Q(\mu_2) \overline{Q(-\nu_{-1})} Q(\mu_3), \\ Q_0^{\epsilon,(\Yvcentermath1\begin{smallmatrix}\cdot&\cdot\\\cdot\end{smallmatrix},\mathbb{I},(3,1))}(\mu_1, \mu_2) &= \int_{E^2} \psi_\circ(k_{[12]} - l_1, k_3)\chi^\epsilon(k_1,k_2,k_3,l_1) \\ &\quad \times Q(\mu_{[12]} + \nu_{-1}) Q(\mu_1) Q(\mu_2) \overline{Q(-\nu_{-1})} \\ &\quad \times Q(\mu_3)\, \delta(\mu_3 + \nu_{-1})\, d\mu_3 d\nu_1. \end{aligned}$$

Proof. Contraction formula of $H_t^{\epsilon,(\tau_1,\tau_2,g)}$ is trivial from the product formula of Wiener chaoses. For example, we see

$$\begin{aligned} X^{\epsilon,}_{(t,x)} &= X^{\epsilon,}_{(t,x)} \odot X^{\epsilon,}_{(t,x)} \\ &= \sum [\triangle_{m_1} X_t^{\epsilon,}](x)[\triangle_{m_2} X_t^{\epsilon,}](x) \\ &= \sum \mathcal{J}_{2,1}(\rho_{m_1} f^{\epsilon,}_{(t,x)}) \mathcal{J}_{1,0}(\rho_{m_2} f^{\epsilon,}_{(t,x)}) \\ &= \mathcal{J}_{3,1}(f^{\epsilon,(,,\emptyset)}_{(t,x)}) + \mathcal{J}_{2,0}(f^{\epsilon,(,,(3,1))}_{(t,x)}), \end{aligned}$$

where the summation runs over integers $m_1, m_2 \geq -1$ with $|m_1 - m_2| \leq 1$. In order to obtain contraction formula of $Q_0^{\epsilon,(\tau_1,\tau_2,g)}$, we use Plancherel's formula

$$\int_{\mathbf{R}^2} f(s)g(t)\delta(s-t)\,dsdt = \int_{\mathbf{R}^2} \hat{f}(\sigma)\hat{g}(\tau)\delta(\sigma+\tau)\,d\sigma d\tau.$$

Note that $\mu_i + \nu_{-j} = 0$ if and only if $\sigma_i + \tau_j = 0, k_i = l_j$. This formula is obtained as follows.

$$\begin{aligned} \int_{\mathbf{R}^2} f(s)g(t)\delta(s-t)\,dsdt &= \int_{\mathbf{R}} f(s)\overline{g(s)}\,ds = \int_{\mathbf{R}} \hat{f}(\sigma)\overline{\hat{\bar{g}}(\sigma)}\,d\sigma \\ &= \int_{\mathbf{R}} \hat{f}(\sigma)\hat{g}(-\sigma)\,d\sigma = \int_{\mathbf{R}^2} \hat{f}(\sigma)\hat{g}(\tau)\delta(\sigma+\tau)\,d\sigma d\tau. \end{aligned}$$

The proof is completed. □

In Table 3, we give a list of all contractions g for each (τ_1, τ_2) and define the corresponding symbols (τ_1, τ_2, g). Note that the graphs in the same line gives the same kernel $f^{\epsilon,(\tau_1,\tau_2,g)}$, so we write (τ_1, τ_2, g) by the same symbol. By taking the renormalization into account, we have the following decompositions:

(5.5)

$$\begin{aligned} X^{\epsilon,} &= \mathcal{J}_{3,1}(f^{\epsilon,}) + \mathcal{J}_{2,0}(f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{2,2}(f^{\epsilon,}) + 2\mathcal{J}_{1,1}(f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{3,1}(f^{\epsilon,}) + 2\mathcal{J}_{2,0}(f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{2,2}(f^{\epsilon,}) + 4\mathcal{J}_{1,1}(f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{2,2}(f^{\epsilon,}) + \mathcal{J}_{1,1}(f^{\epsilon,}) + \mathcal{J}_{1,1}(f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{1,3}(f^{\epsilon,}) + 2\mathcal{J}_{0,2}(f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{3,2}(f^{\epsilon,}) + 2\mathcal{J}_{2,1}(f^{\epsilon,}) + \mathcal{J}_{2,1}(f^{\epsilon,}) + 2\mathcal{J}_{1,0}(\Re f^{\epsilon,}), \\ X^{\epsilon,} &= \mathcal{J}_{2,3}(f^{\epsilon,}) + 4\mathcal{J}_{1,2}(f^{\epsilon,}) + 2\mathcal{J}_{0,1}(\Re f^{\epsilon,}) \end{aligned}$$

where

$$\mathfrak{R}f^{\epsilon,\tau} = f^{\epsilon,\tau} - \mathfrak{c}^{\epsilon}_{2,2} f^{\epsilon,\tau'}, \qquad \mathfrak{R}f^{\epsilon,\tau} = f^{\epsilon,\tau} - \mathfrak{c}^{\epsilon}_{2,1} f^{\epsilon,\tau''}.$$

Table 3: List of graphical symbols for contractions

τ_1	τ_2	g	(τ_1, τ_2, g)
		$\emptyset$	
		$\{(3,1)\}$	
		$\emptyset$	
		$\{(1,2)\},\{(2,2)\}$	
		$\emptyset$	
		$\{(1,1)\},\{(2,1)\}$	
		$\emptyset$	
		$\{(1,1)\},\{(1,2)\},\{(2,1)\},\{(2,2)\}$	
		$\{(1,1),(2,2)\},\{(1,2),(2,1)\}$	
		$\emptyset$	
		$\{(1,2)\}$	
		$\{(2,1)\}$	
		$\{(1,2),(2,1)\}$	
		$\emptyset$	
		$\{(1,2)\},\{(1,3)\}$	
		$\emptyset$	
		$\{(1,2)\},\{(2,2)\}$	
		$\{(3,1)\}$	
		$\{(1,2),(3,1)\},\{(2,2),(3,1)\}$	
		$\emptyset$	
		$\{(1,2)\},\{(1,3)\},\{(2,2)\},\{(2,3)\}$	
		$\{(1,2),(2,3)\},\{(1,3),(2,2)\}$	

Finally, we define a process X^{τ}, which is a candidate of the limit of $\{X^{\epsilon,\tau}\}_{0<\epsilon<1}$. It may be natural to define $H_t^{(\tau_1,\tau_2,g)}$ by the same way as in Proposition 5.7 by replacing H_t^{ϵ,τ_i} by $H_t^{\tau_i}$ for $i = 1, 2$. This definition makes sense if $\#g = 0, 1$, however, does not if $\#g = 2$. In Section 5.3.3, we will show that there exist kernels $\mathfrak{R}f^{\tau}$ for $\tau =$ such that

$$\mathfrak{R}f^{\epsilon,\tau} \to \mathfrak{R}f^{\tau}, \qquad \mathfrak{R}f^{\epsilon,\tau} \to \mathfrak{R}f^{\tau}$$

as $\epsilon \to 0$.

Definition 5.8. For $\tau \in \{$ $\}$, we define

$$X^{\tau}_{(t,x)} = \sum_{\#g=0,1} \mathcal{J}_{p-\#g,q-\#g}(f^{\tau}_{(t,x)}).$$

For $\tau =$, we define

$$X^{\text{}} = \mathcal{J}_{3,2}(f^{\text{}}) + 2\mathcal{J}_{2,1}(f^{\text{}}) + \mathcal{J}_{2,1}(f^{\text{}}) + 2\mathcal{J}_{1,0}(\mathfrak{R}f^{\text{}}),$$
$$X^{\text{}} = \mathcal{J}_{2,3}(f^{\text{}}) + 4\mathcal{J}_{1,2}(f^{\text{}}) + 2\mathcal{J}_{0,1}(\mathfrak{R}f^{\text{}}).$$

The following is the main theorem in this section:

Theorem 5.9. *Let $\kappa > 0$ and $T > 0$. Then, we have*

$$\lim_{\epsilon \downarrow 0} \boldsymbol{E}[\|X - X^{\epsilon}\|^{p}_{\mathcal{X}^{\kappa}_{T}}] = 0$$

for every $1 < p < \infty$.

Remark 5.10. The limit process X in Theorem 5.9 is given explicitly by generalized Itô-Wiener integrals. Since the expression of kernels are independent of χ, so is X.

The proof of this theorem will be given in the next section.

5.3 Proof of convergence of driving vectors

In this section, we show the convergence $X^{\epsilon,\tau} \to X^{\tau}$ for all τ. As stated above, they have the good kernels. Hence, it is sufficient to estimate Q^{τ}_{0} and $Q^{\tau}_{0} - Q^{\epsilon,\tau}_{0}$, due to Proposition 5.4.

5.3.1 Useful estimates

In order to estimate Q^{τ}_{0} and $Q^{\tau}_{0} - Q^{\epsilon,\tau}_{0}$, we use the following lemmas many times.

Lemma 5.11. *If $\alpha, \beta \in (0,5)$ and $\alpha + \beta > 5$, we have*

$$\int_{E} |\mu|_{*}^{-\alpha} |\nu - \mu|_{*}^{-\beta} d\mu \lesssim |\nu|_{*}^{-\alpha-\beta+5}.$$

Proof. We modify [GP17, Lemma 9.8] to the three dimensional case. □

Lemma 5.12. *The function $\psi_\circ$ defined by (5.4) is bounded and supported in the set $\{(k,l); C^{-1}|l|_* \le |k|_* \le C|l|_*\}$ for some $C > 0$. Moreover, we have*

$$|\psi_\circ(k,l)| \lesssim |k+l|_*^{-\theta}|l|_*^{\theta}$$

for every $\theta > 0$.

Proof. The properties $|\psi_\circ(k,l)| \le 1$ and $\psi_\circ(k,l) > 0$ imply $|k|_* \approx |l|_*$ are trivial. We show the last property. Since if $(k,l) \in \mathrm{supp}(\psi_\circ)$ then $|l|_*/|k|_* \gtrsim 1$ and $|k+l|_* \le |k|_* + |l|_* \lesssim |k|_*$, we have

$$|\psi_\circ(k,l)| \lesssim |k|_*^{-\theta}|l|_*^{\theta} \lesssim |k+l|_*^{-\theta}|l|_*^{\theta}$$

for every $\theta > 0$. □

5.3.2 Lower order terms

Now we consider X^τ for $\tau = \mathbf{I}, \mathbf{I}\!\!\mathbf{I}, \mathbf{V}, \mathbf{V}\!\!\mathbf{I}, \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}, \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}\!\!\mathbf{I}, \mathbf{Y}, \mathbf{Y}\!\!\mathbf{I}, \mathbf{Y}\!\!\mathbf{I}\!\!\mathbf{I}$.

Proposition 5.13. *For $\tau = \mathbf{I}, \mathbf{I}\!\!\mathbf{I}, \mathbf{V}, \mathbf{V}\!\!\mathbf{I}, \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}, \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}\!\!\mathbf{I}, \mathbf{Y}, \mathbf{Y}\!\!\mathbf{I}, \mathbf{Y}\!\!\mathbf{I}\!\!\mathbf{I}$, we have*

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^\tau(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim |\mu|_*^{-2\gamma_\tau},$$

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |(Q_0^\tau - Q_0^{\epsilon,\tau})(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim \epsilon^\lambda |\mu|_*^{-2\gamma_\tau+\lambda},$$

for every $\lambda \in (0,1]$, where $\gamma_\tau = 2\,(\tau = \mathbf{I}, \mathbf{I}\!\!\mathbf{I}), \frac{3}{2}\,(\tau = \mathbf{V}, \mathbf{V}\!\!\mathbf{I}, \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}), 1\,(\tau = \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}), \frac{7}{2}\,(\tau = \mathbf{Y}, \mathbf{Y}\!\!\mathbf{I}), 3\,(\tau = \mathbf{Y}\!\!\mathbf{I}\!\!\mathbf{I})$.

Proof. For $\tau = \mathbf{I}, \mathbf{I}\!\!\mathbf{I}$, the required estimates is trivial from $|Q(\mu)| \lesssim |\mu|_*^{-2}$. Indeed,

$$|Q_0^{\mathbf{I}}(\mu_1)|^2 = |Q(\mu_1)|^2 \lesssim |\mu_1|_*^{-4}, \qquad |Q_0^{\mathbf{I}\!\!\mathbf{I}}(\nu_1)|^2 = |Q(\nu_{-1})|^2 \lesssim |\nu_1|_*^{-4}.$$

For $\tau = \mathbf{V}$, from Lemma 5.11 we have

$$\int_{\mu_{[12]}=\mu} |Q_0^{\mathbf{V}}(\mu_{1,2})|^2 \lesssim \int_{\mu_{[12]}=\mu} |\mu_1|_*^{-4}|\mu_2|_*^{-4} \lesssim |\mu|_*^{-3}.$$

The case $\tau = \mathbf{V}\!\!\mathbf{I}, \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}$ are parallel. For $\tau = \mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}\!\!\mathbf{I}$, we have

$$\int_{\mu_{[12]}+\nu_{-1}=\mu} |Q_0^{\mathbf{V}\!\!\mathbf{I}\!\!\mathbf{I}\!\!\mathbf{I}}(\mu_{1,2},\nu_1)|^2 \lesssim \int_{\mu_{[12]}+\nu_{-1}=\mu} |\mu_1|_*^{-4}|\mu_2|_*^{-4}|\nu_{-1}|_*^{-4} \lesssim |\mu|_*^{-2}.$$

For $\tau = \mathbf{Y}, \mathbf{Y}\!\!\mathbf{I}, \mathbf{Y}\!\!\mathbf{I}\!\!\mathbf{I}$, we have

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^\tau(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2$$
$$\lesssim |\mu|_*^{-4} \int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^{\tau_0}(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2$$
$$\lesssim \begin{cases} |\mu|_*^{-7}, & \tau = \Upsilon, \Upsilon^{\bullet}, \\ |\mu|_*^{-6}, & \tau = \Upsilon^{\bullet\bullet}. \end{cases}$$

Here, we used Proposition 5.5 (2). The required estimates of $Q_0^\tau - Q_0^{\epsilon,\tau}$ is obtained by similar computations by using $Q_0^\tau - Q_0^{\epsilon,\tau} = (1-\chi^\epsilon)Q_0^\tau$ and the inequality

$$|1-\chi^\epsilon(k_{1,\dots,p},l_{1,\dots,q})| \tag{5.6}$$
$$\lesssim \epsilon^\lambda \left(\sum_{i=1}^p |k_i|^\lambda + \sum_{j=1}^q |l_j|^\lambda\right) \lesssim \epsilon^\lambda \left(\sum_{i=1}^p |\mu_i|_*^\lambda + \sum_{j=1}^q |\nu_j|_*^\lambda\right)$$

for every $\lambda \in (0,1]$. □

Proof of Theorem 5.9 *for* $\mathfrak{l}$, $\mathcal{V}$, $\mathcal{V}^{\bullet}$, Υ, $\Upsilon^{\bullet}$, $\Upsilon^{\bullet\bullet}$. Propositions 5.4 and 5.13 imply the conclusion. Note that we need to prove $X^{\Upsilon^{\bullet\bullet}}$ is Hölder continuous in time. □

5.3.3 Higher order terms

Now we consider X^τ for $\tau = \mathcal{I}_{\mathcal{V}}(\tau_1)\tau_2$, $\mathcal{I}_{\mathcal{V}}(\tau_1)\tau_2^{\bullet}$, $\mathcal{I}_{\mathcal{V}^{\bullet}}(\tau_1)\tau_2$, $\mathcal{I}_{\mathcal{V}^{\bullet}}(\tau_1)\tau_2^{\bullet}$, $\mathcal{I}_{\Upsilon}(\tau_1)\tau_2$, $\mathcal{I}_{\Upsilon}(\tau_1)\tau_2^{\bullet}$, $\mathcal{I}_{\Upsilon^{\bullet}}(\tau_1)\tau_2$, $\mathcal{I}_{\Upsilon^{\bullet}}(\tau_1)\tau_2^{\bullet}$. We define (τ_1,τ_2) for each τ as in Proposition 5.7. We note that X^τ is written as a sum of Itô-Wiener integrals which have good kernels $f^{(\tau_1,\tau_2,g)}$ for $\#g = 0,1$ and $\mathfrak{R}f^{(\tau_1,\tau_2,g)}$ for $\#g = 2$ such as (5.5). We will estimate these functions for the case $\#g = 0,1,2$ separately.

First we consider the functions $Q_0^\tau = Q_0^{(\tau_1,\tau_2,\emptyset)}$.

Proposition 5.14. *For* $\tau = \mathcal{I}_{\mathcal{V}}(\tau_1)\tau_2$, $\mathcal{I}_{\mathcal{V}}(\tau_1)\tau_2^{\bullet}$, $\mathcal{I}_{\mathcal{V}^{\bullet}}(\tau_1)\tau_2$, $\mathcal{I}_{\mathcal{V}^{\bullet}}(\tau_1)\tau_2^{\bullet}$, $\mathcal{I}_{\Upsilon}(\tau_1)\tau_2$, $\mathcal{I}_{\Upsilon}(\tau_1)\tau_2^{\bullet}$, $\mathcal{I}_{\Upsilon^{\bullet}}(\tau_1)\tau_2$, $\mathcal{I}_{\Upsilon^{\bullet}}(\tau_1)\tau_2^{\bullet}$, *we have*

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^\tau(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim |\mu|_*^{-2\gamma_\tau+\kappa}|k|_*^{-2\delta_\tau-\kappa},$$
$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |(Q_0^\tau - Q_0^{\epsilon,\tau})(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim \epsilon^\lambda |\mu|_*^{-2\gamma_\tau+\kappa+\lambda}|k|_*^{-2\delta_\tau-\kappa+\lambda},$$

for every small $\kappa > 0$ *and* $\lambda > 0$, *where*

$$(\gamma_\tau,\delta_\tau) = \begin{cases} (2,\frac{1}{2}), & \tau = \mathcal{I}_{\mathcal{V}}(\tau_1)\tau_2, \mathcal{I}_{\mathcal{V}}(\tau_1)\tau_2^{\bullet}, \\ (\frac{3}{2},1), & \tau = \mathcal{I}_{\mathcal{V}^{\bullet}}(\tau_1)\tau_2, \mathcal{I}_{\mathcal{V}^{\bullet}}(\tau_1)\tau_2^{\bullet}, \mathcal{I}_{\Upsilon}(\tau_1)\tau_2, \mathcal{I}_{\Upsilon}(\tau_1)\tau_2^{\bullet}, \\ (\frac{3}{2},\frac{1}{2}), & \tau = \mathcal{I}_{\Upsilon^{\bullet}}(\tau_1)\tau_2, \mathcal{I}_{\Upsilon^{\bullet}}(\tau_1)\tau_2^{\bullet}. \end{cases}$$

Proof. We have

$$\begin{aligned}
&\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^\tau(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \\
&\quad\lesssim \int_{\mu_1'+\mu_2'=\mu} \psi_\circ(k_1',k_2')^2 \int_{\mu_{[1\dots p_1]}+\nu_{[(-1)\dots(-q_1)]}=\mu_1'} |Q_0^{\tau_1}(\mu_{1,\dots,p_1},\nu_{1,\dots,q_1})|^2 \\
&\qquad\times \int_{\mu_{[(p_1+1)\dots p]}+\nu_{[(-q_1-1)\dots(-q)]}=\mu_1'} |Q_0^{\tau_2}(\mu_{(p_1+1),\dots,p},\nu_{(q_1+1),\dots,q})|^2 \\
&\quad\lesssim \int_{\mu_1'+\mu_2'=\mu} \psi_\circ(k_1',k_2')^2 |\mu_1'|_*^{-2\gamma_{\tau_1}} |\mu_2'|_*^{-2\gamma_{\tau_2}}.
\end{aligned}$$

We estimate them by using Proposition 5.13 and Lemma 5.12. In the case $\tau =$, Proposition 5.13 implies $(\gamma_{\tau_1},\gamma_{\tau_2}) = (3,2)$. Applying Lemma 5.12, we have

$$\begin{aligned}
\psi_\circ(k_1',k_2')^2|\mu_1'|_*^{-6}|\mu_2'|_*^{-4} &\lesssim |k_1'+k_2'|_*^{-1-\kappa}|k_1'|_*^{1+\kappa}|\mu_1'|_*^{-6}|\mu_2'|_*^{-4} \\
&\le |k|_*^{-1-\kappa}|\mu_1'|_*^{-5+\kappa}|\mu_2'|_*^{-4}.
\end{aligned}$$

for any $\mu_1'+\mu_2'=\mu$. Hence

$$\begin{aligned}
\int_{\mu_1'+\mu_2'=\mu} \psi_\circ(k_1',k_2')^2|\mu_1'|_*^{-6}|\mu_2'|_*^{-4} &\lesssim |k|_*^{-1-\kappa} \int_{\mu_1'+\mu_2'=\mu} |\mu_1|_*^{-5+\kappa}|\mu_2|_*^{-4} \\
&\lesssim |k|_*^{-1-\kappa}|\mu|_*^{-4+\kappa}.
\end{aligned}$$

For $\tau =$, , , , $(\gamma_{\tau_1},\gamma_{\tau_2}) = (\frac{7}{2},\frac{3}{2})$ implies

$$\begin{aligned}
\int_{\mu_1'+\mu_2'=\mu} \psi_\circ(k_1',k_2')^2|\mu_1'|_*^{-7}|\mu_2'|_*^{-3} &\lesssim |k|_*^{-2-\kappa} \int_{\mu_1'+\mu_2'=\mu} |\mu_1|_*^{-5+\kappa}|\mu_2|_*^{-3} \\
&\lesssim |k|_*^{-2-\kappa}|\mu|_*^{-3+\kappa}.
\end{aligned}$$

For $\tau =$, , $(\gamma_{\tau_1},\gamma_{\tau_2}) = (3,\frac{3}{2})$ implies

$$\begin{aligned}
\int_{\mu_1'+\mu_2'=\mu} \psi_\circ(k_1',k_2')^2|\mu_1'|_*^{-6}|\mu_2'|_*^{-3} &\lesssim |k|_*^{-1-\kappa} \int_{\mu_1'+\mu_2'=\mu} |\mu_1|_*^{-5+\kappa}|\mu_2|_*^{-3} \\
&\lesssim |k|_*^{-1-\kappa}|\mu|_*^{-3+\kappa}.
\end{aligned}$$

By noting (5.6), we can estimate $Q_0^\tau - Q_0^{\epsilon,\tau}$ in a similar way. □

Next we consider the functions Q_0^τ for $\tau =$, , , , , , , , , . We show three propositions; Propositions 5.15, 5.16 and 5.17.

Proposition 5.15. *For $\tau =$* , , *we have*

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^\tau(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim |\mu|_*^{-5},$$

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |(Q_0^{\tau}-Q_0^{\epsilon,\tau})(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim \epsilon^{\lambda}|\mu|_*^{-5+\lambda},$$

for every small $\lambda>0$.

Proof. We consider the case $\tau=$. Note

$$\begin{aligned} Q_0^{\tau}(\mu_{1,2}) &= \int_{E^2} \psi_\circ(k_{[12]}-l_1,k_3)Q(\mu_{[12]}+\nu_{-1}) \\ &\qquad \times Q(\mu_1)Q(\mu_2)\overline{Q(-\nu_{-1})}Q(\mu_3)\,\delta(\mu_3+\nu_{-1})\,d\mu_3 d\nu_1 \\ &= Q(\mu_1)Q(\mu_2)\int_E \psi_\circ(k_{[12]}-k_3,k_3)Q(\mu_{[12]}-\mu_3)|Q(\mu_3)|^2\,d\mu_3. \end{aligned}$$

By using the estimate $|\psi_\circ(k_{[12]}-k_3,k_3)|\le 1$, we have

$$|Q_0^{\tau}(\mu_{1,2})| \lesssim |\mu_1|_*^{-2}|\mu_2|_*^{-2}\int_E |\mu_{[12]}-\mu_3|_*^{-2}|\mu_3|_*^{-4}\,d\mu_3 \lesssim |\mu_1|_*^{-2}|\mu_2|_*^{-2}|\mu_{[12]}|_*^{-1}.$$

Hence

$$\int_{\mu_{[12]}=\mu} |Q_0^{\tau}(\mu_{1,2})|^2 \lesssim |\mu|_*^{-2}\int_{\mu_{[12]}=\mu} |\mu_1|_*^{-4}|\mu_2|_*^{-4} \lesssim |\mu|_*^{-5}.$$

We will show the second inequality for $\tau=$. The inequality (5.6) implies

$$|Q_0^{\tau}(\mu_{1,2})-Q_0^{\epsilon,\tau}(\mu_{1,2})| \lesssim |\mu_1|_*^{-2}|\mu_2|_*^{-2}\cdot\epsilon^{\lambda/2}\{|\mu_1|_*^{\lambda}+|\mu_2|_*^{\lambda}+|\mu_{[12]}|_*^{\lambda}\}^{1/2}|\mu_{[12]}|_*^{-1}.$$

Hence we obtain the second inequality for $\tau=$.

The assertion for is verified in the same way. □

Proposition 5.16. *For* $\tau=$, , , , , *we have*

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |Q_0^{\tau}(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim |\mu|_*^{-4+\kappa}|k|_*^{-1-\kappa},$$

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu} |(Q_0^{\tau}-Q_0^{\epsilon,\tau})(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2 \lesssim \epsilon^{\lambda}|\mu|_*^{-4+\kappa+\lambda}|k|_*^{-1-\kappa},$$

for every small $\kappa>0$ *and* $\lambda>0$.

Proof. We give a proof for the case $\tau=$ only, because we can show the other cases in the same way. Note

$$\begin{aligned} Q_0^{\tau}(\mu_{1,3}) &= \int_{E^2} \psi_\circ(k_{[12]},k_3-l_1)Q(\mu_{[12]}) \\ &\qquad \times Q(\mu_1)Q(\mu_2)Q(\mu_3)\overline{Q(-\nu_{-1})}\,\delta(\mu_2+\nu_{-1})\,d\mu_2 d\nu_1 \end{aligned}$$

$$= Q(\mu_1)Q(\mu_3)\int_E \psi_\circ(k_1+k_2,k_3-k_2)Q(\mu_{[12]})|Q(\mu_2)|^2\,d\mu_2.$$

Lemma 5.12 implies

$$\psi_\circ(k_1+k_2,k_3-k_2)\lesssim |k_1+k_3|_*^{-\frac{1+\kappa}{2}}|k_1+k_2|_*^{\frac{1+\kappa}{2}}\le |k_1+k_3|_*^{-\frac{1+\kappa}{2}}|\mu_{[12]}|_*^{\frac{1+\kappa}{2}}.$$

Combining them, we have

$$\begin{aligned}|Q_0^{\text{ }}(\mu_{1,3})| &\lesssim |\mu_1|_*^{-2}|\mu_3|_*^{-2}|k_1+k_3|_*^{-\frac{1+\kappa}{2}}\int_E |\mu_{[12]}|_*^{-\frac{3-\kappa}{2}}|\mu_2|_*^{-4}\,d\mu_2\\ &\lesssim |\mu_1|_*^{-2}|\mu_3|_*^{-2}|k_1+k_3|_*^{-\frac{1+\kappa}{2}}|\mu_1|_*^{-\frac{1-\kappa}{2}}\end{aligned}$$

Hence

$$\int_{\mu_{[13]}=\mu}|Q_0^{\text{}}(\mu_{1,3})|^2\lesssim |k|_*^{-1-\kappa}\int_{\mu_{[13]}=\mu}|\mu_1|_*^{-5+\kappa}|\mu_3|_*^{-4}\lesssim |k|_*^{-1-\kappa}|\mu|_*^{-4+\kappa}.$$

In a similar way, we see

$$\begin{aligned}&|Q_0^{\text{ }}(\mu_{1,3})-Q_0^{\epsilon,\text{}}(\mu_{1,3})|\\ &\quad\lesssim |\mu_1|_*^{-2}|\mu_3|_*^{-2}|k_1+k_3|_*^{-\frac{1+\kappa}{2}}\cdot\epsilon^{\lambda/2}\{|\mu_1|_*^\lambda+|\mu_2|_*^\lambda+|\mu_{[12]}|_*^\lambda\}^{1/2}|\mu_1|_*^{-\frac{1-\kappa}{2}},\end{aligned}$$

which implies the second assertion. The proof has been completed. □

Proposition 5.17. *For* $\tau=$, , , *we have*

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu}|Q_0^\tau(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2\lesssim|\mu|_*^{-4+\kappa}|k|_*^{-\kappa},$$

$$\int_{\mu_{[1\dots p]}+\nu_{[(-1)\dots(-q)]}=\mu}|(Q_0^\tau-Q_0^{\epsilon,\tau})(\mu_{1,\dots,p},\nu_{1,\dots,q})|^2\lesssim\epsilon^\lambda|\mu|_*^{-4+\kappa+\lambda}|k|_*^{-\kappa}$$

for every small $\kappa>0$ *and* $\lambda>0$.

Proof. Here, we will show the assertion for $\tau=$ only. Note

$$\begin{aligned}Q_0^{\text{ }}(\mu_{1,3},\nu_1) &= \int_{E^2}\psi_\circ(k_{[12]}-l_1,k_3-l_2)Q(\mu_{[12]}+\nu_{-1})Q(\mu_1)Q(\mu_2)\overline{Q(-\nu_{-1})}\\ &\qquad\times Q(\mu_3)\overline{Q(-\nu_{-2})}\,\delta(\mu_2+\nu_{-2})\,d\mu_2d\nu_2\\ &= Q(\mu_1)Q(\mu_3)\overline{Q(-\nu_{-1})}\\ &\qquad\times\int_E\psi_\circ(k_{[12]}-l_1,k_3-k_2)Q(\mu_{[12]}+\nu_{-1})|Q(\mu_2)|^2\,d\mu_2.\end{aligned}$$

We use Lemma 5.12 to obtain

$$\psi_\circ(k_{[12]}-l_1,k_3-k_2)\lesssim|k_{[13]}-l_1|_*^{-\frac{\kappa}{2}}|k_{[12]}-l_1|_*^{\frac{\kappa}{2}}\lesssim|k_{[13]}-l_1|_*^{-\frac{\kappa}{2}}|\mu_{[12]}+\nu_{-1}|_*^{\frac{\kappa}{2}}.$$

Hence

$$
\begin{aligned}
&|Q_0^{\tau_1}(\mu_{1,3},\nu_1)| \\
&\qquad \lesssim |\mu_1|_*^{-2}|\mu_3|_*^{-2}|\nu_{-1}|_*^{-2}|k_{[13]}-l_1|_*^{-\frac{\kappa}{2}}\int_E |\mu_{[12]}+\nu_{-1}|_*^{-2+\frac{\kappa}{2}}|\mu_2|_*^{-4}\,d\mu_2 \\
&\qquad \lesssim |\mu_1|_*^{-2}|\mu_3|_*^{-2}|\nu_{-1}|_*^{-2}|k_{[13]}-l_1|_*^{-\frac{\kappa}{2}}|\mu_1+\nu_{-1}|_*^{-1+\frac{\kappa}{2}},
\end{aligned}
$$

which implies

$$
\begin{aligned}
&\int_{\mu_{[13]}+\nu_{-1}=\mu}|Q_0^{\tau_1}(\mu_{1,3},\nu_1)|^2 \\
&\qquad\qquad \lesssim |k|_*^{-\kappa}\int_{\mu_{[13]}+\nu_{-1}=\mu}|\mu_1|_*^{-4}|\mu_3|_*^{-4}|\nu_{-1}|_*^{-4}|\mu_1+\nu_{-1}|_*^{-2+\kappa} \\
&\qquad\qquad \lesssim |k|_*^{-\kappa}|\mu|_*^{-4+\kappa}.
\end{aligned}
$$

In addition, we have

$$
\begin{aligned}
|Q_0^{\tau_1}(\mu_{1,3},\nu_1)-Q_0^{\epsilon,\tau_1}(\mu_{1,3},\nu_1)| &\lesssim |\mu_1|_*^{-2}|\mu_3|_*^{-2}|\nu_{-1}|_*^{-2}|k_{[13]}-l_1|_*^{-\frac{\kappa}{2}} \\
&\quad\times \epsilon^{\lambda/2}\{|\mu_1|_*^{\lambda}+|\mu_3|_*^{\lambda}+|\nu_{-1}|_*^{\lambda}+|\mu_1+\nu_{-1}|_*^{\lambda}\}^{1/2}|\mu_1+\nu_{-1}|_*^{-1+\frac{\kappa}{2}},
\end{aligned}
$$

which implies the conclusion. The proof is completed. □

Finally we consider the functions $\mathfrak{R}Q_0^{\tau}$ for $\tau = \tau_2, \tau_3$. First of all, we have to define the renormalized kernels $\mathfrak{R}f^{\tau}$. Since

$$
\begin{aligned}
\mathfrak{R}Q_0^{\epsilon,\tau_2} &= Q_0^{\epsilon,\tau_2}-Q_0^{\epsilon,\tau_2\tau_1}Q_0^{\epsilon,\bullet}, \\
\mathfrak{R}Q_0^{\epsilon,\tau_3} &= Q_0^{\epsilon,\tau_3}-Q_0^{\epsilon,\tau_3\tau_1}Q_0^{\epsilon,\bullet},
\end{aligned}
$$

we have

$$
\begin{aligned}
\mathfrak{R}Q_0^{\epsilon,\tau_2}(\mu_1) &= \chi^{\epsilon}(k_1)Q(\mu_1)\int_{E^2}\chi^{\epsilon}(k_{2,3})^2|Q(\mu_2)|^2|Q(\mu_3)|^2 \\
&\qquad\times \delta_{0,\mu_1}\{\psi_\circ(\cdot+k_2-k_3,k_3-k_2)Q(\cdot+\mu_2-\mu_3)\}\,d\mu_2 d\mu_3, \\
\mathfrak{R}Q_0^{\epsilon,\tau_3}(\nu_1) &= \chi^{\epsilon}(k_1)\overline{Q(-\nu_{-1})}\int_{E^2}\chi^{\epsilon}(k_{1,2})^2|Q(\mu_1)|^2|Q(\mu_2)|^2 \\
&\qquad\times \delta_{0,-\nu_{-1}}\{\psi_\circ(\cdot+k_{[12]},-k_{[12]})Q(\cdot+\mu_{[12]})\}\,d\mu_1 d\mu_2,
\end{aligned}
$$

where $\delta_{\mu_1,\mu_2}f = f(\mu_2)-f(\mu_1)$ is the difference operator. We set

$$
\mathfrak{R}Q_0^{\tau_1}(\mu_1) = Q(\mu_1)\int_{E^2}|Q(\mu_2)|^2|Q(\mu_3)|^2
$$

$$\times \delta_{0,\mu_1}\{\psi_\circ(\cdot + k_2 - k_3, k_3 - k_2)Q(\cdot + \mu_2 - \mu_3)\}\, d\mu_2 d\mu_3,$$

$$\mathfrak{R}Q_0^{\Lambda}(\nu_1) = \overline{Q(-\nu_{-1})} \int_{E^2} |Q(\mu_1)|^2 |Q(\mu_2)|^2$$
$$\times \delta_{0,-\nu_{-1}}\{\psi_\circ(\cdot + k_{[12]}, -k_{[12]})Q(\cdot + \mu_{[12]})\}\, d\mu_1 d\mu_2,$$

if they are well-defined. The required kernels $\mathfrak{R}f^\tau$ is defined by good kernels with $\mathfrak{R}H_t^\tau := \mathcal{F}_{\text{time}}^{-1}(e^{-2\pi \mathrm{i} R t} Q_0^\tau)$. The following proposition implies that these kernels are well-defined and $\mathfrak{R}f^{\epsilon,\tau}$ converges to $\mathfrak{R}f^\tau$.

Proposition 5.18. *For* $\tau = \Lambda, \Lambda$, *we have*

$$|\mathfrak{R}Q_0^\tau(\mu)|^2 \lesssim |\mu|_*^{-4+\kappa},$$
$$|(\mathfrak{R}Q_0^\tau - \mathfrak{R}Q_0^{\epsilon,\tau})(\mu)|^2 \lesssim \epsilon^\lambda |\mu|_*^{-4+\kappa+\lambda}$$

for every small $\kappa > 0$ *and* $\lambda > 0$.

In order to prove this proposition, we extend the domains of $\psi_\circ = \psi_\circ(k,l)$ and $Q = Q(\sigma, k)$ into $\mathbf{R}^6$ and $\mathbf{R}^4$ in a natural way, respectively. We write $k = (k^1, k^2, k^3) \in \mathbf{R}^3$. Then, we have the following estimate of their derivatives:

Lemma 5.19. *For every* $0 < \kappa < 1$, *it holds that*

$$|\partial_{k^\alpha}\psi_\circ(k,l)| \lesssim 1_{|k|\approx|l|}|k|_*^{-1+\kappa}, \quad |\partial_\sigma Q(\mu)| \lesssim |\mu|_*^{-4}, \quad |\partial_{k^\alpha} Q(\mu)| \lesssim |\mu|_*^{-3}.$$

Proof. The latter two inequality can be shown easily. We show the first inequality. Note that $(k,l) \in \operatorname{supp}\psi_\circ$ implies $|k| \approx |l|$. We see

$$\partial_{k^\alpha}\psi_\circ(k,l) = \sum_{|i-j|\le 1} 2^{-i}\partial_{k^\alpha}\rho_0(2^{-i}k)\rho_j(l) = \sum_{i\ge -1} 2^{-i}\partial_{k^\alpha}\rho_0(2^{-i}k) \sum_{j;|i-j|\le 1}\rho_j(l).$$

In this calculation, we abused the symbols ρ_{-1} and ρ_0. We see that the compactness of $\operatorname{supp}\rho_0$ implies $|\partial_{k^\alpha}\rho_0(2^{-i}k)| \lesssim |2^{-i}k|_*^{-1+\kappa} \le 2^{-i(1-\kappa)}|k|_*^{-1+\kappa}$ and the summation $\sum_{j;|i-j|\le 1}\rho_j(l)$ has an upper bound independent of i. Hence,

$$|\partial_{k^\alpha}\psi_\circ(k,l)| \lesssim \sum_{|i-j|\le 1} 2^{-i(2-\kappa)}|k|_*^{-1+\kappa} \lesssim |k|_*^{-1+\kappa}.$$

The proof is completed. □

Proof of Proposition 5.18. We focus on $\tau = \Lambda$.

First we esimate the difference operator part in $\mathfrak{R}Q_0^{\Lambda}$. We write $\tau^s\mu = (\tau^2\sigma, \tau k)$ for $\mu \in E$ and $\tau \in [0,1]$. The fundamental theorem of calculus and Lemma 5.19 imply

$$|\delta_{0,\mu_1}\{\psi_\circ(\cdot+k_2,-k_2)Q(\cdot+\mu_2)\}| = \left|\int_0^1 \frac{d}{d\tau}\{\psi_\circ(\cdot,-k_2)Q\}(\tau^s\mu_1+\mu_2)\,d\tau\right|$$
$$\lesssim |k_1|\int_0^1 \mathtt{1}_{|\tau k_1+k_2|\approx|k_2|}|\tau k_1+k_2|_*^{-1+\kappa}|\tau^s\mu_1+\mu_2|_*^{-2}\,d\tau$$
$$+|\sigma_1|\int_0^1 \tau|\tau^s\mu_1+\mu_2|_*^{-4}\,d\tau + |k_1|\int_0^1 |\tau^s\mu_1+\mu_2|_*^{-3}\,d\tau.$$

Hence we have

$$\int_{E^2}|Q(\mu_2)|^2|Q(\mu_3)|^2|\delta_{0,\mu_1}\{\psi_\circ(\cdot+k_2-k_3,k_3-k_2)Q(\cdot+\mu_2-\mu_3)\}|\,d\mu_2 d\mu_3$$
$$\lesssim A_1+A_2+A_3,$$

where

$$A_1 = \int_{E^2} d\mu_2 d\mu_3\,|\mu_2|_*^{-4}|\mu_3|_*^{-4}|k_1|\int_0^1 \mathtt{1}_{|\tau k_1+k_2-k_3|\approx|k_2-k_3|}$$
$$\times|\tau k_1+k_2-k_3|_*^{-1+\kappa}|\tau^s\mu_1+\mu_2-\mu_3|_*^{-2}\,d\tau,$$
$$A_2 = \int_{E^2} d\mu_2 d\mu_3\,|\mu_2|_*^{-4}|\mu_3|_*^{-4}|\sigma_1|\int_0^1 \tau|\tau^s\mu_1+\mu_2-\mu_3|_*^{-4}\,d\tau,$$
$$A_3 = \int_{E^2} d\mu_2 d\mu_3\,|\mu_2|_*^{-4}|\mu_3|_*^{-4}|k_1|\int_0^1 |\tau^s\mu_1+\mu_2-\mu_3|_*^{-3}\,d\tau.$$

We estimate the terms A_1, A_2 and A_3. Note that Lemma 5.11 holds even if $\nu\in\mathbf{R}^4$. We start the estimates with A_1. By changing variables with $\mu_2'=\mu_2$ and $\mu_3'=\mu_2-\mu_3$ and the Fubini theorem, we have

$$A_1 = |k_1|\int_{E^2} d\mu_2' d\mu_3'\,|\mu_2'|_*^{-4}|\mu_2'-\mu_3'|_*^{-3}$$
$$\times\left(\int_0^1 \mathtt{1}_{|\tau k_1+k_3'|\approx|k_3'|}|\tau k_1+k_3'|_*^{-1+\kappa}|\tau^s\mu_1+\mu_3'|_*^{-2}\,d\tau\right)$$
$$\lesssim |k_1|\int_E d\mu_3'\,|\mu_3'|^{-3}\left(\int_0^1 \mathtt{1}_{|\tau k_1+k_3'|\approx|k_3'|}|\tau k_1+k_3'|_*^{-1+\kappa}|\tau^s\mu_1+\mu_3'|_*^{-2}\,d\tau\right)$$
$$= |k_1|\int_0^1 d\tau\int_E d\mu_3'\,\mathtt{1}_{|\tau k_1+k_3'|\approx|k_3'|}|\tau k_1+k_3'|_*^{-1+\kappa}|\tau^s\mu_1+\mu_3'|_*^{-2}|\mu_3'|^{-3}.$$

The Young inequality implies

$$\int_{\mathbf{R}} d\sigma_3'\,|\tau^s\mu_1+\mu_3'|_*^{-2}|\mu_3'|_*^{-3} \le \int_{\mathbf{R}} d\sigma_3'\,\left(|\tau^s\mu_1+\mu_3'|_*^{-5}+|\mu_3'|_*^{-5}\right)$$
$$= |\tau k_1+k_3'|_*^{-3}+|k_3'|_*^{-3}.$$

Hence

$$\begin{aligned}\int_E d\mu_3' \, \mathbf{1}_{|\tau k_1+k_3'|\approx|k_3'|}|\tau k_1+k_3'|_*^{-1+\kappa}|\tau^s\mu_1+\mu_3'|_*^{-2}|\mu_3'|^{-3}\\ \lesssim \sum_{k_3'} \mathbf{1}_{|\tau k_1+k_3'|\approx|k_3'|}|\tau k_1+k_3'|_*^{-1+\kappa}\left(|\tau k_1+k_3'|_*^{-3}+|k_3'|_*^{-3}\right)\\ \lesssim \sum_{k_3'} \mathbf{1}_{|\tau k_1+k_3'|\approx|k_3'|}|k_3'|_*^{-4+\kappa} \lesssim \sum_{k_3':|\tau k_1|\lesssim|k_3'|}|k_3'|_*^{-4+\kappa} \lesssim |\tau k_1|_*^{-1+\kappa}.\end{aligned}$$

Here, we used that $|\tau k_1| \le |\tau k_1+k_3'|+|k_3'| \lesssim |k_3'|$ in the case that $|\tau k_1+k_3'| \approx |k_3'|$. Combining them and using that $|\tau^s\mu|_* \ge \tau|\mu|_*$ for every $\tau \in [0,1]$, we obtain

$$A_1 \lesssim |k_1| \int_0^1 d\tau (\tau|k_1|_*)^{-1+\kappa} \lesssim |k_1|_*^\kappa \le |\mu_1|_*^\kappa.$$

The estimate of A_1 has finished.

The estimates of A_2 and A_3 is easy. Indeed, we have

$$A_2 \lesssim |\sigma_1| \int_0^1 \tau|\tau^s\mu_1|_*^{-2}\, d\tau, \qquad A_3 \lesssim |k_1| \int_0^1 |\tau^s\mu_1|_*^{-1}\, d\tau.$$

Since $|\tau^s\mu|_* \ge \tau|\mu|_*$ for every $\tau \in [0,1]$, we have

$$\begin{aligned}A_2 &\lesssim |\sigma_1| \int_0^1 \tau|\tau^s\mu_1|_*^{\kappa-2}\, d\tau \lesssim |\sigma_1||\mu_1|_*^{\kappa-2} \int_0^1 \tau^{\kappa-1}\, d\tau \lesssim |\mu_1|_*^\kappa,\\ A_3 &\lesssim |k_1| \int_0^1 |\tau^s\mu_1|_*^{\kappa-1}\, d\tau \lesssim |k_1||\mu_1|_*^{\kappa-1} \int_0^1 \tau^{\kappa-1}\, d\tau \lesssim |\mu_1|_*^\kappa\end{aligned}$$

for every $\kappa \in (0,1)$.

Hence

$$|\Re Q_0^\tau(\mu)|^2 \lesssim |\mu|_*^{-4+\kappa}.$$

We obtain the estimate of $|(\Re Q_0^\tau - \Re Q_0^{\epsilon,\tau})(\mu)|^2$ in a similar way. We can see the assertion is valid for $\tau =$ in the same way. □

Proof of Theorem 5.9 *for* , , , , , , , . We will use Proposition 5.4. The constant $(\gamma_\tau, \delta_\tau)$ in Proposition 5.14 satisfies

$$\gamma_\tau + \delta_\tau = \begin{cases} \frac{5}{2}, & \tau = \text{}, \text{},\\ \frac{5}{2}, & \tau = \text{}, \text{ }, \text{ }, \text{},\\ 2, & \tau = \text{}, \text{}. \end{cases} \tag{5.7}$$

The assertions for the case and follow from (5.7) and Proposition 5.15. For , , and , we see the assertion from (5.7) and Proposition 5.16. For and , we use (5.7), Propositions 5.17 and 5.18. □

5.4 Properties of renormalization constants

In this subsection, we study the renormalization constants $\mathfrak{c}_1^\epsilon$, $\mathfrak{c}_{2,1}^\epsilon$ and $\mathfrak{c}_{2,2}^\epsilon$ defined by (5.3). We use Propositions 5.5 and 5.7 to show that they are independent of (t,x) and the choice of the dyadic partition $\{\rho_m\}_{m=-1}^\infty$ of unity (Proposition 5.20) and obtain the divergence rate (Proposition 5.21). Proposition 5.20 and Theorem 4.29 imply the renormalized equation (1.2) does not depend on the choice of the dyadic partition of unity. Hence the solution to (1.2) is independent of the partition.

5.4.1 Expression of renormalization constants

We obtain explicit expressions of $\mathfrak{c}_1^\epsilon$, $\mathfrak{c}_{2,1}^\epsilon$ and $\mathfrak{c}_{2,2}^\epsilon$ as follows:

Proposition 5.20. *We have the following:*

$$\mathfrak{c}_1^\epsilon = \sum_{k\in\mathbf{Z}^3} \frac{\chi^\epsilon(k)^2}{2(4\pi^2|k|^2+1)}, \tag{5.8}$$

$$\mathfrak{c}_{2,1}^\epsilon = \sum_{k_1,k_2\in\mathbf{Z}^3} \frac{\chi^\epsilon(k_1,k_2)^2}{4(4\pi^2\mu|k_1|^2+1)(4\pi^2\mu|k_2|^2+1)(\alpha_1+\mathsf{i}\beta_1)}, \tag{5.9}$$

$$\mathfrak{c}_{2,2}^\epsilon = \sum_{k_1,l_1\in\mathbf{Z}^3} \frac{\chi^\epsilon(k_1,l_1)^2}{4(4\pi^2\mu|k_1|^2+1)(4\pi^2\mu|l_1|^2+1)(\alpha_2+\mathsf{i}\beta_2)}, \tag{5.10}$$

where

$$\alpha_1 = 4\pi^2\mu(|k_1+k_2|^2+|k_1|^2+|k_2|^2)+3, \quad \beta_1 = 4\pi^2(|k_1+k_2|^2-|k_1|^2-|k_2|^2),$$
$$\alpha_2 = 4\pi^2\mu(|k_1-l_1|^2+|k_1|^2+|l_1|^2)+3, \quad \beta_2 = 4\pi^2(|k_1-l_1|^2-|k_1|^2+|l_1|^2).$$

Proof. From Proposition 5.5, we have

$$\begin{aligned}\mathfrak{c}_1^\epsilon &= \sum_{k\in\mathbf{Z}^3}\int_{\mathbf{R}} f_{(t,x)}^{\epsilon,\mathsf{I}}(s,k) f_{(t,x)}^{\epsilon,\vdots}(s,k)\,ds \\ &= \sum_{k\in\mathbf{Z}^3}\int_{\mathbf{R}} \chi^\epsilon(k)^2 |H_t(s,k)|^2\,ds = \sum_{k\in\mathbf{Z}^3}\frac{\chi^\epsilon(k)^2}{2(4\pi^2|k|^2+1)},\end{aligned}$$

which is (5.8).

We show (5.9). From Propositions 5.5 and 5.7, we have

$$\begin{aligned}\mathfrak{c}_{2,1}^\epsilon &= H_t^{\epsilon,(\curlyvee,\vdots,\{(1,1),(2,2)\})} \\ &= \int_{E^2} \psi_\circ(k_1+k_2,-(k_1+k_2)) H_t^{\epsilon,\curlyvee}(m_{1,2}) H_t^{\epsilon,\vdots}(m_{1,2})\,dm_1 dm_2.\end{aligned}$$

By using $\psi_\circ(k_1+k_2,-(k_1+k_2))=1$ and

$$\int_{\mathbf{R}} ds\, H^{\shortuparrow}_u(s,k)H^{\shortdownarrow}_t(s,k) = \frac{1}{2(4\pi^2\mu|k|^2+1)}H^{\shortdownarrow}_t(u,k),$$
$$\int_{\mathbf{R}} dt\, H^{\shortdownarrow}_u(t,l)H^{\shortuparrow}_t(t,l) = \frac{1}{2(4\pi^2\mu|l|^2+1)}H^{\shortuparrow}_t(u,l),$$

we obtain the assertion. We can show (5.10) in the same way. □

5.4.2 Divergence rate of renormalization constants

Here, we show the following proposition concerning divergence rate of the the renormalization constants:

Proposition 5.21. *There exists a positive constant C such that*

$$C^{-1}\epsilon^{-1} \le \mathfrak{c}_1^\epsilon \le C\epsilon^{-1}, \tag{5.11}$$
$$C^{-1}\log\epsilon^{-1} \le |\mathfrak{c}_{2,1}^\epsilon| \le C\log\epsilon^{-1}, \tag{5.12}$$
$$C^{-1}\log\epsilon^{-1} \le |\mathfrak{c}_{2,2}^\epsilon| \le C\log\epsilon^{-1} \tag{5.13}$$

for any $0<\epsilon<1$.

Since the estimate (5.11) follows easily from (5.8), we show (5.12) and (5.13) for the rest of this subsection.

Lemma 5.22. *Let α_1, β_1, α_2 and β_2 be as in Proposition* 5.20. *There exist positive constants C_1 and C_2 such that*

$$\Re\frac{1}{\alpha_1+\mathsf{i}\beta_1} \ge C_1(|k_1|^2+|k_2|^2+1)^{-1}, \quad \left|\frac{1}{\alpha_1+\mathsf{i}\beta_1}\right| \le C_2(|k_1|^2+|k_2|^2+1)^{-1},$$
$$\Re\frac{1}{\alpha_2+\mathsf{i}\beta_2} \ge C_1(|k_1|^2+|l_1|^2+1)^{-1}, \quad \left|\frac{1}{\alpha_2+\mathsf{i}\beta_2}\right| \le C_2(|k_1|^2+|l_1|^2+1)^{-1},$$

for any k_1, k_2 and l_1.

Proof. We show the assertion for $1/(\alpha_1+\mathsf{i}\beta_1)$. Since $|k_1|^2+|k_2|^2+1 \lesssim \alpha_1 \lesssim |k_1|^2+|k_2|^2+1$ and $|\beta_1| \lesssim |k_1|^2+|k_2|^2+1$, we see

$$\Re\frac{1}{\alpha_1+\mathsf{i}\beta_1} = \Re\frac{\alpha_1-\mathsf{i}\beta_1}{\alpha_1^2+\beta_1^2} = \frac{\alpha_1}{\alpha_1^2+\beta_1^2} \gtrsim \frac{|k_1|^2+|k_2|^2+1}{(|k_1|^2+|k_2|^2+1)^2}$$

and

$$\left|\frac{1}{\alpha_1+\mathsf{i}\beta_1}\right| = \left|\frac{\alpha_1-\mathsf{i}\beta_1}{\alpha_1^2+\beta_1^2}\right| \lesssim \frac{|k_1|^2+|k_2|^2+1}{(|k_1|^2+|k_2|^2+1)^2}.$$

The assertion is verified. We can show the assertion for $1/(\alpha_2+\mathsf{i}\beta_2)$ in the same way. □

Proof of Proposition 5.21. We first prove of the lower estimate. We show that there exists a constant C_1 such that

$$\Re \mathfrak{c}_{2,1}^{\epsilon}, \Re \mathfrak{c}_{2,2}^{\epsilon} \geq C_1 \log \epsilon^{-1},$$

for any $0 < \epsilon < 1$.

We consider only $\mathfrak{c}_{2,1}^{\epsilon}$ and estimate the real part of summands in (5.9). Proposition 5.20 and Lemma 5.22 imply

$$\begin{aligned}
\Re \mathfrak{c}_{2,1}^{\epsilon} &\gtrsim \sum_{k_1,k_2 \in \mathbf{Z}^3} \chi^{\epsilon}(k_1)^2 \chi^{\epsilon}(k_2)^2 (|k_1|^2+1)^{-1} (|k_2|^2+1)^{-1} (|k_1|^2+|k_2|^2+1)^{-1} \\
&\geq \sum_{k_1,k_2 \in \mathbf{Z}^3} \chi^{\epsilon}(k_1)^2 \chi^{\epsilon}(k_2) (|k_1|^2+|k_2|^2+1)^{-3} \\
&\geq 1 + \log \epsilon^{-1},
\end{aligned}$$

which implies the lower estimate of $\Re \mathfrak{c}_{2,1}^{\epsilon}$. We can obtain that of $\Re \mathfrak{c}_{2,2}^{\epsilon}$ by the same way.

Next we prove the upper estimate. We show that there exists a constant C_1 such that

$$|\mathfrak{c}_{2,1}^{\epsilon}|, |\mathfrak{c}_{2,2}^{\epsilon}| \leq C_2 \log \epsilon^{-1}$$

for any $0 < \epsilon < 1$.

We consider only $\mathfrak{c}_{2,1}^{\epsilon}$. Proposition 5.20 and Lemma 5.22 imply

$$\begin{aligned}
|\mathfrak{c}_{2,1}^{\epsilon}| &\lesssim \sum_{k_1,k_2 \in \mathbf{Z}^3} \chi^{\epsilon}(k_1)^2 \chi^{\epsilon}(k_2)^2 (|k_1|^2+1)^{-1} (|k_2|^2+1)^{-1} (|k_1|^2+|k_2|^2+1)^{-1} \\
&\leq \sum_{|k_1|,|k_2| \leq \epsilon^{-1}} (|k_1|^2+1)^{-1} (|k_2|^2+1)^{-1} (|k_1|^2+|k_2|^2+1)^{-1}.
\end{aligned}$$

In this estimate, we used $\operatorname{supp} \chi \subset B(0,1)$. We divide the region of the summation $\{(k_1,k_2); |k_1| \leq \epsilon^{-1} \text{ and } |k_2| \leq \epsilon^{-1}\} \subset \mathbf{Z}^3 \times \mathbf{Z}^3$ into

$$\begin{aligned}
A_1 &= \{(k_1,k_2); |k_1| \leq 2 \text{ or } |k_2| \leq 2\}, \\
A_2 &= \{(k_1,k_2); 2 \leq |k_1| \leq |k_2| \leq \epsilon^{-1}\}, \\
A_3 &= \{(k_1,k_2); 2 \leq |k_2| \leq |k_1| \leq \epsilon^{-1}\}.
\end{aligned}$$

The summation over A_1 is estimated as follows:

$$\begin{aligned}
\sum_{(k_1,k_2) \in A_1} (|k_1|^2+1)^{-1} (|k_2|^2+1)^{-1} (|k_1|^2+|k_2|^2+1)^{-1} &\lesssim \sum_{k \in \mathbf{Z}^3} (|k|^2+1)^{-2} \\
&< \infty.
\end{aligned}$$

For the summation over A_2, we have

$$\begin{aligned}
&\sum_{(k_1,k_2)\in A_2} (|k_1|^2+1)^{-1}(|k_2|^2+1)^{-1}(|k_1|^2+|k_2|^2+1)^{-1} \\
&\qquad \le \sum_{(k_1,k_2)\in A_2} |k_1|^{-2}|k_2|^{-2}|k_2|^{-2} \\
&\qquad \le \sum_{k_2:2\le|k_2|\le\epsilon^{-1}} \left(\sum_{k_1:2\le|k_1|\le|k_2|} |k_1|^{-2} \right) |k_2|^{-4} \\
&\qquad \lesssim \sum_{2\le|k_2|\le\epsilon^{-1}} |k_2||k_2|^{-4} \\
&\qquad \lesssim \log\epsilon^{-1}.
\end{aligned}$$

The summation over A_3 has the same upper bound. Hence we see $|\mathfrak{c}^\epsilon_{2,1}| \lesssim \log\epsilon^{-1}$. We can prove $|\mathfrak{c}^\epsilon_{2,2}| \lesssim \log\epsilon^{-1}$ by the same way. The proof is completed. □

A Complex multiple Itô-Wiener integral

We recall some notations and properties of complex multiple Wiener integrals from [Itô52].

A complex random variable Z is called isotropic complex normal if $\Re Z$ and $\Im Z$ are independent, has the same law with mean 0 and $(\Re Z, \Im Z)$ is jointly normal. A system of complex random variables $\{Z_\lambda\}$ is called jointly isotropic complex normal if $\sum_{i=1}^n c_i Z_{\lambda_i}$ is isotropic complex normal for any n, any $c_1,\dots,c_n \in \mathbf{C}$, and any indices $\lambda_1,\dots,\lambda_n$. Note that the isotropic complex normal $\{Z_\lambda\}$ satisfies $\boldsymbol{E}[Z_\lambda Z_\mu] = 0 = \boldsymbol{E}[\overline{Z_\mu Z_\nu}]$. The distribution of jointly isotropic complex normal system $\{Z_\lambda\}$ is uniquely determined by the positive-definite matrix $V_{\lambda\mu} = \boldsymbol{E}[Z_\lambda \overline{Z_\mu}]$ ([Itô52, Theorem 2.3]).

Let $(E, \mathcal{B}, \mathfrak{m})$ be a σ-finite, atomless measure space, and $\mathcal{B}^*$ be the set of all elements $A \in \mathcal{B}$ such that $\mathfrak{m}(A) < \infty$. Then there exists a jointly isotropic complex normal system $\{M(A)\,; A \in \mathcal{B}^*\}$ defined on a probability space $(\Omega, \mathcal{F}, \boldsymbol{P})$ such that

$$\boldsymbol{E}[M(A)\overline{M(B)}] = \mathfrak{m}(A \cap B),$$

and its distribution is uniquely determined ([Itô52, Theorem 3.1]).

Now the complex multiple Itô-Wiener integral of $f \in L^2_{p,q} = L^2(E^p \times E^q)$ is

defined as follows. Let $S_{p,q}$ be the set of $L^2_{p,q}$ functions of the form

$$f = \sum_{i_1,\dots,i_p,j_1,\dots,j_q=1}^{n} a_{i_1\dots i_p j_1\dots j_q} \mathbf{1}_{E_{i_1}\times\cdots\times E_{i_p}\times E_{j_1}\times\cdots\times E_{j_q}}$$

with $n \in \mathbf{Z}_+$, where $E_1, \dots, E_n$ are any disjoint sets of $\mathcal{B}^*$ and $\{a_{i_1\dots i_p j_1\dots j_q}\}$ is a set of complex numbers such that $= 0$ unless $i_1, \dots, i_p, j_1, \dots, j_q$ are all different. For f of this form, we define

$$\mathcal{J}_{p,q}(f) = \sum_{i_1,\dots,i_p,j_1,\dots,j_q=1}^{n} a_{i_1\dots i_p j_1\dots j_q} M(E_{i_1})\cdots M(E_{i_p})\overline{M(E_{j_1})}\cdots\overline{M(E_{j_q})}.$$

This functional has the property $\boldsymbol{E}[|\mathcal{J}_{p,q}(f)|^2] \le p!q!\|f\|^2_{L^2_{p,q}}$. We defined the Itô-Wiener integral for non-symmetric functions, hence, we cannot expect the equality in this inequality. Since $S_{p,q}$ is dense in $L^2_{p,q}$, the integral $\mathcal{J}_{p,q}$ is uniquely extended into continuous linear map from $L^2_{p,q}$ to $L^2(\boldsymbol{P})$. We set $L^2_{0,0} = \mathbf{C}$ and $\mathcal{J}_{0,0}(c) = c$. From [Itô52, Theorem 7], we have

$$\begin{aligned}\boldsymbol{E}[|\mathcal{J}_{p,q}(f)|^2] &\le p!q!\|f\|_{L^2_{p,q}},\\ \boldsymbol{E}[\mathcal{J}_{p,q}(f)\overline{\mathcal{J}_{r,s}(g)}] &= 0 \quad \text{for } (p,q) \ne (r,s).\end{aligned}$$

The product formula is important. For $0 \le r_1 \le p_1 \wedge q_2$ and $0 \le r_1 \le q_1 \wedge p_2$, we denote by $\mathcal{F}(p_1, q_1; p_2, q_2; r_1, r_2)$ the set of graphs consisting of disjoint $r_1 + r_2$ edges

$$(p', q') \in \{1, \dots, p_1\} \times \{q_1 + 1, \dots, q_1 + q_2\} \cup \{p_1 + 1, \dots, p_1 + p_2\} \times \{1, \dots, q_1\}.$$

For $(f, g) \in L^2_{p_1,q_1} \times L^2_{p_2,q_2}$ and $\gamma \in \mathcal{F}(p_1, q_1; p_1, q_2; r_1, r_2)$, we define $f \otimes_\gamma g \in L^2_{p_1+p_2-(r_1+r_2),q_1+q_2-(r_1+r_2)}$ by

$$\begin{aligned}&(f \otimes_\gamma g)(t_1, \dots, t_{p_1+p_2}, s_1, \dots, s_{q_1+q_2} \setminus \{(t_{p'}, s_{q'})\}_{(p',q')\in\gamma})\\ &\qquad = \int_{E^{r_1+r_2}} h(\{(t_{p'}, s_{q'})\}_{(p',q')\in\gamma}) \prod_{(p',q')\in\gamma} dm(t_{p'}, s_{q'}),\end{aligned}$$

where $h : E^{r_1+r_2} \to \mathbf{C}$ is defined by

$$\begin{aligned}h(\{(t_{p'}, s_{q'})\}_{(p',q')\in\gamma}) &= f(t_1, \dots, t_{p_1}, s_1, \dots, s_{q_1})\\ &\quad \times g(t_{p_1+1}, \dots, t_{p_1+p_2}, s_{q_1+1}, \dots, s_{q_1+q_2})|_{t_{p'}=s_{q'},(p',q')\in\gamma}.\end{aligned}$$

Theorem A.1 ([Itô52, Theorem 9], [Nua06, Proposition 1.1.2])**.** *For every $f \in L^2_{p_1,q_1}$ and $g \in L^2_{p_2,q_2}$, we have*

$$\mathcal{J}_{p_1,q_1}(f)\mathcal{J}_{p_2,q_2}(g)$$
$$= \sum_{r_1=0}^{p_1\wedge q_2} \sum_{r_2=0}^{p_2\wedge q_1} \sum_{\gamma\in\mathcal{F}(p_1,q_1;p_2,q_2;r_1,r_2)} \mathcal{J}_{p_1+p_2-(r_1+r_2),q_1+q_2-(r_1+r_2)}(f\otimes_\gamma g).$$

For example, we have

$$\begin{aligned}\mathcal{J}_{2,1}(f)\mathcal{J}_{0,2}(g) = {} & \mathcal{J}_{2,3}(f\otimes_\emptyset g)\\ & + \mathcal{J}_{1,2}(f\otimes_{\{(1,1)\}} g) + \mathcal{J}_{1,2}(f\otimes_{\{(1,2)\}} g)\\ & + \mathcal{J}_{1,2}(f\otimes_{\{(2,2)\}} g) + \mathcal{J}_{1,2}(f\otimes_{\{(2,3)\}} g)\\ & + \mathcal{J}_{0,1}(f\otimes_{\{(1,2),(2,3)\}} g) + \mathcal{J}_{0,1}(f\otimes_{\{(1,3),(2,2)\}} g).\end{aligned}$$

Acknowledgement

The authors thank Professor Reika Fukuizumi of Tohoku University for stimulating discussions and valuable advice. They also thank Professor Makoto Katori of Chuo University for helpful comments.

References

[AK02] Igor S. Aranson and Lorenz Kramer. The world of the complex ginzburg-landau equation. *Rev. Mod. Phys.*, 74:99–143, Feb 2002.

[BBF15] Ismaël Bailleul, Frédéric Bernicot, and Dorothee Frey. Higher order paracontrolled calculus, 3d-PAM and multiplicative Burgers equations, 2015, arXiv:1506.08773.

[BCD11] Hajer Bahouri, Jean-Yves Chemin, and Raphaël Danchin. *Fourier analysis and nonlinear partial differential equations*, volume 343 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2011.

[BK16] Nils Berglund and Christian Kuehn. Regularity structures and renormalisation of FitzHugh-Nagumo SPDEs in three space dimensions. *Electron. J. Probab.*, 21:Paper No. 18, 1–48, 2016.

[BS04a] Marc Barton-Smith. Global solution for a stochastic Ginzburg-Landau equation with multiplicative noise. *Stochastic Anal. Appl.*, 22(1):1–18, 2004.

[BS04b] Marc Barton-Smith. Invariant measure for the stochastic Ginzburg Landau equation. *NoDEA Nonlinear Differential Equations Appl.*, 11(1):29–52, 2004.

[CC13] Rémi Catellier and Khalil Chouk. Paracontrolled distributions and the 3-dimensional stochastic quantization equation, 2013, arXiv:1310.6869.

[FH14] Peter K. Friz and Martin Hairer. *A course on rough paths*. Universitext. Springer, Cham, 2014.

[FH16] Tadahisa Funaki and Masato Hoshino. A coupled KPZ equation, its two types of approximations and existence of global solutions, 2016, arXiv:1611.00498.

[GIP15] Massimiliano Gubinelli, Peter Imkeller, and Nicolas Perkowski. Paracontrolled distributions and singular PDEs. *Forum Math. Pi*, 3:e6, 75, 2015.

[GP17] Massimiliano Gubinelli and Nicolas Perkowski. KPZ Reloaded. *Comm. Math. Phys.*, 349(1):165–269, 2017.

[Hai02] Martin Hairer. Exponential mixing properties of stochastic PDEs through asymptotic coupling. *Probab. Theory Related Fields*, 124(3):345–380, 2002.

[Hai14] Martin Hairer. A theory of regularity structures. *Invent. Math.*, 198(2):269–504, 2014.

[Hos16a] Masato Hoshino. KPZ equation with fractional derivatives of white noise. *Stoch. Partial Differ. Equ. Anal. Comput.*, 4(4):827–890, 2016.

[Hos16b] Masato Hoshino. Paracontrolled calculus and Funaki-Quastel approximation for the KPZ equation, 2016, arXiv:1605.02624.

[Itô52] Kiyosi Itô. Complex multiple Wiener integral. *Jap. J. Math.*, 22:63–86, 1952.

[KS04] Sergei Kuksin and Armen Shirikyan. Randomly forced CGL equation: stationary measures and the inviscid limit. *J. Phys. A*, 37(12):3805–3822, 2004.

[Kup16] Antti Kupiainen. Renormalization group and stochastic PDEs. *Ann. Henri Poincaré*, 17(3):497–535, 2016.

[MW16] Jean-Christophe Mourrat and Hendrik Weber. Global well-posedness of the dynamic ϕ^4_3 model on the torus. 2016, arXiv:1605.02624.

[Nua06] David Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006.

[Oda06] Cyril Odasso. Ergodicity for the stochastic complex Ginzburg-Landau equations. *Ann. Inst. H. Poincaré Probab. Statist.*, 42(4):417–454, 2006.

[PG11] Xueke Pu and Boling Guo. Momentum estimates and ergodicity for the 3D stochastic cubic Ginzburg-Landau equation with degenerate noise. *J. Differential Equations*, 251(7):1747–1777, 2011.

[Yan04] Desheng Yang. The asymptotic behavior of the stochastic Ginzburg-Landau equation with multiplicative noise. *J. Math. Phys.*, 45(11):4064–4076, 2004.

[ZZ15] Rongchan Zhu and Xiangchan Zhu. Three-dimensional Navier-Stokes equations driven by space-time white noise. *J. Differential Equations*, 259(9):4443–4508, 2015.

Masato Hoshino

Graduate School of Mathematical Sciences
The University of Tokyo
Komaba, Meguro-ku, Tokyo, 153-8914
Japan

E-mail address: hoshino@ms.u-tokyo.ac.jp

Yuzuru Inahama

Graduate School of Mathematics
Kyushu University
Motooka, Nishi-ku, Fukuoka, 819-0395
Japan

E-mail address: inahama@math.kyushu-u.ac.jp

Nobuaki Naganuma

Graduate School of Engineering Science
Osaka University
Machikaneyama, Toyonaka, Osaka, 560-8531
Japan

E-mail address: naganuma@sigmath.es.osaka-u.ac.jp